

Volume 11: Five-Membered Heterenes with One Chalcogen and One Additional Heteroatom

	Preface	V
	Volume Editor's Preface	VII
	Table of Contents	XI
	Introduction	
	E. Schaumann	1
11.1	Product Class 1: 1,3-Dioxolium Salts and Annulated Analogues	
	H. Perst	13
11.2	Product Class 2: 1,2-Oxatholium Salts	
	H. Perst	31
11.3	Product Class 3: 1,3-Oxatholium Salts and Related Compounds	
	H. Perst and C. Klenke	35
11.4	Product Class 4: 1,2-Oxaselenolium Salts	
	H. Perst	89
11.5	Product Class 5: 1,3-Oxaselenolium Salts	
	H. Perst	93
11.6	Product Class 6: 1,2-Oxatellurolium Compounds and Benzoannulated Analogues	
	H. Perst	97
11.7	Product Class 7: 1,2-Dithiolium Salts and Related Compounds	
	C. Th. Pedersen	107
11.8	Product Class 8: 1,3-Dithiolium Salts, Se and Te Analogues, and Benzoannulated Analogues	
	G. Schukat and E. Fanghänel	191
11.9	Product Class 9: Isoxazoles	
	B. J. Wakefield	229
11.10	Product Class 10: 1,2-Benzisoxazoles and Related Compounds	
	R. K. Smalley	289
11.11	Product Class 11: 2,1-Benzisoxazoles and Related Compounds	
	R. K. Smalley	337
11.12	Product Class 12: Oxazoles	
	G. V. Boyd	383
11.13	Product Class 13: Benzoxazoles and Other Annulated Oxazoles	
	G. V. Boyd	481

11.14	Product Class 14: 1,3-Oxaphospholes and Benzoannulated Analogues D. Gudat	493
11.15	Product Class 15: Isothiazoles D. W. Brown and M. Sainsbury	507
11.16	Product Class 16: Benzisothiazoles D. W. Brown and M. Sainsbury	573
11.17	Product Class 17: Thiazoles D. Kikelj and U. Urleb	627
11.18	Product Class 18: Benzothiazoles and Related Compounds H. Ulrich	835
11.19	Product Class 19: 1,2- and 1,3-Thiaphospholes and Benzoannulated Analogues D. Gudat	913
11.20	Product Class 20: Isoselenazoles W.-D. Pfeiffer	921
11.21	Product Class 21: Annulated Isoselenazole Compounds W.-D. Pfeiffer	931
11.22	Product Class 22: Selenazoles W.-D. Pfeiffer	941
11.23	Product Class 23: Annulated Selenazole Compounds W.-D. Pfeiffer	991
11.24	Product Class 24: 1,3-Selenaphospholes D. Gudat	1001
11.25	Product Class 25: Isotellurazoles, and Annulated Isotellurazole and Tellurazole Compounds W.-D. Pfeiffer	1005
	Keyword Index	1021
	Author Index	1093
	Abbreviations	1155

Table of Contents

Introduction

E. Schaumann

Introduction	1
---------------------------	---

11.1 Product Class 1: 1,3-Dioxolium Salts and Annulated Analogues

H. Perst

11.1	Product Class 1: 1,3-Dioxolium Salts and Annulated Analogues	13
11.1.1	Product Subclass 1: Monocyclic 1,3-Dioxolium Salts	15
11.1.1.1	Synthesis by Ring-Closure Reactions	15
11.1.1.1.1	By Formation of Two O—C Bonds	15
11.1.1.1.1.1	Method 1: Reaction of α -Diazo Ketones with <i>N</i> -Carbonylmethaniminium Salts	15
11.1.1.1.1.2	Method 2: Reaction of α -Diazo Ketones with Acyl Trifluoromethanesulfonates	16
11.1.1.2	Synthesis by Substituent Modification	18
11.1.1.2.1	Addition Reactions	18
11.1.1.2.1.1	Method 1: Addition of Electrophiles to 1,3-Dioxol-2-one	18
11.1.1.2.1.2	Method 2: Addition of Electrophiles to 1,3-Dioxole-2-thione	18
11.1.1.2.2	Substitution of Existing Substituents	19
11.1.1.2.2.1	Method 1: Elimination of a Methoxy Group from a 2-Methoxy-1,3-dioxole	19
11.1.2	Product Subclass 2: Monocyclic Mesoionic 1,3-Dioxolium-4-olates	20
11.1.2.1	Synthesis by Ring-Closure Reactions	20
11.1.2.1.1	By Formation of One O—C Bond	20
11.1.2.1.1.1	Method 1: Catalytic Decomposition of Aryldiazoacetic Benzoic Anhydrides	20
11.1.2.1.1.2	Method 2: Thermal Dehydration of <i>O</i> -Acetylmandelic Acid	22
11.1.3	Product Subclass 3: Annulated 1,3-Dioxolium Salts	23
11.1.3.1	Synthesis by Ring-Closure Reactions	24
11.1.3.1.1	By Annulation to an Arene	24
11.1.3.1.1.1	By Formation of Two O—C Bonds	24
11.1.3.1.1.1.1	Method 1: From Catechol and Carbenium Salts	24
11.1.3.2	Synthesis by Substituent Modification	25
11.1.3.2.1	Addition Reactions	25
11.1.3.2.1.1	Method 1: Addition of Electrophiles to 1,3-Benzodioxol-2-one	25

11.1.3.2.2	Substitution of Existing Substituents	25
11.1.3.2.2.1	Method 1: Elimination of Chloride from 2-Chloro-1,3-benzodioxoles	25
11.1.3.2.2.1.1	Variation 1: Ionization of 2,2-Dichloro-1,3-benzodioxole	25
11.1.3.2.2.1.2	Variation 2: Ionization of 2,2-Dichloro-1,3-benzodioxole in the Presence of Nucleophiles	26
11.1.3.2.2.2	Method 2: Elimination of a Methoxy Group from 2-Methoxy-1,3-benzodioxoles and Related Systems	27
11.2	Product Class 2: 1,2-Oxathiolium Salts H. Perst	
<hr/>		
11.2	Product Class 2: 1,2-Oxathiolium Salts	31
11.2.1	Synthesis by Ring-Closure Reactions	31
11.2.1.1	By Formation of One O—S Bond	31
11.2.1.1.1	Method 1: Oxidation of a Monothio β -Diketone	31
11.2.2	Synthesis by Substituent Modification	31
11.2.2.1	Addition Reactions	31
11.2.2.1.1	Method 1: Protonation of 1,6-Dioxa-6a-thiapentalenes	31
11.3	Product Class 3: 1,3-Oxathiolium Salts and Related Compounds H. Perst and C. Klenke	
<hr/>		
11.3	Product Class 3: 1,3-Oxathiolium Salts and Related Compounds	35
11.3.1	Product Subclass 1: Monocyclic 1,3-Oxathiolium Salts	39
11.3.1.1	Synthesis by Ring-Closure Reactions	41
11.3.1.1.1	By Formation of One O—C and One S—C Bond	41
11.3.1.1.1.1	Method 1: From Ketones and N-Substituted S-Chloroisothiocarbamoyl Chlorides	41
11.3.1.1.1.2	Method 2: From α -Halo Ketones and Thiocarboxamide Derivatives	43
11.3.1.1.1.2.1	Variation 1: From α -Bromo Ketones and Tetramethylthiourea (or S-Alkyl Dithiocarbamates)	43
11.3.1.1.1.2.2	Variation 2: From α -Bromo Ketones and Arenethiocarboxamides	45
11.3.1.1.1.2.3	Variation 3: From α -Bromo Ketones and Alkanethiocarboxamides in the Presence of Aryl (or Cinnamyl) Aldehydes	47
11.3.1.1.1.3	Method 3: From α -Bromo Ketones and S-Alkyl Dithiocarboxylates	48
11.3.1.1.2	By Formation of One O—C Bond	49
11.3.1.1.2.1	Method 1: Acid-Catalyzed Cyclization of 2-Oxo-2-phenylethyl Thiocyanates	49
11.3.1.1.2.2	Method 2: Cyclocondensations of Thiocarbamate (or Dithiocarboxylate) Derivatives	50
11.3.1.1.2.2.1	Variation 1: Via Substitution of an Oxygen Substituent: From S-(2-Oxoalkyl) Thiocarbamates (or Thiocarboxylates)	51

11.3.1.1.2.2.2	Variation 2: Via Substitution of a Sulfur Substituent: From <i>S</i> -(2-Oxoalkyl) Dithiocarbamates	54
11.3.1.1.2.2.3	Variation 3: Via Substitution of an Amino Substituent: From <i>S</i> -(2-Oxoalkyl) Thiocarboximidium Salts in the Presence of Aldehydes	56
11.3.1.1.3	By Formation of One S—C Bond	57
11.3.1.1.3.1	Method 1: Cyclocondensation of $\{[(\text{Methylsulfanyl})\text{carbonothioyl}]\text{oxy}\}$ (phenyl)acetic Acid	57
11.3.1.1.3.2	Method 2: Cyclocondensation of a 2-Oxoalkyl Benzoate in the Presence of Hydrogen Sulfide	58
11.3.1.2	Synthesis by Substituent Modification	59
11.3.1.2.1	Substitution of Existing Substituents	59
11.3.1.2.1.1	Method 1: Elimination of a Methoxy Group from 2-Methoxy-1,3-oxathioles	59
11.3.2	Product Subclass 2: Monocyclic Mesoionic 1,3-Oxatholium-4-olates and -5-olates	59
11.3.2.1	Synthesis by Ring-Closure Reactions	61
11.3.2.1.1	By Formation of One O—C Bond	61
11.3.2.1.1.1	1,3-Oxatholium-5-olates	61
11.3.2.1.1.1.1	Method 1: Cyclocondensations of 2-(Aroylsulfanyl)acetic Acids	61
11.3.2.1.2	By Formation of One S—C Bond	62
11.3.2.1.2.1	1,3-Oxatholium-4-olates	62
11.3.2.1.2.1.1	Method 1: Cyclocondensation of Substituted Acetic Acids	62
11.3.2.1.2.1.1.1	Variation 1: Cyclocondensation of $\{[(\text{Dialkylamino})\text{carbonothioyl}]\text{oxy}\}$ acetic Acids	63
11.3.2.1.2.1.1.2	Variation 2: Cyclocondensation of $\{[(\text{Alkylsulfanyl})\text{carbonothioyl}]\text{oxy}\}$ acetic Acids	63
11.3.2.2	Synthesis by Substituent Modification	65
11.3.2.2.1	Substitution of Existing Substituents	65
11.3.2.2.1.1	Method 1: Substitution of a Methylsulfanyl Group in 2-(Methylsulfanyl)-1,3-oxathiol-1-ium-4-olate	65
11.3.2.2.1.2	Method 2: Substitution of the 4-Olate by a 4-Thiolate Group in 2-(Dialkylamino)-1,3-oxatholium-4-olates	66
11.3.3	Product Subclass 3: Annulated 1,3-Oxatholium Salts	66
11.3.3.1	Synthesis by Ring-Closure Reactions	69
11.3.3.1.1	By Annulation to an Arene	70
11.3.3.1.1.1	By Formation of One O—C and One S—C Bond	71
11.3.3.1.1.1.1	Method 1: From 2-Sulfanylphenol and Carboxylic Acid Derivatives or Equivalents	71
11.3.3.1.1.1.1.1	Variation 1: From 2-Sulfanylphenol and Dichloro- <i>N,N</i> -dimethylmethaniminium Chloride	71
11.3.3.1.1.1.1.2	Variation 2: From 2-Sulfanylphenol and Trihalomethyl- (or Triethoxymethyl)-Substituted Arenes	71
11.3.3.1.1.1.1.3	Variation 3: From 2-Sulfanylphenol and Carboxylic Acids or Derivatives	72

11.3.3.1.1.1.2	Method 2:	From 1,4-Quinones and Thiocarboxylic Acid (or Thiourea) Derivatives	75
11.3.3.1.1.1.2.1	Variation 1:	From 1,4-Quinones and Thioureas	76
11.3.3.1.1.1.2.2	Variation 2:	From 1,4-Quinones and Thiocarboxylic Acids or 4-(Arylthiioyl)morpholines	77
11.3.3.1.1.1.2.3	Variation 3:	From 1,4-Quinones and 4-(2-Phenylethanethiioyl)morpholine in the Presence of Aryl Aldehydes	78
11.3.3.1.1.2	By Formation of One O—C Bond		80
11.3.3.1.1.2.1	Method 1:	From Cyclocondensations of <i>S</i> -(2,5-Dihydroxyaryl) Thioesters or Thiocarbamate Derivatives	80
11.3.3.1.1.3	By Formation of One S—C Bond		82
11.3.3.1.1.3.1	Method 1:	Rearrangement of <i>O</i> -(2,4,6-Trihalophenyl) <i>N,N</i> -Dimethylthiocarbamates	82
11.3.3.2	Synthesis by Substituent Modification		83
11.3.3.2.1	Addition Reactions		83
11.3.3.2.1.1	Method 1:	<i>N</i> -Protonation of a 1,3-Benzoxathiol-2-imine	83
11.3.3.2.2	Substitution of Existing Substituents		84
11.3.3.2.2.1	Method 1:	Elimination of Hydride from a 1,3-Benzoxathiole	84
11.3.3.2.2.2	Method 2:	Elimination of an Ethoxy Group from a 2-Ethoxy-1,3-benzoxathiole	85
11.3.3.2.3	Modification of Substituents		86
11.4	Product Class 4: 1,2-Oxaselenolium Salts		
	H. Perst		
11.4	Product Class 4: 1,2-Oxaselenolium Salts		89
11.4.1	Synthesis by Ring-Closure Reactions		89
11.4.1.1	By Formation of One O—Se Bond		89
11.4.1.1.1	Method 1:	Rearrangement of <i>Z</i> -3-(Arylselanyl)propenoyl Chlorides	89
11.4.2	Synthesis by Substituent Modification		91
11.4.2.1	Addition Reactions		91
11.4.2.1.1	Method 1:	Protonation of 1,6-Dioxa-6a λ^4 -selenapentalenes	91
11.5	Product Class 5: 1,3-Oxaselenolium Salts		
	H. Perst		
11.5	Product Class 5: 1,3-Oxaselenolium Salts		93
11.5.1	Synthesis by Ring-Closure Reactions		93
11.5.1.1	By Formation of One O—C and One Se—C Bond		93
11.5.1.1.1	Method 1:	Cyclocondensation of α -Bromo Ketones with Tetramethylselenourea	93

11.5.1.1.1.1	Variation 1: Cyclocondensation of α -Bromo Ketones with N,N-Disubstituted Selenoamides	94
11.5.2	Synthesis by Substituent Modification	94
11.5.2.1	Addition Reactions	94
11.5.2.1.1	Method 1: N-Protonation of 1,3-Oxaselenol-2-imines	94
11.6	Product Class 6: 1,2-Oxatellurolium Compounds and Benzoannulated Analogues H. Perst	
11.6	Product Class 6: 1,2-Oxatellurolium Compounds and Benzoannulated Analogues	97
11.6.1	Product Subclass 1: Monocyclic 1,2-Oxatellurolium Compounds	97
11.6.1.1	Synthesis by Ring-Closure Reactions	98
11.6.1.1.1	By Formation of One O—Te Bond	98
11.6.1.1.1.1	Method 1: Rearrangement of Z-3-(Aryltellanyl)propenoyl Chlorides	98
11.6.1.2	Synthesis by Substituent Modification	101
11.6.1.2.1	Substitution of Existing Substituents	101
11.6.1.2.1.1	Method 1: Substitution of Chloride by Other Anions	101
11.6.2	Product Subclass 2: 2,1-Benzoxatellurolium Compounds	101
11.6.2.1	Synthesis by Ring-Closure Reactions	102
11.6.2.1.1	By Annulation to an Arene	103
11.6.2.1.1.1	Method 1: Rearrangement of 2-(Aryltellanyl)benzoyl Chlorides	103
11.6.2.2	Synthesis by Ring Transformation	103
11.6.2.2.1	Hetero-Ring Size Retained Overall	103
11.6.2.2.1.1	Method 1: By Oxidation of Preformed 3 <i>H</i> -2,1-Benzoxatelluroles	103
11.6.2.2.2	By Ring Contraction	104
11.6.2.2.2.1	Method 1: By Rearrangement of 1,1-Dihalo-4 <i>H</i> -1-benzotellurin-4-ones	104
11.7	Product Class 7: 1,2-Dithiolium Salts and Related Compounds C. Th. Pedersen	
11.7	Product Class 7: 1,2-Dithiolium Salts and Related Compounds	107
11.7.1	Product Subclass 1: Simple 1,2-Dithiolium Salts	107
11.7.1.1	Synthesis by Ring-Closure Reactions	109
11.7.1.1.1	By Formation of One S—S and Two S—C Bonds	109
11.7.1.1.1.1	Method 1: From 1,3-Diketones, 1,3-Dialdehydes, or 3-Oxoaldehydes	109
11.7.1.1.1.2	Method 2: From Polyhalo-Substituted Propenes	110
11.7.1.1.2	By Formation of One S—S and One S—C Bond	110

11.7.1.1.2.1	Method 1: From 3-Oxothiocarboxylic Acid Derivatives	110
11.7.1.1.2.2	Method 2: From β -Thioxo Ketones	111
11.7.1.1.2.3	Method 3: From α,β -Unsaturated β -Amino Ketones	111
11.7.1.1.3	By Formation of Two S—C Bonds	111
11.7.1.1.3.1	Method 1: From 1,3-Dioxo Compounds and Disulfane or Polysulfanes	111
11.7.1.1.3.2	Method 2: From 1,3-Dioxo Compounds and Diacetyl Disulfide	113
11.7.1.1.4	By Formation of One S—S Bond	113
11.7.1.2	Aromatization	114
11.7.1.2.1	Method 1: From 3-Alkylidene-3 <i>H</i> -1,2-dithioles	114
11.7.1.2.2	Method 2: By Oxidation of 3 <i>H</i> -1,2-Dithiole-3-thiones	114
11.7.1.2.3	Method 3: From 3 <i>H</i> -1,2-Dithiol-3-ones or 3 <i>H</i> -1,2-Dithiole-3-thiones by Halogenation	116
11.7.1.3	Synthesis by Substituent Modification	117
11.7.1.3.1	Substitution of Existing Substituents	117
11.7.1.3.1.1	Of Hydrogen	117
11.7.1.3.1.2	Of Halogen or Alkylsulfanyl Groups	117
11.7.1.3.2	Addition Reactions	119
11.7.1.3.2.1	Protonation	119
11.7.1.3.3	Modification of Substituents	120
11.7.1.3.3.1	Alkylation of 3 <i>H</i> -1,2-Dithiol-3-ones and 3 <i>H</i> -1,2-Dithiole-3-thiones	120
11.7.1.3.3.1.1	Method 1: 3 <i>H</i> -1,2-Dithiol-3-ones	120
11.7.1.3.3.1.2	Method 2: 3 <i>H</i> -1,2-Dithiole-3-thiones	121
11.7.1.3.3.2	Reactions at the α -Methyl Group	122
11.7.1.3.3.3	Exchange of Anions	125
11.7.2	Product Subclass 2: Mesoionic and Zwitterionic 1,2-Ditholium Salts	125
11.7.2.1	1,2-Ditholium-4-olates	125
11.7.2.1.1	Synthesis by Ring-Closure Reactions	126
11.7.2.1.1.1	By Formation of One S—S and Two S—C Bonds	126
11.7.2.1.1.1.1	Method 1: From Bromo-Substituted Acetones	126
11.7.2.1.1.1.2	Method 2: From Propane-1,2,3-triones	126
11.7.2.1.1.1.3	Method 3: From 1,3-Dichloroacetone	126
11.7.2.1.2	Aromatization	127
11.7.2.1.2.1	Method 1: From 4-Hydroxy-3 <i>H</i> -1,2-dithiole-3-thiones	127
11.7.2.1.2.2	Method 2: From 4-Hydroxy-1,2-ditholium Salts	127
11.7.2.1.2.3	Method 3: From 4-Alkoxy- or 4-Acyloxy-1,2-ditholium Salts	127
11.7.2.1.3	Synthesis by Substituent Modification	127
11.7.2.2	4-(1,2-Ditholium-3-yl)phenolates and Analogous Compounds	128
11.7.2.2.1	Aromatization	128
11.7.2.2.2	Synthesis by Substituent Modification	129

11.7.3	Product Subclass 3: Annulated 1,2-Dithiolium Salts	130
11.7.3.1	Synthesis by Ring-Closure Reactions	130
11.7.3.1.1	By Formation of Two S—C Bonds	130
11.7.3.1.2	By Formation of One C—C Bond	131
11.7.3.2	Synthesis by Ring Transformation	132
11.7.3.3	Aromatization	132
11.7.3.4	Synthesis by Substituent Modification	133
11.7.3.4.1	Substitution of Existing Substituents	133
11.7.3.4.2	Modification of Substituents	134
11.7.4	Product Subclass 4: Heterapentalenes Related to the 1,2-Dithiolium System	134
11.7.4.1	1,6-Dioxa-6a λ^4 -thiapentalenes	135
11.7.4.1.1	Synthesis by Ring-Closure Reactions	135
11.7.4.1.2	Synthesis by Ring Transformation	136
11.7.4.1.3	Synthesis by Substituent Modification	136
11.7.4.1.3.1	Substitution of Existing Substituents	136
11.7.4.1.3.1.1	Of Hydrogen	136
11.7.4.1.3.1.2	Of Halogen	137
11.7.4.2	1-Oxa-6,6a λ^4 -dithiapentalenes	137
11.7.4.2.1	Synthesis by Ring-Closure Reactions	138
11.7.4.2.1.1	By Formation of One O—S, One S—S, and Two S—C Bonds	138
11.7.4.2.1.2	By Formation of One O—S and One O—C Bond	138
11.7.4.2.1.3	By Formation of One O—S and One C—C Bond	139
11.7.4.2.1.3.1	Method 1: From 1,2-Dithiolium Salts	139
11.7.4.2.1.3.2	Method 2: From 3-[(2-Aryl-2-oxoethyl)sulfanyl]-1,2-dithiolium Salts	142
11.7.4.2.1.3.3	Method 3: From 3 <i>H</i> -1,2-Dithiol-3-ones or 3 <i>H</i> -1,2-Dithiole-3-thiones	143
11.7.4.2.1.4	By Formation of One O—S Bond	143
11.7.4.2.2	Synthesis by Ring Transformation	143
11.7.4.2.2.1	Method 1: From 4 <i>H</i> -Pyran- or 4 <i>H</i> -Thiopyran-4-thiones	143
11.7.4.2.2.2	Method 2: From Other Heterapentalenes	144
11.7.4.2.3	Synthesis by Substituent Modification	147
11.7.4.2.3.1	Substitution of Existing Substituents	147
11.7.4.2.3.2	Modification of Substituents	147
11.7.4.3	1,6-Dioxa-6a λ^4 -selenapentalenes	147
11.7.4.3.1	Synthesis by Ring Transformation	148
11.7.4.4	1,6-Dioxa-6a λ^4 -tellurapentalenes	148
11.7.4.4.1	Synthesis by Ring-Closure Reactions	148

11.7.4.5	1,6,6a λ^4 -Trithiapentalenes	149
11.7.4.5.1	Synthesis by Ring-Closure Reactions	150
11.7.4.5.1.1	By Formation of Two S—S, Three S—C, and One C—C Bond	150
11.7.4.5.1.2	By Formation of Two S—S and Three S—C Bonds	150
11.7.4.5.1.3	By Formation of Two S—S and Two S—C Bonds	151
11.7.4.5.1.4	By Formation of Two S—S and One S—C Bond	152
11.7.4.5.1.5	By Formation of One S—S, One S—C, and One C—C Bond	152
11.7.4.5.1.5.1	Method 1: From Dialkylamino-Substituted 1,2-Dithiolium Salts	152
11.7.4.5.1.5.2	Method 2: From 3 <i>H</i> -1,2-Dithiole-3-thiones	153
11.7.4.5.1.6	By Formation of One S—S and One S—C Bond	153
11.7.4.5.1.6.1	Method 1: From 3-[2-(Dialkylamino)vinyl]-1,2-dithiolium Salts	153
11.7.4.5.1.6.2	Method 2: From 2-(5-Aryl-3 <i>H</i> -1,2-dithiol-3-ylidene)malononitriles	154
11.7.4.5.1.7	By Formation of One S—S and One C—C Bond	155
11.7.4.5.1.7.1	Method 1: From 1,2-Dithiolium Salts	155
11.7.4.5.1.7.2	Method 2: From 3-(Alkylsulfanyl)-1,2-dithiolium Salts	155
11.7.4.5.1.7.3	Method 3: From 3 <i>H</i> -1,2-Dithiole-3-thiones	156
11.7.4.5.1.8	By Formation of One S—S Bond	157
11.7.4.5.1.8.1	Method 1: From Other 1,2-Dithiole Derivatives	157
11.7.4.5.2	Synthesis by Ring Transformation	158
11.7.4.5.2.1	Method 1: From 4 <i>H</i> -Thiopyran-4-thiones	158
11.7.4.5.2.2	Method 2: From 1-Oxa-6,6a λ^4 -dithiapentalenes	158
11.7.4.5.3	Synthesis by Substituent Modification	160
11.7.4.5.3.1	Substitution of Existing Substituents	160
11.7.4.5.3.1.1	Of Hydrogen	160
11.7.4.5.3.1.1.1	Method 1: Substitution by Bromination	160
11.7.4.5.3.1.1.2	Method 2: Substitution by Nitration or Nitrosation	160
11.7.4.5.3.1.1.3	Method 3: Substitution by Formylation	161
11.7.4.5.3.1.2	Of Heteroatoms	161
11.7.4.5.3.2	Modification of Substituents	161
11.7.4.5.3.2.1	Method 1: Condensation in the Side Chain by Aldehydes	161
11.7.4.5.3.2.2	Method 2: Condensation in the Side Chain by Carbon Disulfide and Thioformamides	162
11.7.4.6	1,6a λ^4 -Dithia-6-selenapentalenes	162
11.7.4.6.1	Synthesis by Ring-Closure Reactions	162
11.7.4.6.2	Synthesis by Ring Transformation	163
11.7.4.7	1,6a λ^4 -Dithia-6-azapentalenes	163
11.7.4.7.1	Synthesis by Ring-Closure Reactions	163
11.7.4.7.1.1	By Formation of One S—N and One N—C Bond	163

11.7.4.7.1.1.1	Method 1: From 3-[2-(Methylsulfanyl)vinyl]-1,2-dithiolium Salts	163
11.7.4.7.1.1.2	Method 2: From 3-[2-(Dimethylamino)vinyl]-1,2-dithiolium Salts	164
11.7.4.7.1.2	By Formation of One S—N and One C—C Bond	164
11.7.4.7.1.2.1	Method 1: From 5-Alkyl-3 <i>H</i> -1,2-dithiole-3-thiones and Isocyanates or Isothiocyanates	164
11.7.4.7.1.2.2	Method 2: From 5-Aryl-3-benzylidene-3 <i>H</i> -1,2-dithioles or 3-Methyl-1,2-dithiolium Salts and Isocyanides	165
11.7.4.7.2	Synthesis by Ring Transformation	165
11.7.4.8	1,6-Dithia-6a λ^4 -selenapentalenes	165
11.7.4.8.1	Synthesis by Ring Transformation	166
11.7.4.9	1-Oxa-6,6a λ^4 -dithia-2-azapentalenes	166
11.7.4.9.1	Synthesis by Ring-Closure Reactions	167
11.7.4.9.2	Synthesis by Ring Transformation	167
11.7.4.10	1-Oxa-6,6a λ^4 -dithia-3-azapentalenes	168
11.7.4.10.1	Synthesis by Ring-Closure Reactions	169
11.7.4.10.1.1	By Formation of One O—S, One S—S, and One N—C Bond	169
11.7.4.10.1.2	By Formation of One O—S and One N—C Bond	169
11.7.4.10.1.2.1	Method 1: From 3-Amino-1,2-dithiolium Salts	169
11.7.4.10.1.2.2	Method 2: From 3 <i>H</i> -1,2-Dithiole-3-thiones	169
11.7.4.10.2	Synthesis by Ring Transformation	170
11.7.4.10.2.1	Method 1: From 1,2-Benzothiazoles	170
11.7.4.10.2.2	Method 2: From 1,6,6a λ^4 -Trithia-3-azapentalenes	170
11.7.4.11	1,6,6a λ^4 -Trithia-3-azapentalenes	170
11.7.4.11.1	Synthesis by Ring-Closure Reactions	171
11.7.4.11.1.1	By Formation of One S—S and One N—C Bond	171
11.7.4.11.1.1.1	Method 1: From 3-Amino-1,2-dithiolium Salts	171
11.7.4.11.2	Synthesis by Ring Transformation	171
11.7.4.12	6,6a λ^4 -Dithia-1,2-diazapentalenes	171
11.7.4.12.1	Synthesis by Ring-Closure Reactions	172
11.7.4.12.1.1	By Formation of One S—N and One N—C Bond	172
11.7.4.12.1.1.1	Method 1: From 1,2-Dithiolium Salts	172
11.7.4.12.1.1.2	Method 2: From 3 <i>H</i> -1,2-Dithiole-3-thiones	172
11.7.4.12.2	Synthesis by Ring Transformation	173
11.7.5	Product Subclass 5: Extended Structures Based on the 1,2-Dithiolium System	173
11.7.5.1	3-[(3 <i>H</i> -1,2-Dithiol-3-ylidene)methyl]-1,2-dithiolium Salts	174
11.7.5.1.1	Synthesis by Ring-Closure Reactions	174
11.7.5.1.1.1	Method 1: From 1,2-Dithiolium Salts	174

11.7.5.1.1.2	Method 2: From 3 <i>H</i> -1,2-Dithiole-3-thiones	175
11.7.5.2	2-[(3 <i>H</i> -1,2-Dithiol-3-ylidene)methyl]-1,6,6aλ ⁴ -trithiapentalenes and Related Systems	175
11.7.5.2.1	2-[(3 <i>H</i> -1,2-Dithiol-3-ylidene)methyl]-1-oxa-6,6aλ ⁴ -dithiapentalenes	176
11.7.5.2.1.1	Synthesis by Ring-Closure Reactions	177
11.7.5.2.1.1.1	Method 1: From 1,2-Dithiolium Salts	177
11.7.5.2.1.1.2	Method 2: From 3 <i>H</i> -1,2-Dithiole-3-thiones	177
11.7.5.2.2	2-[(3 <i>H</i> -1,2-Dithiol-3-ylidene)methyl]-1,6,6aλ ⁴ -trithiapentalenes	178
11.7.5.2.2.1	Synthesis by Ring Transformation	178
11.7.6	Product Subclass 6: 1,2-Diselenolium Salts and Related Compounds	179
11.7.6.1	1,2-Diselenolium Salts	179
11.7.6.1.1	Synthesis by Ring-Closure Reactions	179
11.7.6.1.1.1	By Formation of One Se—Se and Two Se—C Bonds	179
11.7.6.1.1.1.1	Method 1: From 1,3-Diketones, 1,3-Dialdehydes, or 3-Oxoaldehydes	179
11.7.6.1.1.1.2	Method 2: From a 1,3-Dichloropropenylium Chloride	180
11.7.6.1.1.2	By Formation of One Se—Se Bond	180
11.7.6.1.2	Aromatization	180
11.7.6.2	1,6,6aλ ⁴ -Triselenapentalenes	181
11.7.6.2.1	Synthesis by Ring-Closure Reactions	181
11.7.6.3	1,2-Ditellurolium Salts	181
11.7.6.3.1	Synthesis by Ring Transformation	182
11.8	Product Class 8: 1,3-Dithiolium Salts, Se and Te Analogues and Benzoannulated Analogues G. Schukat and E. Fanghänel	
11.8	Product Class 8: 1,3-Dithiolium Salts, Se and Te Analogues and Benzoannulated Analogues	191
11.8.1	Product Subclass 1: 1,3-(Benzo)Dithiolium Salts	191
11.8.1.1	Synthesis by Ring-Closure Reactions	193
11.8.1.1.1	By Formation of Two S—C Bonds	193
11.8.1.1.1.1	Fragments S—C—C—S and C	193
11.8.1.1.1.1.1	Method 1: From 1,2-Dithiols	193
11.8.1.1.1.1.1.1	Variation 1: Reaction with Carboxylic Acid Derivatives	193
11.8.1.1.1.1.1.2	Variation 2: Reaction with Aldehydes and Oxidation of the Intermediate	194
11.8.1.1.1.2	Fragments S—C—S and C—C	195
11.8.1.1.1.2.1	Method 1: From Thioacids or Dithioacids and α-Substituted Ketones	195
11.8.1.1.2	By Formation of One S—C Bond	197
11.8.1.1.2.1	Fragment S—C—S—C—C	197

11.8.1.1.2.1.1	Method 1:	From <i>S</i> -(2-Oxoalkyl) Thiocarboxylates	197
11.8.1.1.2.1.2	Method 2:	From <i>S</i> -(2-Oxoalkyl) Dithiocarboxylates	197
11.8.1.1.2.1.3	Method 3:	From <i>S</i> -(2-Carboxyalkyl) Dithiocarboxylates	199
11.8.1.1.2.1.4	Method 4:	From <i>S</i> -(1-Cyanoalkyl) Dithiocarboxylates	201
11.8.1.1.2.1.5	Method 5:	From <i>N,N</i> -Disubstituted Dithiocarbamates	201
11.8.1.1.2.1.5.1	Variation 1:	From 2-Oxoalkyl <i>N,N</i> -Disubstituted Dithiocarbamates	201
11.8.1.1.2.1.5.2	Variation 2:	From Alkenyl or Allyl <i>N,N</i> -Dialkyldithiocarbamates	203
11.8.1.2	Synthesis by Ring Transformation		204
11.8.1.2.1	Method 1:	From 3-(Methylsulfanyl)-1,4,2-dithiazines	204
11.8.1.2.2	Method 2:	Protonation of 1,3-Dithiol-2-ylidenes	204
11.8.1.2.3	Method 3:	From 1,3-Dithioles	205
11.8.1.2.3.1	Variation 1:	Abstraction of a Hydride Ion	205
11.8.1.2.3.2	Variation 2:	Elimination of Alkanolate or Alkanethiolate Ions or Amines	206
11.8.1.2.4	Method 4:	Oxidation of 1,3-Dithiole-2-thiones with Hydroperoxides	207
11.8.1.2.5	Method 5:	Oxidation of 2-(1,3-Dithiol-2-ylidene)-1,3-dithioles	208
11.8.1.2.6	Method 6:	Halogenation of 1,3-Dithiole-2-thiones	209
11.8.1.2.7	Method 7:	Alkylation of 1,3-Dithiole-2-thiones	210
11.8.1.2.8	Method 8:	Arylation of 1,3-Dithiole-2-thiones or -selones	211
11.8.1.2.9	Method 9:	Reaction of 2-Alkyl-1,3-dithiolium Salts with Aldehydes	212
11.8.1.2.10	Method 10:	Reaction of 2-(Alkylsulfanyl)-1,3-dithiolium Salts with Amines	213
11.8.1.2.11	Method 11:	Reaction of 1,3-Dithiolium Salts with Azides	213
11.8.1.2.12	Method 12:	Reaction of 2-Chloro-1,3-dithiolium Salts with Arenesulfinates	214
11.8.2	Product Subclass 2: 1,3-(Benzo)Thiaselenolium Salts		215
11.8.2.1	Synthesis by Ring-Closure Reactions		216
11.8.2.1.1	By Formation of One Se—C Bond		216
11.8.2.1.1.1	Fragment Se—C—S—C—C		216
11.8.2.1.1.1.1	Method 1:	From 2-Oxoalkyl <i>N,N</i> -Dimethylselenothiocarbamates	216
11.8.2.2	Synthesis by Ring Transformation		216
11.8.2.2.1	Method 1:	From 2-(Ethylsulfanyl)-1,4,3-thiaselenazines	216
11.8.2.2.2	Method 2:	Elimination of Alkanethiolate Ions from 1,3-Thiaselenoles	217
11.8.2.2.3	Method 3:	Alkylation of 1,3-Thiaselenole-2-thiones	217
11.8.2.2.4	Method 4:	Reaction of 2-(Alkylsulfanyl)-1,3-thiaselenolium Salts with Amines	218
11.8.3	Product Subclass 3: 1,3-(Benzo)Diselenolium Salts		218
11.8.3.1	Synthesis by Ring-Closure Reactions		219
11.8.3.1.1	By Formation of One Se—C Bond		219
11.8.3.1.1.1	Fragment Se—C—Se—C—C		219
11.8.3.1.1.1.1	Method 1:	From 2-Oxoalkyl <i>N,N</i> -Dialkyldiselenocarbamates	219
11.8.3.2	Synthesis by Ring Transformation		220
11.8.3.2.1	Method 1:	From 1,3-Diselenoles	220
11.8.3.2.1.1	Variation 1:	Abstraction of a Hydride Ion	220
11.8.3.2.1.2	Variation 2:	Elimination of Alkaneselenolate Ions	221

11.8.3.2.2	Method 2: Alkylation of 1,3-Diselenole-2-thiones or -2-selones	221
11.8.4	Product Subclass 4: 1,3-(Benzo)Ditellurolium Salts	222
11.8.4.1	Synthesis by Ring Transformation	222
11.8.4.1.1	Method 1: Elimination of Alkanolate Ions from 1,3-Ditelluroles	222
11.8.5	Product Subclass 5: 1,3-(Benzo)Thiatellurolium and 1,3-(Benzo)Selenatellurolium Salts	223
11.9	Product Class 9: Isoxazoles B. J. Wakefield	
11.9	Product Class 9: Isoxazoles	229
11.9.1	Synthesis by Ring-Closure Reactions	231
11.9.1.1	By Formation of Two Heteroatom—Carbon Bonds	231
11.9.1.1.1	Fragments C—C—C and N—O	231
11.9.1.1.1.1	Method 1: By Reactions of Hydroxylamine with 1,3-Dialdehydes and Their Acetals	231
11.9.1.1.1.2	Method 2: By Reactions of Hydroxylamine with 3-Oxoaldehydes	232
11.9.1.1.1.3	Method 3: By Reactions of Hydroxylamine with 1,3-Diketones	233
11.9.1.1.1.4	Method 4: Synthesis of Hydroxyisoxazoles and Isoxazolones by Reactions of Hydroxylamine with 3-Oxo Esters and 1,3-Diesters	234
11.9.1.1.1.5	Method 5: Synthesis of Isoxazolamines by Reactions of Hydroxylamine with 3-Oxonitriles	236
11.9.1.1.1.6	Method 6: Synthesis of Isoxazolamines by Reactions of Hydroxylamine with 2-Cyano Esters	238
11.9.1.1.1.6.1	Variation 1: In the Absence of Excess Base	238
11.9.1.1.1.6.2	Variation 2: In the Presence of Excess Base	239
11.9.1.1.1.7	Method 7: By Reactions of Hydroxylamine with Miscellaneous Analogues of 1,3-Dicarbonyl Compounds	239
11.9.1.1.1.8	Method 8: By Reactions of Hydroxylamine with α,β -Acetylenic Carbonyl and Related Compounds	239
11.9.1.1.1.9	Method 9: Synthesis of Isoxazolamines by Reactions of Hydroxylamine with α,β -Acetylenic Nitriles	241
11.9.1.1.1.10	Method 10: By Reactions of Hydroxylamine with α,β -Unsaturated Carbonyl Compounds	242
11.9.1.1.1.10.1	Variation 1: Reactions of β -Amino α,β -Unsaturated Carbonyl Compounds with Hydroxylamine	243
11.9.1.1.1.10.2	Variation 2: Reactions of [2-(Alkylsulfanyl)alk-1-enyl] Ketones with Hydroxylamine	244
11.9.1.1.1.11	Method 11: By Reactions of Hydroxylamine with α,β -Unsaturated Nitriles	245
11.9.1.1.1.12	Method 12: By Reactions of Hydroxylamine with α,β -Dihalo Ketones, Esters, and Nitriles	245
11.9.1.1.1.13	Method 13: Use of the Nitrosonium Ion as the N—O Component	246
11.9.1.1.1.14	Method 14: Synthesis of Nitroisoxazoles from Propargyl Bromides and Sodium Nitrite	247
11.9.1.2	By Formation of One Heteroatom—Carbon and One C—C Bond	248

11.9.1.2.1	Fragments O—N—C and C—C	248
11.9.1.2.1.1	Method 1: From Nitrile Oxides and Alkynes	249
11.9.1.2.1.2	Method 2: By Cycloaddition of Nitrile Oxides to C=C Bonds, Accompanied by Elimination or Dehydrogenation	251
11.9.1.2.1.2.1	Variation 1: Reactions with Enamines	252
11.9.1.2.1.2.2	Variation 2: Reactions with Enols and Enolates	253
11.9.1.2.2	Fragments O—N—C—C and C	254
11.9.1.2.2.1	Method 1: From Dianions Derived from Oximes	254
11.9.1.2.2.1.1	Variation 1: Reactions of Oxime-Derived Dianions with Esters	254
11.9.1.2.2.1.2	Variation 2: Reactions of Oxime-Derived Dianions with Amides	255
11.9.1.2.2.2	Method 2: Synthesis of Isoxazol-5-amines from α -Halooximes and Isocyanides	256
11.9.1.3	By Formation of One Heteroatom—Carbon Bond	256
11.9.1.3.1	Fragment O—N—C—C—C	256
11.9.1.3.1.1	Method 1: Cyclization of β -Hydroxyimino Carbonyl Compounds	256
11.9.1.3.1.1.1	Variation 1: Isoxazol-3-ols by Cyclization of Hydroxamic Acids Derived from Protected β -Oxo Esters	256
11.9.1.3.1.2	Method 2: Oxidative Cyclization of α,β -Unsaturated Oximes	257
11.9.1.3.1.3	Method 3: Reductive Cyclization of 2-Nitroalkenyl Ketones and 2-Nitroalkenyl Nitriles	258
11.9.2	Synthesis by Ring Transformation	259
11.9.2.1	Ring Expansion	259
11.9.2.1.1	Method 1: From 2-Acyl-2 <i>H</i> -azirines	259
11.9.2.2	Ring Transformation	260
11.9.2.3	Ring Contraction	260
11.9.2.3.1	Method 1: (2-Hydroxyaryl)isoxazoles from Chromones	260
11.9.3	Aromatization	261
11.9.3.1	Method 1: Dehydrogenation of Dihydroisoxazoles	261
11.9.3.1.1	Variation 1: Dehydrogenation with <i>N</i> -Bromosuccinimide and Base	261
11.9.3.1.2	Variation 2: Dehydrogenation with γ -Manganese(IV) Oxide	262
11.9.3.2	Method 2: Elimination of HX from Dihydroisoxazoles	262
11.9.3.3	Method 3: Cycloelimination from Dihydroisoxazoles	263
11.9.3.4	Method 4: C=C Bond Migration	263
11.9.4	Synthesis by Substituent Modification	264
11.9.4.1	Addition Reactions	264
11.9.4.1.1	Protonation	264
11.9.4.1.2	N-Alkylation	265
11.9.4.1.2.1	Method 1: Direct N-Alkylation	265
11.9.4.1.3	Addition of Heteroatoms	266
11.9.4.2	Substitution of Existing Substituents	266
11.9.4.2.1	Of Hydrogen	266

11.9.4.2.1.1	Deuteration	266
11.9.4.2.1.2	Metalation	267
11.9.4.2.1.3	By Carbofunctional Groups	267
11.9.4.2.1.3.1	Method 1: Acylation of Isoxazol-5(4 <i>H</i>)-ones	267
11.9.4.2.1.4	Halogenation	267
11.9.4.2.1.5	By Sulfur Functionalities	268
11.9.4.2.1.5.1	Method 1: Chlorosulfonation and Sulfonation	268
11.9.4.2.1.5.2	Method 2: Thiocyanation	269
11.9.4.2.1.6	By Nitrogen Functionalities	269
11.9.4.2.1.6.1	Method 1: Nitration	269
11.9.4.2.1.6.2	Method 2: Diazo Coupling	270
11.9.4.2.2	Of Metals	271
11.9.4.2.2.1	By Hydrogen and by Metals	271
11.9.4.2.2.2	By Halogens	271
11.9.4.2.2.2.1	Method 1: Reaction of Metal Derivatives of Isoxazoles with Halogens	271
11.9.4.2.2.3	By Carbofunctional Groups	271
11.9.4.2.2.3.1	Method 1: Reactions of Isoxazol-4-yllithium Compounds	271
11.9.4.2.3	Of Carbon Functionalities	272
11.9.4.2.4	Of Halogens	272
11.9.4.2.4.1	By Metals	272
11.9.4.2.4.2	By Oxygen Functionalities	272
11.9.4.2.4.2.1	Method 1: Reaction of Haloisoxazoles with Alkoxides	272
11.9.4.2.4.3	By Sulfur Functionalities	273
11.9.4.2.4.3.1	Method 1: Reaction of 5-Chloroisoxazoles with Alkylthiolates	273
11.9.4.2.4.4	By Nitrogen Functionalities	274
11.9.4.2.4.4.1	Method 1: Reaction of 5-Chloroisoxazoles with Amines and Lithium Amides	274
11.9.4.2.4.4.2	Method 2: Reaction of a 5-Chloroisoxazole with Sodium Azide	274
11.9.4.2.4.5	By Carbofunctional Groups	275
11.9.4.2.4.5.1	Method 1: By Palladium-Catalyzed Cross Coupling	275
11.9.4.2.5	Of Oxygen Functionalities	275
11.9.4.2.5.1	By Halogens	275
11.9.4.2.5.1.1	Method 1: Reactions of Isoxazol-5(4 <i>H</i>)-ones with Phosphoryl Halides	275
11.9.4.2.5.2	By Oxygen Functionalities	276
11.9.4.2.5.3	By Nitrogen Functionalities	276
11.9.4.2.5.4	By Carbon Functionalities	276

11.9.4.2.5.4.1	Method 1: Reaction of <i>N</i> -(Tetrahydropyran-2-yl)isoxazol-5(2 <i>H</i>)-ones with Organolithium Compounds	276
11.9.4.2.6	Of Nitrogen Functionalities	276
11.9.4.2.6.1	By Halogens	276
11.9.4.2.6.2	By Oxygen Functionalities	277
11.9.4.3	Modification of Substituents	277
11.9.4.3.1	Reduction of Substituents	277
11.9.4.3.1.1	Reduction of Carbonyl Compounds	277
11.9.4.3.1.1.1	Method 1: Reduction of Isoxazolecarboxylic Acids via Mixed Anhydrides	277
11.9.4.3.1.2	Reduction of Nitroisoxazoles	278
11.9.4.3.2	Reactions Involving α -Deprotonation of Side Chains	278
11.9.4.3.2.1	Method 1: Condensation of 5-Methylisoxazoles with Aldehydes	278
11.9.4.3.2.2	Method 2: Generation of (Isoxazolylmethyl)lithium Compounds	279
11.9.4.3.2.3	Method 3: Reactions Involving Deprotonation of Isoxazolium Salts	280
11.9.4.3.3	Preparation and Reactions of Halomethylisoxazoles	280
11.9.4.3.3.1	Method 1: α -Halogenation of Methylisoxazoles	281
11.9.4.4	Rearrangement of Substituents	281
11.10	Product Class 10: 1,2-Benzisoxazoles and Related Compounds R. K. Smalley	
11.10	Product Class 10: 1,2-Benzisoxazoles and Related Compounds	289
11.10.1	Synthesis by Ring-Closure Reactions	291
11.10.1.1	By Annulation to an Arene or Hetarene	291
11.10.1.1.1	By Formation of One O—C and One N—C Bond	291
11.10.1.1.2	By Formation of One O—C and One C—C Bond	292
11.10.1.1.3	By Formation of One O—N Bond	292
11.10.1.1.3.1	Method 1: By Thermal and Base-Catalyzed Cyclization of 2-Hydroxyaryl Ketoxime and Aldoxime <i>O</i> -Acylates	292
11.10.1.1.3.2	Method 2: By Cyclization of 2-Hydroxyaryl Aldoxime <i>O</i> -Sulfonates	295
11.10.1.1.3.3	Method 3: By Cyclization of 2-Hydroxyaryl Aldoxime <i>O</i> -Carbamates	296
11.10.1.1.3.4	Method 4: By Thermal Cyclization of <i>N</i> -Hydroxy-2-hydroxyarenecarboximidamide Derivatives	297
11.10.1.1.3.5	Method 5: By Intramolecular Dehydration of 2-Hydroxyaryl Ketoximes and Aldoximes	298
11.10.1.1.3.6	Method 6: By Cyclization of <i>N</i> ,2-Dihydroxybenzamides	299
11.10.1.1.3.7	Method 7: By Oxidative Cyclization of 2-Hydroxyaryl Ketoximes	300
11.10.1.1.4	By Formation of One O—C Bond	300
11.10.1.1.4.1	Method 1: By Cyclization of 2-Haloaryl Ketoximes	301
11.10.1.1.4.2	Method 2: By Cyclization of <i>N</i> -Hydroxy-2-fluoroarenecarboximidamides	304
11.10.1.1.4.3	Method 3: By Cyclization of 2-Nitroaryl Ketoximes	305

11.10.1.1.4.4	Method 4: By Cyclization of 2-Methoxy-, 2-Diazo-, 2-Amino-, and 2-(4-Tosyloxy)aryl Ketoximes	306
11.10.1.1.5	By Formation of One N—C Bond	307
11.10.1.1.5.1	Method 1: By Intramolecular Transoximation of 2-[(Isopropylideneamino)oxy]diaryl Ketones	308
11.10.1.1.5.2	Method 2: By Cyclization of 2-[(Isopropylideneamino)oxy]benzoxonitriles	308
11.10.1.2	By Annulation to the Isoxazole Ring	309
11.10.1.2.1	Annulation of an Arene Ring	309
11.10.1.2.2	Annulation of a Five-Membered Heteroarene	310
11.10.1.2.2.1	Method 1: To a Disubstituted Isoxazole	310
11.10.1.2.2.2	Method 2: To a Monosubstituted Isoxazole	311
11.10.1.2.3	Annulation of a Six-Membered Heteroarene	311
11.10.1.2.3.1	Method 1: Isoxazolopyridines	311
11.10.1.2.3.2	Method 2: Isoxazolodiazines	313
11.10.2	Synthesis by Ring Transformation	315
11.10.2.1	Hetero-Ring Size Retained Overall	315
11.10.2.1.1	Method 1: By Decomposition of 1,3,2,4-Dioxathiazole 2-Oxides	315
11.10.2.1.2	Method 2: By Decomposition of 1,4,2-Dioxazol-5-one	316
11.10.2.2	With Ring Contraction of 4-Hydroxy-2 <i>H</i> -benzopyran-2-ones	317
11.10.3	Aromatization	317
11.10.3.1	Method 1: Aromatization of Hydro-1,2-benzisoxazoles	317
11.10.4	Synthesis by Substituent Modification	319
11.10.4.1	Addition Reactions: Protonation and N-Alkylation	319
11.10.4.2	Substitution of Existing Substituents	320
11.10.4.2.1	Of Hydrogen	320
11.10.4.2.1.1	Method 1: Metalation	320
11.10.4.2.1.2	Method 2: Acylation and Alkylation	321
11.10.4.2.1.3	Method 3: Halogenation	321
11.10.4.2.1.4	Method 4: Sulfonation	323
11.10.4.2.1.5	Method 5: Nitration	324
11.10.4.2.2	Of Metals	326
11.10.4.2.3	Of Heteroatoms	326
11.10.4.3	Modification of Substituents	327
11.10.4.4	Rearrangement of Substituents	330
11.11	Product Class 11: 2,1-Benzisoxazoles and Related Compounds R. K. Smalley	
11.11	Product Class 11: 2,1-Benzisoxazoles and Related Compounds	337
11.11.1	Synthesis by Ring-Closure Reactions	339

11.11.1.1	By Annulation to an Arene or Hetarene	339
11.11.1.1.1	By Formation of One O—C and One C—C Bond	339
11.11.1.1.1.1	Method 1: Base-Catalyzed Condensation of an Arylacetonitrile with a 1-Halo-4-nitroarene	339
11.11.1.1.1.2	Method 2: Aminoisoxazolobenzazines by Reaction of Nitrobenzazines with Potassium Cyanide in Methanol	342
11.11.1.1.2	By Formation of One O—N Bond	343
11.11.1.1.2.1	Method 1: Thermal Decomposition of 2-Azidoaryl Carbonyl Compounds	343
11.11.1.1.2.2	Method 2: Deoxygenation of 2-Nitroacylbenzenes	346
11.11.1.1.2.3	Method 3: Deoxygenation of 2-Nitrosoacylbenzenes	347
11.11.1.1.3	By Formation of One O—C Bond	347
11.11.1.1.3.1	Method 1: Reduction of 2-Nitroaryl Carbonyl Compounds	348
11.11.1.1.3.2	Method 2: Reduction of 2-Nitrobenzonitriles	351
11.11.1.1.3.3	Method 3: Cyclization of 2-Nitrobenzyl Derivatives	353
11.11.1.1.3.4	Method 4: Cyclization of 2-Nitrosoaryl Ketones	357
11.11.1.1.3.5	Method 5: From 2-(Hydroxyamino)benzonitriles	358
11.11.1.1.3.6	Method 6: Oxidation of 2-Aminoacylbenzenes	360
11.11.1.1.3.7	Method 7: Condensation of 2-Nitroaryl Aldehydes with Electron-Rich Arenes	361
11.11.1.2	By Annulation to the Isoxazole Ring	363
11.11.1.2.1	Method 1: Annulation of a Six-Membered Ring	363
11.11.2	Synthesis by Ring Transformation	364
11.11.2.1	Hetero-Ring Size Retained Overall	364
11.11.2.1.1	Method 1: Isomerization of Indole and Isoindole Derivatives	365
11.11.2.2	By Ring Contraction	366
11.11.3	Aromatization	367
11.11.4	Synthesis by Substituent Modification	367
11.11.4.1	Addition Reactions	367
11.11.4.1.1	Method 1: Protonation and Complex Formation	367
11.11.4.1.2	Method 2: N-Alkylation	368
11.11.4.2	Substitution of Existing Substituents	370
11.11.4.2.1	Method 1: Of Hydrogen	370
11.11.4.2.1.1	Variation 1: By Carbofunctional Groups	370
11.11.4.2.1.2	Variation 2: Halogenation	372
11.11.4.2.1.3	Variation 3: Methoxylation	372
11.11.4.2.1.4	Variation 4: Nitration	373
11.11.4.3	Modification of Substituents	374

11.12	Product Class 12: Oxazoles G. V. Boyd	
11.12	Product Class 12: Oxazoles	383
11.12.1	Synthesis by Ring-Closure Reactions	387
11.12.1.1	By Formation of One O—C and One N—C Bond	387
11.12.1.1.1	Fragments O—C—N and C—C	387
11.12.1.1.1.1	Method 1: From Carbonylnitrenes and Alkynes	387
11.12.1.1.2	Fragments O—C—C and C—N	388
11.12.1.1.2.1	Method 1: From α -Halo Ketones and Amides	388
11.12.1.1.2.2	Method 2: From α -Hydroxy Aldehydes or α -Hydroxy Ketones and Cyanamides	391
11.12.1.1.2.3	Method 3: From α -Hydroxy Ketones and Carbamic Acid Derivatives	393
11.12.1.1.2.4	Method 4: From α -Hydroxy Carbonyl Compounds and Thiocyanic Acid	394
11.12.1.1.2.5	Method 5: From Desyl Chloride and Alkyl or Aryl Thiocyanates	395
11.12.1.1.2.6	Method 6: By Oxidative Addition of α -Methylene Ketones to Nitriles	395
11.12.1.1.2.7	Method 7: From <i>C,N</i> -Diacylimines and Nitriles	397
11.12.1.1.2.8	Method 8: From Acylcarbenes and Nitriles	397
11.12.1.1.2.9	Method 9: From Nitriles and Alkynes	401
11.12.1.1.3	Fragments N—C—C and C—O	402
11.12.1.1.3.1	Method 1: From Vinyl Azides and Carboxylic Acids or Acyl Chlorides	402
11.12.1.1.3.2	Method 2: From Aminomalononitrile and Carboxylic Acids or Their Derivatives	403
11.12.1.1.3.3	Method 3: From <i>S,S</i> -Disubstituted Dithiooxaldiimidates and Aldehydes	404
11.12.1.1.4	Fragments O—C—C—N and C	405
11.12.1.1.4.1	Method 1: By the Fischer Synthesis: Aldehyde Cyanohydrins and Aldehydes	405
11.12.1.1.4.2	Method 2: From Aroyl Cyanides and Aldehydes	405
11.12.1.1.4.3	Method 3: From Aroylthioamides, Aldehydes, and Iodomethane	407
11.12.1.1.4.4	Method 4: Oxazole 3-Oxides from Monooximes of α -Diketones and Aldehydes	407
11.12.1.1.4.5	Method 5: From α -(Hydroxyamino)phenylacetic Acid and Aldehydes	408
11.12.1.1.4.6	Method 6: From α -Azidoacetophenones, Dimethylformamide, and Phosphoryl Chloride	409
11.12.1.1.4.7	Method 7: From α -Azido Ketones	409
11.12.1.1.4.8	Method 8: From Phenacylammonium Salts and Carbon Disulfide	410
11.12.1.2	By Formation of Two N—C Bonds	410
11.12.1.2.1	Fragments C—O—C—C and N	410
11.12.1.2.1.1	Method 1: From α -Acyloxy Ketones and an Ammonia Source	410
11.12.1.3	By Formation of One O—C and One C—C Bond	412
11.12.1.3.1	Fragments C—N—C and C—O	412
11.12.1.3.1.1	Method 1: From Isocyanides and Carbonyl Compounds	412
11.12.1.3.1.2	Method 2: From <i>N</i> -(Tosylmethyl)imines and Aldehydes	416
11.12.1.3.1.3	Method 3: From <i>N</i> -(Tosylmethyl)- <i>N'</i> -tritylcarbodiimide and Aldehydes	416

11.12.1.3.1.4	Method 4: From Dimethyl N-(Ethoxycarbonylmethyl)dithioimidocarbonate and S-Phenyl Thioesters	417
11.12.1.3.2	Fragments O—C—N—C and C	418
11.12.1.3.2.1	Method 1: From Aroyl Isocyanates and Diazomethane	418
11.12.1.3.2.2	Method 2: From <i>tert</i> -Butyl Isocyanide and an <i>N</i> -Acylimine	419
11.12.1.3.2.3	Method 3: From Tris(ethoxycarbonyl)methanimine and Dichlorocarbene ..	419
11.12.1.3.2.4	Method 4: From 4-(Acylaminomethyl)pyridinium Salts and Acetic Anhydride	419
11.12.1.4	By Formation of One O—C Bond	420
11.12.1.4.1	Fragment O—C—N—C—C	420
11.12.1.4.1.1	Method 1: Cyclization of α -Acylamino Ketones	420
11.12.1.4.1.2	Method 2: Cyclization of α -Acylamino Nitriles	423
11.12.1.4.1.3	Method 3: Cyclization of α -Acylamino Carboxylic Acids and Their Derivatives	425
11.12.1.4.1.4	Method 4: Cyclization of (Acylcarbamoyl)carbenes	433
11.12.1.4.1.5	Method 5: Cyclization of <i>N</i> -Propargylamides and <i>E</i> - β -(Acylaminomethyl)- β -iodovinyl Phenyl Sulfones	433
11.12.1.4.1.6	Method 6: Cyclization of <i>N</i> -Acyl- β -haloenamines	435
11.12.1.4.1.7	Method 7: Cyclization of <i>N</i> -Acylhexafluoropropan-2-imines	436
11.12.1.4.1.8	Method 8: Cyclization of <i>N,N'</i> -Diaroyl-1,2-bis(1 <i>H</i> -benzotriazol-1-yl)ethane- 1,2-diamines	436
11.12.1.4.2	Fragment O—C—C—N—C	437
11.12.1.4.2.1	Method 1: Cyclization of <i>N</i> -Phenacylnitrilium Salts	437
11.12.1.4.2.2	Method 2: Cyclization of Phenacylnitrones	438
11.12.1.4.2.3	Method 3: From α -Azido Ketones or Esters, Acyl Halides, and Triphenylphosphine	439
11.12.1.4.2.4	Method 4: Cyclization of Anilides of 2-Isocyanoalkanoic Acids	439
11.12.1.4.2.5	Method 5: From Derivatives of Isocyanoacetic Acid and Arylsulfanyl Chlorides	440
11.12.1.5	By Formation of One N—C Bond	441
11.12.1.5.1	Fragment N—C—C—O—C	441
11.12.1.5.1.1	Method 1: Intramolecular Aza-Wittig Reaction of β -(Acyloxy)vinyl Azides	441
11.12.1.5.1.2	Method 2: Cyclization of β -(Acyloxy)- <i>N,N</i> -bis(trimethylsilyl)enamines ...	442
11.12.2	Synthesis by Ring Transformation	443
11.12.2.1	Method 1: From 2 <i>H</i> -Azirines or Aziridines	443
11.12.2.2	Method 2: From an Azetidinone	445
11.12.2.3	Method 3: From Vinylene Carbonate and Amides	446
11.12.2.4	Method 4: From 4-Benzamido-3 <i>H</i> -1,2-dithiole-3-thione	446
11.12.2.5	Method 5: From Isoxazoles	447
11.12.2.6	Method 6: From a 1,3,2-Dioxaphosphole and Acyl Isocyanates	449
11.12.2.7	Method 7: From 1-Acyl-1 <i>H</i> -1,2,3-triazoles	449
11.12.2.8	Method 8: From a Pyridin-3(4 <i>H</i>)-one	451
11.12.3	Aromatization	451

11.12.3.1	Method 1: By Dehydrogenation of 2-Oxazolines	451
11.12.3.2	Method 2: By Decarboxylation	452
11.12.3.3	Method 3: By Retro-Diels–Alder Reactions	452
11.12.4	Synthesis by Substituent Modification	453
11.12.4.1	Substitution Reactions	453
11.12.4.1.1	Method 1: Electrophilic Substitution	453
11.12.4.1.2	Method 2: Reactions of Metalated Oxazoles with Electrophiles	456
11.12.4.1.3	Method 3: Nucleophilic Substitution	467
11.12.4.2	Elimination Reactions	469
11.12.4.2.1	Method 1: Reduction of Oxazole 3-Oxides	469
11.12.4.3	Addition Reactions	469
11.12.4.3.1	Method 1: O-Alkylation, O-Acylation and S-Alkylation	469
11.12.4.3.2	Method 2: N-Alkylation and N-Acylation	472
11.12.4.4	By Rearrangement	473
11.12.4.4.1	Method 1: The Cornforth Rearrangement	473
11.12.4.5	Modification of Substituents	473
11.12.4.5.1	Method 1: Oxidation of Oxazolylmethanols	473
11.13	Product Class 13: Benzoxazoles and Other Annulated Oxazoles G. V. Boyd	
11.13	Product Class 13: Benzoxazoles and Other Annulated Oxazoles	481
11.13.1	Synthesis by Ring-Closure Reactions	481
11.13.1.1	By Formation of One O–C and One N–C Bond	481
11.13.1.1.1	Fragments O–C–C–N and C	481
11.13.1.1.1.1	Method 1: From 2-Aminophenols and Carboxylic Acids or Acid Anhydrides	481
11.13.1.1.1.2	Method 2: From 2-Nitrophenols and Tertiary Amines	482
11.13.1.1.1.3	Method 3: From 3-Aminotropolones and Ortho Esters	482
11.13.1.1.1.4	Method 4: From Phenanthra-9,10-quinone Imines and Aldehydes	483
11.13.1.2	By Formation of One O–C Bond	484
11.13.1.2.1	Fragment O–C–N–C–C	484
11.13.1.2.1.1	Method 1: Cyclization of 1-(Carboxymethyl)pyridin-2(1H)-one	484
11.13.1.2.2	Fragment O–C–C–N–C	485
11.13.1.2.2.1	Method 1: From <i>N</i> -Acyl- and <i>O,N</i> -Diacyl-2-aminophenols and Related Compounds	485
11.13.1.2.2.2	Method 2: Oxidative Cyclization of <i>N</i> -(2-Hydroxyphenyl)thiobenzamides	487
11.13.1.2.2.3	Method 3: From 2-Aminophenols or 2-Fluoroanilines, Aryl Iodides, and Carbon Monoxide	487
11.13.1.2.2.4	Method 4: Beckmann Rearrangement of 2-Hydroxy-Substituted Acetophenone and Propiophenone Oximes	489
11.13.1.2.2.5	Method 5: Cyclization of <i>N</i> -(2-Hydroxyphenyl)amidines	490

11.13.1.3	By Formation of One N—C Bond	490
11.13.1.3.1	Fragment N—C—C—O—C	490
11.13.1.3.1.1	Method 1: Reductive Cyclization of 2-Nitrophenyl and 2-Azidophenyl Benzoate	490
11.13.2	Synthesis by Ring Transformation	491
11.13.2.1	Method 1: Benzoxazole from Benzisoxazole	491
11.13.3	Synthesis by Substituent Modification	491
11.14	Product Class 14: 1,3-Oxaphospholes and Benzoannulated Analogues D. Gudat	
<hr/>		
11.14	Product Class 14: 1,3-Oxaphospholes and Benzoannulated Analogues	493
11.14.1	Synthesis by Ring-Closure Reactions	496
11.14.1.1	By Formation of Two P—C Bonds	496
11.14.1.1.1	Fragments C—O—C—C and P	496
11.14.1.1.1.1	Method 1: From Ketene <i>O,N</i> -Acetals and Phosphorus Trihalides	496
11.14.1.2	By Formation of One P—C and One O—C Bond	496
11.14.1.2.1	Fragments P—C—C—O and C	496
11.14.1.2.1.1	Method 1: From 2-Phosphinophenols	496
11.14.1.2.1.1.1	Variation 1: Condensation with Acyl Chlorides	497
11.14.1.2.1.1.2	Variation 2: Condensation with <i>N</i> -Arylimidoyl Chlorides	497
11.14.1.2.2	Fragments P—C, C and O—C	499
11.14.1.2.2.1	Method 1: From Carbene Complexes and Phosphaalkynes	499
11.14.1.2.3	Fragments O—C—C and P—C	502
11.14.1.2.3.1	Method 1: From 2-Diazo-1,3-diketones and Phosphaalkynes	502
11.14.2	Synthesis by Ring Transformation	503
11.14.2.1	Method 1: From Phosphaalkynes and Isomünchnones	503
11.15	Product Class 15: Isothiazoles D. W. Brown and M. Sainsbury	
<hr/>		
11.15	Product Class 15: Isothiazoles	507
11.15.1	Synthesis by Ring-Closure Reactions	514
11.15.1.1	By Formation of One S—C and One N—C Bond	514
11.15.1.1.1	Method 1: By Cycloaddition of Thiazyl Chloride to a Furan or a Pyrrole	514
11.15.1.1.2	Method 2: By Addition of Thiazyl Chloride to Alkenes	515
11.15.1.1.3	Method 3: By 1,3-Dipolar Cycloaddition of Nitrile Sulfides to Alkynes and Alkenes	516
11.15.1.1.3.1	Variation 1: From 1,3,4-Oxathiazol-2-ones	516
11.15.1.1.3.2	Variation 2: From 1,3,4-Oxathiazoles	518

11.15.1.1.3.3	Variation 3:	From Difluoro- λ^4 -sulfanes	518
11.15.1.1.3.4	Variation 4:	From Thiobenzamides	521
11.15.1.2	By Formation of One S—N Bond		521
11.15.1.2.1	Method 1:	By Oxidative Ring Closure of 3-Aminopropenethioamides or 3-Aminoprop-2-enethiones	521
11.15.1.2.1.1	Variation 1:	From α -Cyano Ketones and Ammonia	522
11.15.1.2.1.2	Variation 2:	From 2,2-Bis(sulfonyl)ethene-1,1-dicarbonitriles or 2-Substituted 2-Aminoethene-1,1-dicarbonitriles and Hydrogen Sulfide	524
11.15.1.2.1.3	Variation 3:	From Acetamides and Phenyl Isothiocyanate	525
11.15.1.2.1.4	Variation 4:	From 3-Aminoprop-2-enenitriles	526
11.15.1.2.1.5	Variation 5:	From 3-Amino-1-phenylprop-2-enethiones	527
11.15.1.2.2	Method 2:	From 1,2-Dithioles and Related Compounds	528
11.15.1.2.2.1	Variation 1:	From 3 <i>H</i> -1,2-Dithioles	528
11.15.1.2.2.2	Variation 2:	From Ethene-1,1-dithiolates	529
11.15.1.2.3	Method 3:	From Bis(2-cyanoethyl) Disulfides and Related Compounds	532
11.15.1.2.4	Method 4:	From 3-Oxo- or 3-Hydrazinoprop-1-enyl Thiocyanates	534
11.15.1.2.5	Method 5:	From Propynones or 3-Chloroprop-2-enals	536
11.15.1.2.6	Method 6:	From 3-Thiocyanatoprop-2-enals	539
11.15.1.2.7	Method 7:	From Propynones and Hydroxylamine <i>O</i> -Sulfonic Acid	539
11.15.1.2.8	Method 8:	From 1-Aminopropan-2-ones, 1-Aminopropene-2-nitriles, or 2-Arylethene-1,1-dicarbonitriles	540
11.15.1.2.9	Method 9:	From But-2-ynedinitrile or But-2-ynediamide	542
11.15.1.3	By Formation of One N—C Bond		542
11.15.1.3.1	Method 1:	From Propenes or Propanols	542
11.15.1.3.2	Method 2:	From Sulfinimidic Acid or Sulfoximides	543
11.15.1.3.3	Method 3:	From Thiolates	543
11.15.1.3.4	Method 4:	From Prop-2-enethiones	545
11.15.1.4	By Formation of One C—C Bond		546
11.15.2	Synthesis by Ring Transformation		547
11.15.2.1	Method 1:	From 1,2-Dithiolium Salts	547
11.15.2.2	Method 2:	From 1,4-Dithiin-2,3,5,6-tetracarbonitrile	549
11.15.2.3	Method 3:	From 1,4,2-Dithiazines	549
11.15.3	Synthesis by Substituent Modification		550
11.15.3.1	Substitution of Existing Substituents		550
11.15.3.1.1	Of Carbon Functionalities		550
11.15.3.1.1.1	Method 1:	Decarboxylation	550
11.15.3.1.1.2	Method 2:	N-Dealkylation of 2-Alkylisothiazolium Salts	551
11.15.3.1.2	Of Heteroatoms		552
11.15.3.1.2.1	Method 1:	Electrophilic Substitution	552
11.15.3.1.2.2	Method 2:	Nucleophilic Substitution	554
11.15.3.1.2.3	Method 3:	Isothiazole-5-carboxylic Acids, 5-carbaldehydes, and 5-Alkylisothiazoles from Isothiazol-5-yllithiums	557
11.15.3.1.2.4	Method 4:	From Isothiazolamines via Diazonium Salts	559

11.15.3.1.2.5	Method 5: Aryl and Heteroarylthiazoles from Isothiazolamines	560
11.15.3.2	Addition Reactions	561
11.15.3.2.1	Addition of Organic Groups	561
11.15.3.2.1.1	Method 1: Intramolecular S-Alkylation	561
11.15.3.2.1.2	Method 2: N-Alkylation and the Synthesis of 2-Alkylisothiazolium Salts	561
11.15.3.2.2	Addition of Heteroatoms	563
11.15.3.2.2.1	Method 1: Oxidation	563
11.15.3.3	Modification of Substituents	565
11.15.3.3.1	Method 1: From Acylthiazoles	565
11.15.3.3.2	Method 2: Modification of Side Chains	565
11.16	Product Class 16: Benzisothiazoles D. W. Brown and M. Sainsbury	
11.16	Product Class 16: Benzisothiazoles	573
11.16.1	Product Subclass 1: 1,2-Benzisothiazoles	574
11.16.1.1	Synthesis by Ring-Closure Reactions	577
11.16.1.1.1	By Formation of Two C—C Bonds	577
11.16.1.1.2	By Formation of One S—N or One N—C Bond	577
11.16.1.1.2.1	Method 1: From Thiols, Disulfides, and Related Compounds	578
11.16.1.1.2.2	Method 2: From Oximes	581
11.16.1.1.2.3	Method 3: From (Aminosulfonyl)arenes	583
11.16.1.1.2.4	Method 4: From Disulfides	586
11.16.1.1.2.5	Method 5: From 2-Acylbenzenesulfonamides or 2-(Sulfinyl)benzamides	587
11.16.1.1.3	By Formation of One S—C Bond	589
11.16.1.2	Synthesis by Ring Transformation	590
11.16.1.3	Synthesis by Substituent Modification	595
11.16.1.3.1	Substitution of Existing Substituents	595
11.16.1.3.1.1	Electrophilic Substitution	595
11.16.1.3.1.2	Nucleophilic Substitution	597
11.16.1.3.2	Addition Reactions	600
11.16.1.3.2.1	Addition of Organic Groups	600
11.16.1.3.2.1.1	Method 1: Alkylation of Saccharins	600
11.16.1.3.2.2	Addition of Heteroatoms	601
11.16.1.3.2.2.1	Method 1: Oxidation	601
11.16.2	Product Subclass 2: 2,1-Benzisothiazoles	603
11.16.2.1	Synthesis by Ring-Closure Reactions	607
11.16.2.1.1	By Formation of One S—N Bond	607

11.16.2.1.1.1	Method 1: From 2-Nitrophenylmethanethiols or 2-Nitrobenzenecarbothioamides	607
11.16.2.1.1.2	Method 2: From 2-Aminophenylmethanethiols or 2-Aminobenzenecarbothioamides	608
11.16.2.1.1.3	Method 3: From Isatoic Anhydride	610
11.16.2.1.1.4	Method 4: From 2-Methylanilines	611
11.16.2.1.1.5	Method 5: From Bis(2-aminophenyl)methane	614
11.16.2.2	Synthesis by Ring Transformation	615
11.16.2.2.1	From 2,1-Benzisoxazoles	615
11.16.2.3	Synthesis by Substituent Modification	616
11.16.2.3.1	Substitution of Existing Substituents	616
11.16.2.3.1.1	Of Hydrogen	616
11.16.2.3.1.2	Of Heteroatoms	617
11.16.2.3.1.2.1	Electrophilic Substitution	617
11.16.2.3.1.2.2	Nucleophilic Substitution	619
11.16.2.3.1.2.2.1	From 3-Chloro-2,1-benzisothiazoles	619
11.16.2.3.1.2.2.2	Method 2: From 2,1-Benzisothiazole 3-Diazonium Salts	620
11.16.2.3.2	Addition Reactions	620
11.16.2.3.2.1	Method 1: N-Alkylation	620
11.17	Product Class 17: Thiazoles D. Kikelj and U. Urleb	
11.17	Product Class 17: Thiazoles	627
11.17.1	Synthesis by Ring-Closure Reactions	632
11.17.1.1	By Formation of Three Heteroatom–Carbon Bonds and One C–C Bond	632
11.17.1.1.1	Fragments C–N, C, C, and S	632
11.17.1.1.1.1	From Alkali Cyanides and Hydrogen Sulfide	632
11.17.1.2	By Formation of Three Heteroatom–Carbon Bonds	633
11.17.1.2.1	By Formation of Two S–C Bonds and One N–C Bond	633
11.17.1.2.1.1	Fragments N–C, C–C, and S	633
11.17.1.2.1.1.1	Method 1: From Enamines, Cyanamide, and Sulfur	633
11.17.1.2.2	By Formation of One S–C Bond and Two N–C Bonds	634
11.17.1.2.2.1	Fragments S–C, C–C, and N	634
11.17.1.2.2.1.1	Method 1: From α -Bromoketimines and Ammonium Thiocyanate	634
11.17.1.2.2.2	Fragments S–C–C, N, and C	634
11.17.1.2.2.2.1	Method 1: From 5-Hydroxy-3-sulfanylpentan-2-one, Acetaldehyde, and Hydrazine	634
11.17.1.3	By Formation of Two Heteroatom–Carbon Bonds and One C–C Bond	635

11.17.1.3.1	Fragments S—C, N—C, and C	635
11.17.1.3.1.1	Method 1: From Chloridimidothiocarbonates, Dimethylthioformamide, and Isocyanides	635
11.17.1.4	By Formation of Two Heteroatom—Carbon Bonds	636
11.17.1.4.1	Fragments C—C—N—C and S	636
11.17.1.4.1.1	Method 1: From <i>N</i> -Vinylimines	636
11.17.1.4.1.2	Method 2: From <i>N</i> -(Polychloroalkyl)imidoyl Chlorides	637
11.17.1.4.1.3	Method 3: From <i>N</i> -Functionalized α -Aminonitriles	637
11.17.1.4.1.4	Method 4: From α -Acylamino Carbonyl Compounds	638
11.17.1.4.1.5	Method 5: From 2-Isocyanoacrylates	639
11.17.1.4.1.6	Method 6: From Sulfanyl-Substituted Alkenones	640
11.17.1.4.1.7	Method 7: From <i>N</i> -Alkylimines	641
11.17.1.4.2	Fragments S—C—N and C—C	642
11.17.1.4.2.1	Method 1: From Hypervalent Alkynyliodonium Salts and Aminothiocarbonyl Compounds	642
11.17.1.4.2.2	Method 2: From α -Functionalized Carboxylic Acid Derivatives and Aminothiocarbonyl Compounds	643
11.17.1.4.2.2.1	Variation 1: From α -Functionalized Carboxylic Acid Derivatives and Aminothiocarbonyl Compounds	643
11.17.1.4.2.2.2	Variation 2: From α -Functionalized Carbonitriles and Aminothiocarbonyl Compounds	645
11.17.1.4.2.3	Method 3: From α,β -Unsaturated Carboxylic Acid Derivatives and Aminothiocarbonyl Compounds	646
11.17.1.4.2.4	Method 4: From α -Functionalized Ketones or Ketone Derivatives and Aminothiocarbonyl Compounds	647
11.17.1.4.2.4.1	Variation 1: From α -Diazo Ketones and Aminothiocarbonyl Compounds	647
11.17.1.4.2.4.2	Variation 2: From α -Halo Ketones and Aminothiocarbonyl Compounds	647
11.17.1.4.2.4.3	Variation 3: From α -Tosyloxy-Substituted Ketones and Aminothiocarbonyl Compounds	654
11.17.1.4.2.4.4	Variation 4: From Acyloxiranes and Aminothiocarbonyl Compounds	656
11.17.1.4.2.4.5	Variation 5: From Ketones Halogenated In Situ and Aminothiocarbonyl Compounds	656
11.17.1.4.2.4.6	Variation 6: From α -Halomethylketimines and Aminothiocarbonyl Compounds	657
11.17.1.4.2.4.7	Variation 7: From 2-(2-Bromo-1-phenylethylene)malononitrile and Aminothiocarbonyl Compounds	658
11.17.1.4.2.5	Method 5: From α,β -Unsaturated Ketones and Aminothiocarbonyl Compounds	658
11.17.1.4.2.6	Method 6: From α -Halo-Substituted Aldehydes and Aminothiocarbonyl Compounds	658
11.17.1.4.2.6.1	Variation 1: From Thioureas	658
11.17.1.4.2.6.2	Variation 2: From Thioamides	659
11.17.1.4.2.7	Method 7: From Enamines and Aminothiocarbonyl Compounds	660
11.17.1.4.2.8	Method 8: From 1-Aryl-Substituted 2,2,2-Trichloroethanols and Aminothiocarbonyl Compounds	661
11.17.1.4.2.9	Method 9: From 3-Haloalk-1-enes or 3-Haloalk-1-yne and Aminothiocarbonyl Compounds	661

11.17.1.4.2.10	Method 10: From Thiocyanates and α -Halo Carbonyl Compounds	662
11.17.1.4.3	Fragments S—C—C—N and C	662
11.17.1.4.3.1	Method 1: From 2-Aminoalkanethioic Acids and “One-Carbon” Reagents	662
11.17.1.4.3.2	Method 2: From Ethanedithioamides and Aldehydes	663
11.17.1.4.4	Fragments C—C—S and C—N	664
11.17.1.4.4.1	Method 1: From 2-Sulfanylalkanoic Acids or Derivatives and Nitriles	664
11.17.1.4.4.2	Method 2: From Chlorothioacetic S-Acid and Nitriles	666
11.17.1.4.4.3	Method 3: From β -Amino α,β -Unsaturated Esters or Ketones and Sulfanylacetic Acid Derivatives	666
11.17.1.4.4.4	Method 4: From α -Sulfanyl Ketones and Nitriles	667
11.17.1.4.4.4.1	Variation 1: Generation of α -Sulfanyl Ketones from S-2-Oxoalkyl Dithiocarbonates	668
11.17.1.4.4.4.2	Variation 2: Generation of α -Sulfanyl Ketones from Their Dimers	669
11.17.1.4.4.5	Method 5: Additional Methods	669
11.17.1.4.5	Fragments C—C—N and C—S	669
11.17.1.4.5.1	Method 1: From α -Aminonitriles and Carbonyl Sulfide, Carbon Disulfide, or Isocyanates	669
11.17.1.4.5.2	Method 2: From Carboxamides and Carbon Disulfide	671
11.17.1.4.5.3	Method 3: From α -Amino Acids or Derivatives and Dithiocarboxylic Acids	671
11.17.1.4.5.4	Method 4: From α -Aminonitriles and Thiocarboxylic/Dithiocarboxylic Acids or Derivatives	672
11.17.1.4.5.5	Method 5: From Enaminonitriles and Thiocarboxamides	673
11.17.1.4.5.6	Method 6: From β -Cyano- or β -Nitroenamines and Thiocyanogen	674
11.17.1.4.5.7	Method 7: From Vinylamines and Chloroformylsulfenyl Chlorides or Imidosulfenyl Chlorides	674
11.17.1.4.5.8	Method 8: From α -Halooximes and Thiocyanates	675
11.17.1.4.5.9	Method 9: From Alk-2-ynylamines and Isothiocyanates or Carbon Disulfide	675
11.17.1.4.5.10	Method 10: Additional Methods	676
11.17.1.4.6	Fragments C—C—S—C and N	676
11.17.1.5	By Formation of One Heteroatom—Carbon Bond and One C—C Bond	677
11.17.1.5.1	Fragments C—N—C—S and C	677
11.17.1.5.1.1	Method 1: From Dipotassium Cyanodithioimidocarbonate and Functionalized Methyl Halides	677
11.17.1.5.1.2	Method 2: From Cyanothioimidocarbonates or Cyanodithioimidocarbonate and “Acidic” Methyl Halides	678
11.17.1.5.1.3	Method 3: From [Amino(nitroamino)methylene]dithiocarbamates and “Acidic” Methyl Halides	680
11.17.1.5.1.4	Method 4: From Alkali Salts of <i>N</i> -Cyanothioureas or <i>N</i> -Cyanocarbothioamides and “Acidic” Methyl Halides	680
11.17.1.5.1.5	Method 5: From <i>N</i> -(Diaminomethylene)thioureas and α -Halo Ketones	681
11.17.1.5.1.6	Method 6: From <i>N</i> -Imidoylthioureas or <i>N</i> -Thiocarbamoylimidates and “Acidic” Methyl Halides	682
11.17.1.5.1.7	Method 7: From <i>N</i> -Acylthiocarbamates or <i>N</i> -Acylcarbamates and “Acidic” Methyl Halides	684
11.17.1.5.1.8	Method 8: From <i>N</i> -Acylthioureas and “Acidic” Methyl Halides	684

11.17.1.5.1.9	Method 9:	From <i>N</i> -Imidoylecarbothioamides or <i>N</i> -Aroylcarbothioamides and “Acidic” Methyl Halides	685
11.17.1.5.1.10	Method 10:	Additional Methods	687
11.17.1.5.2	Fragments C—N—C and S—C		687
11.17.1.5.2.1	Method 1:	From <i>N</i> -Thiocarbamoylimidates and Methyl Sulfanylacetate ..	687
11.17.1.5.2.2	Method 2:	From <i>N</i> -Cyanoimidothioates, <i>S</i> -Methyl- <i>N</i> -cyanoisoureas, or Dimethyl Cyanocarbonodithioimidoate and Sulfanylacetic Acid Derivatives	688
11.17.1.5.2.3	Method 3:	From <i>N</i> -(Chloromethylene)imidamidium Salts and Sulfanylacetates or Sulfanylacetone	688
11.17.1.5.2.4	Method 4:	From Alkyl Isothiocyanates and S—C Synthons	689
11.17.1.5.2.5	Method 5:	From (Ethoxycarbonylmethyl)imidodithiocarbonate or Ethyl [[1-(Methylsulfanyl)alkylidene]amino]acetates	690
11.17.1.5.2.6	Method 6:	From Alkyl Isocyanides and S—C Synthons	692
11.17.1.5.2.7	Method 7:	From Nitrile Ylides and S—C Synthons	693
11.17.1.5.2.8	Method 8:	Additional Methods	694
11.17.1.5.3	Fragments C—S—C—N and C		694
11.17.1.5.3.1	Method 1:	From Isothioureas and Carboxylic Acid Derivatives	694
11.17.1.5.4	Fragments C—S—C and N—C		695
11.17.1.5.4.1	Method 1:	From Phenacyl Thiocyanates and Nitriles	695
11.17.1.6	By Formation of One Heteroatom—Carbon Bond		696
11.17.1.6.1	By Formation of One S—C Bond		696
11.17.1.6.1.1	Fragment C—C—N—C—S		696
11.17.1.6.1.1.1	Method 1:	From <i>N</i> -Thiocarbonyl α -Amino Acids and Derivatives	697
11.17.1.6.1.1.2	Method 2:	From α -(Thioacylamino)carbothioamides	698
11.17.1.6.1.1.3	Method 3:	From α -Thioacylamino Ketones, α -Thioureido Ketones and α -Thioacylamino Acetals	699
11.17.1.6.1.1.4	Method 4:	From α -(Aroylamino) Thioketones	700
11.17.1.6.1.1.5	Method 5:	From α -(Thioacylamino)carbonitriles	701
11.17.1.6.1.1.6	Method 6:	From <i>N</i> -(α -Haloacyl)-, <i>N</i> -(α -Haloalkylidene), <i>N</i> -(α -Haloalkyl)-, <i>N</i> -(α -Haloalkenyl)-Substituted Thioureas, Isothiocyanates, or Carbothioamides	701
11.17.1.6.1.1.7	Method 7:	From Isothiocyanatoallenes	704
11.17.1.6.1.1.8	Method 8:	Additional Methods	704
11.17.1.6.1.2	Fragment C—N—C—C—S		704
11.17.1.6.1.2.1	Method 1:	From α -(Acylamino)carbothioamides	705
11.17.1.6.1.2.2	Method 2:	From 2-(Benzoylamino)-Substituted Triphenylphosphonium Salts	705
11.17.1.6.1.2.3	Method 3:	Thiazoles from <i>N</i> ¹ , <i>N</i> ² -Bis[amino(aryl)methyl]ethanedithioamides	706
11.17.1.6.1.2.4	Method 4:	Additional Methods	706
11.17.1.6.2	By Formation of One N—C Bond		707
11.17.1.6.2.1	Fragment C—C—S—C—N		707

11.17.1.6.2.1.1	Method 1:	From α -Thiocyanato Carboxylic Acids and Derivatives	707
11.17.1.6.2.1.2	Method 2:	From S-(1-Iminoalkyl)thioglycolic Acids	708
11.17.1.6.2.1.3	Method 3:	From Cyanomethyl Carbimidothioates	709
11.17.1.6.2.1.4	Method 4:	From α -[(Iminomethyl)sulfanyl] Ketones	710
11.17.1.6.2.1.5	Method 5:	From α -Thiocyanato Ketones	711
11.17.1.6.2.1.6	Method 6:	From 2-(Hydroxyimino)alkyl Carbamodithioates or 2-[2-(Hydroxyimino)alkyl]isothioureas	714
11.17.1.6.2.1.7	Method 7:	From α -Halo- β -thiocyanatoalkenes	714
11.17.1.6.2.1.8	Method 8:	From Alk-1-ynyl Thiocyanates	715
11.17.1.6.2.2	Fragment C—S—C—C—N		715
11.17.1.6.2.2.1	Method 1:	From 2-(Hydroxyimino)alkyl Dithiocarboxylates and Carbonodithioates	715
11.17.1.6.2.2.2	Method 2:	From β -Thiocyanatoenamines	716
11.17.1.6.2.2.3	Method 3:	From N-Substituted 2-[(Thioacyl)sulfanyl]acetamides	717
11.17.1.6.2.2.4	Method 4:	From Benzyl α -(Hydroxyimino)imidothioates	717
11.17.1.6.2.2.5	Method 5:	Additional Methods	718
11.17.1.7	By Formation of One C—C Bond		718
11.17.1.7.1	Fragment C—S—C—N—C		718
11.17.1.7.1.1	Method 1:	From Dialkyl Cyanodithioimidocarbonates or N-Alkyl N'-Cyanoimidothiocarbamates	718
11.17.1.7.1.2	Method 2:	From N-[1-(Alkylsulfanyl)alkylidene]-N'-dimethylimidamidium and -guanidinium Salts	719
11.17.1.7.1.3	Method 3:	From Methyl Acylcarbonodithioimidoates, Methyl Acylcarbamiimidothioates, or Methyl N-Acylarenecarbimidothioates	720
11.17.2	Synthesis by Ring Transformation		721
11.17.2.1	Method 1:	By Ring Enlargement of Three- or Four-Membered Heterocycles	721
11.17.2.2	Method 2:	By Formal Exchange of Ring Atoms with Retention of Ring Size	722
11.17.2.3	Method 3:	By Ring Contraction of Heterocycles	724
11.17.3	Aromatization		725
11.17.3.1	Method 1:	By Dehydration and Dehydroamination of Dihydrothiazoles	725
11.17.3.2	Method 2:	Aromatization by Addition Reactions	726
11.17.3.2.1	Variation 1:	Nucleophilic Additions to Exocyclic Double Bonds	726
11.17.3.2.2	Variation 2:	S-Alkylation of Thiazole-2(3H)-thiones	728
11.17.3.2.3	Variation 3:	N-Alkylation or N-Acylation of 3-Hydroxythiazol-2(3H)-imine	728
11.17.3.2.4	Variation 4:	Elimination of Sulfur from Thiazole-2(3H)-thiones	729
11.17.3.2.5	Variation 5:	Of Carbofunctional Groups	730
11.17.3.2.5.1	Oxidation of Dihydrothiazoles		730
11.17.3.2.5.1.1	2,3-Dihydrothiazoles		730
11.17.3.2.5.1.2	2,5-Dihydrothiazoles		730
11.17.3.2.5.2	Oxidation of Thiazolidines		732
11.17.3.3	Method 3:	Aromatization by Rearrangement (Isomerization)	733
11.17.4	Synthesis from Other Thiazoles		735

11.17.4.1	Addition Reactions	735
11.17.4.1.1	Quaternization	735
11.17.4.1.3	Addition of Heteroatoms	738
11.17.4.1.3.1	Formation of Thiazole 3-Oxides	738
11.17.4.1.3.2	Formation of 3-Aminothiazolium Salts	739
11.17.4.2	Synthesis by Substituent Modification	739
11.17.4.2.1	Method 1: Replacement of Hydrogen by Deuterium	739
11.17.4.2.2	Method 2: Replacement of Hydrogen by Metals	740
11.17.4.2.2.1	Variation 1: Replacement of Hydrogen by Lithium, Sodium, or Potassium	740
11.17.4.2.2.2	Variation 2: Replacement of Hydrogen by Magnesium, Aluminum or Zinc	741
11.17.4.2.2.3	Variation 3: Replacement of Hydrogen by Silicon or Tin	742
11.17.4.2.2.4	Variation 4: Replacement of Hydrogen by Mercury	743
11.17.4.2.3	Method 3: Replacement of Hydrogen by Carbon Electrophiles	743
11.17.4.2.4	Method 4: Introduction of Nucleophiles	746
11.17.4.2.5	Method 5: Via Radical Reactions	747
11.17.4.2.6	Method 6: By Rearrangement of 2-(2-Arylhiazino)thiazoles	749
11.17.4.2.7	Method 7: By Electrophiles Other Than Carbon Electrophiles	749
11.17.4.2.7.1	Variation 1: Halogenation of Thiazoles and Thiazole 3-Oxides	749
11.17.4.2.7.2	Variation 2: Introduction of Chlorosulfonyl, Sulfo, or Thiocyanato Groups	752
11.17.4.2.7.3	Variation 3: Introduction of Alkylsulfanyl, Arylsulfanyl, Thiolate, or Arylselanyl Groups	753
11.17.4.2.7.4	Variation 4: Nitration and Nitrosation of Thiazoles	754
11.17.4.2.7.5	Variation 5: Introduction of Arylazo Groups	756
11.17.4.2.8	Method 8: Introduction of Amino Groups	757
11.17.4.2.9	Additional Methods	758
11.17.4.3	Synthesis of Thiazoles from Metalated Thiazoles	758
11.17.4.3.1	Method 1: Replacement of Lithium, Magnesium, or Zinc by Carbon Electrophiles	758
11.17.4.3.2	Method 2: Replacement of Sodium by Carbon Electrophiles	762
11.17.4.3.3	Method 3: Replacement of Silyl or Mercurio Groups by Hydrogen	763
11.17.4.3.4	Method 4: Replacement of Silyl and Stannyl Groups by Carbon Electrophiles	764
11.17.4.3.5	Method 5: Replacement of Silyl, Stannyl or Mercury Containing Groups by Halogens or Thiocyanate	766
11.17.4.3.6	Method 6: Replacement of Metals by Other Groups	767
11.17.4.3.6.1	Variation 1: Replacement of Lithium by Sulfur- or Phosphorus-Containing Reagents	767
11.17.4.3.7	Method 7: Modification of Other Substituents	770
11.17.4.3.7.1	Variation 1: Carbon Replacement by Hydrogen	770
11.17.4.3.7.1.1	Decarboxylation of Thiazolecarboxylic Acids and Carboxylates	770
11.17.4.3.7.1.2	Removal of the Benzoyl Group from 5-Benzoylthiazoles	771
11.17.4.3.7.1.3	Additional Methods	772
11.17.4.3.7.2	Variation 2: Replacement of One Carbofunctional Group by Another	772
11.17.4.3.7.3	Variation 3: Replacement of a Carbon Substituent by Nitrogen	772

11.17.4.3.7.4	Variation 4: Replacement of a Nitrogen Substituent by Hydrogen	773
11.17.4.3.7.5	Variation 5: Replacement of a Nitrogen Substituent by Carbon	775
11.17.4.3.7.6	Variation 6: Substitution of One Nitrogen Substituent by Another	775
11.17.4.3.7.6.1	Additional Methods	776
11.17.4.3.7.7	Variation 7: Nitrogen Substituents Replacement by Halogen	777
11.17.4.3.7.8	Variation 8: Replacement of a Nitrogen Substituent by Oxygen or Sulfur	778
11.17.4.3.7.9	Variation 9: Replacement of Phosphorus Groups by Hydrogen	779
11.17.4.3.7.10	Variation 10: Replacement of Oxygen and Sulfur Groups by Hydrogen	780
11.17.4.3.7.11	Variation 11: Replacement of Sulfur or Oxygen Groups by Carbon-Containing Substituents	781
11.17.4.3.7.12	Variation 12: Replacement of Oxygen or Sulfur Substituents by Nitrogen	782
11.17.4.3.7.13	Variation 13: Replacement of Oxygen or Selenium by Halogen	784
11.17.4.3.7.14	Variation 14: Replacement of Oxygen by Sulfur	784
11.17.4.3.7.15	Variation 15: Replacement of Sulfur by Oxygen	785
11.17.4.3.7.16	Variation 16: Replacement of Sulfur by Sulfur	785
11.17.4.3.7.17	Variation 17: Replacement of Halogen by Hydrogen	786
11.17.4.3.7.18	Variation 18: Replacement of Halogen by Deuterium	788
11.17.4.3.7.19	Variation 19: Replacement of Halogen by Carbon Nucleophiles	788
11.17.4.3.7.20	Variation 20: Replacement of Halogen by Nitrogen	789
11.17.4.3.7.21	Variation 21: Replacement of Halogen by Oxygen or Sulfur	791
11.17.4.3.7.22	Variation 22: Replacement of Halogens by Other Halogens	793
11.17.5	Modification of Substituents	794
11.17.5.1	Modification of Carbofunctional Substituents	794
11.17.5.1.1	Method 1: Condensation at Alkyl Groups of Alkylthiazoles, Alkylthiazolium Salts, and Thiazole 3-Oxides	794
11.17.5.1.2	Method 2: Other Transformations of Carbofunctional Substituents of Thiazoles	799
11.17.5.2	Modification of Heterofunctional Substituents	800
11.17.5.2.1	Method 1: Modification of O- and S-Substituents	801
11.17.5.2.2	Modification of N-Substituents	803
11.17.5.2.2.1	Method 1: Reduction of Thiazole 3-Oxides	803
11.17.5.3	Rearrangement of Substituents	804
11.17.5.3.1	Rearrangement during Transmetalation Reaction	804
11.17.5.3.2	Rearrangement of Thiazole 3-Oxides	804
11.17.5.3.3	Rearrangement of 3-Substituted Thiazol-2(3 <i>H</i>)-imines and Thiazole-2(3 <i>H</i>)-thiones	805
11.17.5.3.4	Rearrangement of Nitrothiazoles	806
11.17.5.3.5	Photochemical Rearrangements	806
11.17.5.3.6	Rearrangement of 3-Benzylthiazolium Salts	806
11.17.5.3.7	Other Rearrangements and Ring Cleavages	806

11.18	Product Class 18: Benzothiazoles and Related Compounds H. Ulrich	
11.18	Product Class 18: Benzothiazoles and Related Compounds	835
11.18.1	Synthesis by Ring-Closure Reactions	837
11.18.1.1	By Annulation to an Arene or Hetarene Ring	837
11.18.1.1.1	By Formation of One S—C and One N—C Bond	837
11.18.1.1.1.1	Method 1: From 2-Amino(het)arenethiols	837
11.18.1.1.1.1.1	Variation 1: With Phosgene	837
11.18.1.1.1.1.2	Variation 2: With Dithiocarbonates	838
11.18.1.1.1.1.3	Variation 3: With Carbon Disulfide	838
11.18.1.1.1.1.4	Variation 4: With Dithiocarbamates	839
11.18.1.1.1.1.5	Variation 5: With Isothiocyanates	840
11.18.1.1.1.1.6	Variation 6: With Cyanamides	840
11.18.1.1.1.1.7	Variation 7: With Derivatives of Cyanamides	841
11.18.1.1.1.1.8	Variation 8: With Orthoesters	841
11.18.1.1.1.1.9	Variation 9: With <i>N</i> -Chloromethylenamines	843
11.18.1.1.1.1.10	Variation 10: With Acyl Chlorides	843
11.18.1.1.1.1.11	Variation 11: With Acyl Chlorides Generated In Situ	844
11.18.1.1.1.1.12	Variation 12: With Acyl Chlorides Using Salts of the Thiol	844
11.18.1.1.1.1.13	Variation 13: With Imidoyl Chlorides and Derivatives	844
11.18.1.1.1.1.14	Variation 14: With Carboxylic Anhydrides	846
11.18.1.1.1.1.15	Variation 15: With Carboxylic or Thiocarboxylic Esters	846
11.18.1.1.1.1.16	Variation 16: With Carboxylic Acids	847
11.18.1.1.1.1.17	Variation 17: With Ketene Acetals	848
11.18.1.1.1.1.18	Variation 18: With Dithioimidocarbonates	849
11.18.1.1.1.1.19	Variation 19: With Nitriles	849
11.18.1.1.1.1.20	Variation 20: With Trifluoromethyl(het)arenes	850
11.18.1.1.1.1.21	Variation 21: With Aliphatic Fluoro Compounds	851
11.18.1.1.1.1.22	Variation 22: With Aldehydes	851
11.18.1.1.1.1.23	Variation 23: With Activated Ketones	852
11.18.1.1.1.1.24	Variation 24: With Carbon Monoxide and Haloarenes	852
11.18.1.1.1.1.25	Variation 25: With Oxiranes	853
11.18.1.1.1.1.26	Variation 26: With a Cyclopropenylum Salt	854
11.18.1.1.1.2	Method 2: From 2-Halonitrobenzenes	854
11.18.1.1.1.3	Method 3: From 2-Unsubstituted (Het)arenamines or Derivatives and a Sulfur Component	854
11.18.1.1.1.3.1	Variation 1: With Carbon Disulfide and Sulfur	854
11.18.1.1.1.3.2	Variation 2: With Chlorocarbonylsulfonyl Chloride	855
11.18.1.1.1.3.3	Variation 3: With Thiocyanates	855
11.18.1.1.1.3.4	Variation 4: With Disulfur Dichloride	856
11.18.1.1.1.3.5	Variation 5: With Phenylphosphonothioic Dichloride	856
11.18.1.1.1.3.6	Variation 6: With Sulfur	857
11.18.1.1.1.4	Method 4: From 2-Substituted (Het)arenamines or Derivatives and a Sulfur Component	857
11.18.1.1.1.4.1	Variation 1: With Dithiocarbonates	857

11.18.1.1.1.4.2	Variation 2:	With Thioureas	858
11.18.1.1.1.4.3	Variation 3:	With <i>O</i> -Alkyl Thiocarboxylates	858
11.18.1.1.1.4.4	Variation 4:	With Thioamides	858
11.18.1.1.1.4.5	Variation 5:	With Benzoylsulfanyl copper	859
11.18.1.1.1.5	Method 5:	From 2-Substituted Nitro(het)arenes and a Sulfur Component	859
11.18.1.1.1.6	Method 6:	From Benzo-1,4-quinones or Derivatives and a Sulfur Component	860
11.18.1.1.1.7	Method 7:	From 2-Chlorohetarene carbonitriles and Thiourea	861
11.18.1.1.2	By Formation of One S—C Bond		861
11.18.1.1.2.1	Method 1:	From <i>N</i> -Arylthioureas	861
11.18.1.1.2.1.1	Variation 1:	2-Unsubstituted <i>N</i> -Arylthioureas	862
11.18.1.1.2.1.2	Variation 2:	With a Leaving Group in the 2-Position	864
11.18.1.1.2.1.3	Variation 3:	In Situ Preparation of Thioureas from Isothiocyanates	865
11.18.1.1.2.1.4	Variation 4:	From Guanidyl Disulfides	865
11.18.1.1.2.2	Method 2:	From Isothioureas	866
11.18.1.1.2.3	Method 3:	From <i>N</i> -Arylthioamides	866
11.18.1.1.2.3.1	Variation 1:	From 2-Unsubstituted <i>N</i> -Arylthioamides	866
11.18.1.1.2.3.2	Variation 2:	From <i>N</i> -Arylthioamides with a Halogen or Hydroxy Leaving Group in the 2-Position	867
11.18.1.1.2.3.3	Variation 3:	In Situ Formation from Amides	868
11.18.1.1.2.4	Method 4:	From 2-Sulfanylacetophenone Oximes	869
11.18.1.1.2.5	Method 5:	From <i>ortho</i> -Nitrogen-Functionalized Aryl Sulfides or Disulfides	869
11.18.1.1.2.5.1	Variation 1:	From Amides	870
11.18.1.1.2.5.2	Variation 2:	From Isocyanides	871
11.18.1.1.2.5.3	Variation 3:	From Azomethines	871
11.18.1.1.2.5.4	Variation 4:	From Iminophosphoranes	872
11.18.1.1.2.5.5	Variation 5:	From Amines	873
11.18.1.1.3	By Formation of One N—C Bond		873
11.18.1.1.3.1	Method 1:	From 2-Thiocyanatoanilines	873
11.18.1.1.3.1.1	Variation 1:	Using 2-Thiocyanatoanilines	873
11.18.1.1.3.1.2	Variation 2:	Masked as a Cycloadduct with Azides	874
11.18.1.1.3.1.3	Variation 3:	Formed In Situ from 2-Nitroaryl Thiocyanates	874
11.18.1.1.3.2	Method 2:	From <i>S</i> -(2-Aminohetaryl) Dithiocarbamates	875
11.18.1.1.3.2.1	Variation 1:	Using <i>S</i> -(2-Aminohetaryl) Dithiocarbamates	875
11.18.1.1.3.2.2	Variation 2:	Formed In Situ from <i>S</i> -(2-Nitroaryl) Dithiocarbamates	875
11.18.1.1.3.3	Method 3:	From 2-(2-Nitroarylsulfanyl)cycloalkanones	876
11.18.1.2	By Annulation to the Thiazole Ring		876
11.18.1.2.1	Method 1:	Of a Pyrido[2,3] System	876
11.18.1.2.1.1	Variation 1:	From 3-(4-Aminothiazol-5-yl)-3-oxopropanoates	876
11.18.1.2.1.2	Variation 2:	From 4-Amino-5-benzoylthiazolium Salts	877
11.18.1.2.1.3	Variation 3:	From 4-Amino-Substituted Thiazoles and Acetoacetates	877
11.18.1.2.2	Method 2:	Of a Pyrido[3,2] System	877
11.18.1.2.3	Method 3:	Of a Pyrido[4,3] System	878
11.18.1.2.4	Method 4:	Of a Pyridazino[4,5] System	878
11.18.1.2.5	Method 5:	Of a Pyrimido[4,5] System	879
11.18.1.2.6	Method 6:	Of a Pyrimido[5,4] System	879

11.18.2	Synthesis by Ring Transformation	880
11.18.2.1	Hetero-Ring Size Retained Overall	880
11.18.2.1.1	Method 1: From Other Heterocycles	880
11.18.2.1.2	Method 2: By Ring Opening and Reclosure	881
11.18.2.2	With Ring Contraction	881
11.18.2.2.1	Method 1: From 1,4-Benzothiazines and Derivatives	881
11.18.2.2.2	Method 2: From 1,2,4-Benzothiadiazines	882
11.18.2.2.3	Method 3: From 1,5-Benzothiazepines	883
11.18.2.2.4	Method 4: From Tetrahydrodibenzo-1,6,3,8-dithiadiazecines	884
11.18.2.3	With Simultaneous Annulation of a Phenyl Substituent	884
11.18.2.3.1	Method 1: From 5-Aryl-1,2,3-dithiazoles	885
11.18.2.3.2	Method 2: From 5-(Arylimino)thiatriazoles	885
11.18.2.3.3	Method 3: From Heteropentalenes	885
11.18.3	Aromatization	886
11.18.3.1	Method 1: Of the Thiazole Unit	886
11.18.3.1.1	Variation 1: By Dehydrogenation	886
11.18.3.1.2	Variation 2: By Elimination	887
11.18.3.2	Method 2: Of the Benzene Unit	888
11.18.4	Synthesis by Substituent Modification	888
11.18.4.1	Addition Reactions	888
11.18.4.1.1	Method 1: Protonation and Complex Formation	888
11.18.4.1.2	Method 2: Alkylation	889
11.18.4.2	Substitution of Existing Substituents	890
11.18.4.2.1	Of Hydrogen	890
11.18.4.2.1.1	Method 1: By Deuterium	890
11.18.4.2.1.2	Method 2: By Carbon Groups	891
11.18.4.2.1.3	Method 3: By Metals	891
11.18.4.2.1.4	Method 4: By Halogen	891
11.18.4.2.1.5	Method 5: Sulfonation	892
11.18.4.2.1.6	Method 6: Nitration	893
11.18.4.2.2	Of Metals	893
11.18.4.2.2.1	Method 1: Alkylation	893
11.18.4.2.2.2	Method 2: Hydroxyalkylation	894
11.18.4.2.2.3	Method 3: Acylation	894
11.18.4.2.2.4	Method 4: Transmetalation	895
11.18.4.2.2.5	Method 5: By Heteroatoms	896
11.18.4.2.3	Of Carbon Functionalities	896
11.18.4.2.4	Of Heteroatoms	896
11.18.4.2.4.1	Method 1: By Addition of Oxygen	897
11.18.4.2.4.2	Method 2: By Elimination of Oxygen	897
11.18.4.2.4.3	Method 3: By Hydrogen	898

11.18.4.2.4.4	Method 4: By Carbon Functionalities	898
11.18.4.2.4.5	Method 5: By Hetero Functions	900
11.18.4.2.4.6	Method 6: By Rearrangement of Substituents	901
11.18.4.3	Modification of Substituents in the α -Position	902
11.18.4.3.1	Method 1: α -Hydroxyalkylation	902
11.18.4.3.2	Method 2: Condensation Reactions	903
11.18.4.3.3	Method 3: Self-Condensation	904
11.18.4.3.4	Method 4: α -Oxygenation	904
11.18.4.3.5	Method 5: Oxirane Formation	904
11.18.4.3.6	Method 6: Aziridine Formation	905
11.18.4.3.7	Method 7: Sulfur Oxidation	905
11.19	Product Class 19: 1,2- and 1,3-Thiaphospholes and Benzoannulated Analogues D. Gudat	
11.19	Product Class 19: 1,2- and 1,3-Thiaphospholes and Benzoannulated Analogues	913
11.19.1	Product Subclass 1: 1,2-Thiaphospholes	913
11.19.1.1	Synthesis by Ring Transformation	913
11.19.1.1.1	Method 1: Desulfurization of 2-Thio-3 <i>H</i> -1,2-thiaphospholes	913
11.19.1.1.2	Method 2: Additional Methods	915
11.19.2	Product Subclass 2: 1,3-(Benzo)Thiaphospholes	915
11.19.2.1	Synthesis by Ring-Closure Reactions	915
11.19.2.1.1	By Formation of Two P—C Bonds	915
11.19.2.1.1.1	Fragments C—S—C—C and P	915
11.19.2.1.1.1.1	Method 1: From Ketene <i>S,N</i> -Acetals and Phosphorus Trihalides	915
11.19.2.1.2	By Formation of One S—C and One P—C Bond	916
11.19.2.1.2.1	Method 1: From 2-Phosphinobenzenethiols	916
11.19.2.2	Synthesis by Ring Transformation	917
11.19.2.2.1	Method 1: From Thiaphosphiranes and Acetylenedicarboxylates	917
11.19.2.2.2	Method 2: From 1,3-Ditholium-4-olates	917
11.19.2.2.2.1	Variation 1: Reaction with Tris(trimethylsilyl)phosphine	917
11.19.2.2.2.2	Variation 2: Reaction with Phosphaalkynes or Phosphaalkenes	918
11.20	Product Class 20: Isoselenazoles W.-D. Pfeiffer	
11.20	Product Class 20: Isoselenazoles	921
11.20.1	Synthesis by Ring-Closure Reactions	922
11.20.1.1	By Formation of One Se—N, One Se—C, and One N—C Bond	922

11.20.1.1.1	Fragments C—C—C, N, and Se	922
11.20.1.1.1.1	Method 1: From Alkynones, Hydroxylamine O-Sulfonic Acid, and Selenium	922
11.20.1.1.1.2	Method 2: From 3-Chloroprop-2-eniminium Salts, Sodium Selenocyanate, and Ammonia	923
11.20.1.2	By Formation of One Se—N and One Se—C Bond	924
11.20.1.2.1	Fragments Se—C—C—C and N	924
11.20.1.2.1.1	Method 1: From 3-Seleno-Substituted Enones and Ammonia	924
11.20.1.2.2	Fragments N—C—C—C and Se	925
11.20.1.2.2.1	Method 1: From an Enamine and Selenium Dioxide	925
11.20.1.3	By Formation of One Se—N Bond	926
11.20.1.3.1	Fragment Se—C—C—N	926
11.20.1.3.1.1	Method 1: From Seleno-Substituted Alkenoic Amides	926
11.20.2	Synthesis by Substituent Modification	927
11.20.2.1	By Substitution of Existing Substituents	927
11.20.2.1.1	Of Hydrogen	927
11.20.2.1.1.1	Method 1: Bromination	927
11.20.2.1.1.2	Method 2: Nitration	928
11.20.2.1.2	Of Heteroatoms	928
11.20.2.1.2.1	Method 1: Hydrolysis of 3-Chloroisoselenazolium Salts	928
11.20.2.2	Modification of Substituents	929
11.20.2.2.1	Method 1: Oxidation of Methyl-Substituted Isoselenazoles	929
11.21	Product Class 21: Annulated Isoselenazole Compounds W.-D. Pfeiffer	
11.21	Product Class 21: Annulated Isoselenazole Compounds	931
11.21.1	Synthesis by Ring-Closure Reactions	933
11.21.1.1	By Ring-Closure of One Se—N and C—N Bond	933
11.21.1.1.1	Fragments Se—Arene—C and N	933
11.21.1.1.1.1	Method 1: From 2-Seleno-Substituted Arenecarbonyl Compounds and a Nitrogen Source	933
11.21.1.1.1.1.1	Variation 1: From 2-Seleno-Substituted Aldehydes or Ketones	933
11.21.1.1.1.1.2	Variation 2: From 2-(Methylselanyl)thiophene-3-carbaldehyde or 2-(Methylselanyl)selenophene-3-carbaldehyde	934
11.21.1.1.1.1.3	Variation 3: From 2-(Chloroselanyl)benzoyl Chloride	935
11.21.1.1.1.1.4	Variation 4: From 2-(Selenocyanato)phenyl Ketones	935
11.21.1.2	By Formation of One Se—N Bond	936
11.21.1.2.1	Fragment Se—Arene—C—N	936
11.21.1.2.1.1	Method 1: Cyclization of 2-(Methylselanyl)benzamides	936

11.21.2	Synthesis by Ring Transformation	936
11.21.2.1	Method 1: 1,2-Benzisoselenazoles from 1-Benzoselenophene-2,3-dione	936
11.21.2.2	Method 2: 1,2-Benzisoselenazoles from 3,4-Diphenyl-1,2,5-selenadiazole	937
11.21.3	Synthesis by Substituent Modification	937
11.21.3.1	Nucleophilic Substitution	937
11.21.3.1.1	Method 1: By Potassium Amide	937
11.21.3.1.2	Method 2: From 3-Chloro-1,2-benzisoselenazolium Salts	938
11.21.4	Modification of Substituents	938
11.22	Product Class 22: Selenazoles W.-D. Pfeiffer	
11.22	Product Class 22: Selenazoles	941
11.22.1	Synthesis by Ring-Closure Reactions	944
11.22.1.1	By Formation of Two Se—C Bonds	944
11.22.1.1.1	Fragments C—C—N—C and Se	944
11.22.1.1.1.1	Method 1: From Vinylphosphonium Salts	944
11.22.1.2	By Formation of One Se—C and One N—C Bond	945
11.22.1.2.1	Fragments Se—C—N and C—C	945
11.22.1.2.1.1	Method 1: From Selenocarboxamides and α -Halo Carbonyl Compounds	945
11.22.1.2.1.1.1	Variation 1: 2-Alkyl- and 2-Arylselenazoles	945
11.22.1.2.1.1.2	Variation 2: Selenazol-2-amines and 2-(Alkylamino)- and 2-(Arylamino)-Substituted Selenazoles	946
11.22.1.2.1.1.3	Variation 3: 2-Hydrazinoselenazoles and Derivatives	949
11.22.1.2.1.1.4	Variation 4: <i>N</i> -Alkylimino- and <i>N</i> -Arylimino-2,3-dihydroselenazol-3-amines	950
11.22.1.2.1.1.5	Variation 5: Bis(selenazol-2-yl)methane and Oxazol-2-yl(selenazol-2-yl)methane Derivatives	952
11.22.1.2.1.1.6	Variation 6: 2-(2-Furyl)-, 2-(2-Thienyl)-, 2-(Selenophen-2-yl)-, 2-(2-Pyridyl)-, and <i>D</i> -Ribofuranosylselenazoles	952
11.22.1.2.1.1.7	Variation 7: Selenazolidines from Selenoureas and α -Halo Carboxylic Acids	953
11.22.1.2.1.1.8	Variation 8: 4,5-Dihydropyrrolo[3,2- <i>e</i>]benzoselenazoles and 4,5-Dihydroselenazolo[4,5- <i>f</i>]quinolines	954
11.22.1.2.1.1.9	Variation 9: 5-Fluoroselenazoles	955
11.22.1.2.1.1.10	Variation 10: 3-(Selenazol-4-yl)alanine	955
11.22.1.2.1.2	Method 2: Using Hypervalent Iodine Compounds	956
11.22.1.2.1.3	Method 3: From Selenocarboxamides and 2-Chlorooxiranes	956
11.22.1.2.1.4	Method 4: From Selenoureas and 3,4,6,7-Tetrachloro-2,3-dihydro-2,5-benzofurandiol	957
11.22.1.2.1.5	Method 5: From α -Bromoacyl Chlorides and Selenocarboxamides	957
11.22.1.2.1.6	Method 6: From Hydrazonoxy Bromides and Selenoureas	958
11.22.1.2.2	Fragments C—C—N and C—Se	958
11.22.1.2.2.1	Method 1: From Aziridine Derivatives and Carbon Diselenide	958
11.22.1.2.3	Fragments N—C—C—Se and C	959

11.22.1.2.3.1	Method 1: 2-Aryl-4,9-dioxonaphtho[2,3- <i>d</i>]selenazoles from Sodium Salts of 2-Amino-3-selanylaphtho-1,4-quinone	959
11.22.1.3	By Formation of One Se—C and One C—C Bond	960
11.22.1.3.1	Fragments C—N—C—Se and C	960
11.22.1.3.1.1	Method 1: From 1,1-Disubstituted 3-Acylselenoureas and Active Halomethylene Compounds	960
11.22.1.3.1.2	Method 2: From <i>N</i> -Selenoacylamidines or Morpholino(selenocarbonyl)amidines and Active Halomethylene Compounds	960
11.22.1.3.1.3	Method 3: From Selenoacylamidines and Sulfoxonium Ylides	961
11.22.1.3.2	Fragments C—N—C and C—Se	962
11.22.1.3.2.1	Method 1: From Isoselenocyanates and α -Lithiated Isocyanides	962
11.22.1.4	By Formation of One Se—C Bond	962
11.22.1.4.1	Fragment Se—C—N—C—C	962
11.22.1.4.1.1	Method 1: Oxidation of <i>N</i> -[1-(Dimethylamino)ethylidene]benzenecarboseleoamide	962
11.22.1.5	By Formation of One N—C Bond	963
11.22.1.5.1	Fragment N—C—Se—C—C	963
11.22.1.5.1.1	Method 1: From 1-Phenyl-2-(selenocyanato)ethanone	963
11.22.2	Synthesis by Ring Transformation	963
11.22.2.1	Method 1: From 1,3,4-Selenadiazines	963
11.22.2.1.1	Variation 1: 2-Imino-2,3-dihydroseleazol-3-amine Derivatives	963
11.22.2.1.2	Variation 2: Acetophenone (4-Phenyl-selenazol-2-yl)hydrazone	964
11.22.3	Synthesis by Substituent Modification	965
11.22.3.1	Addition of Organic Groups	965
11.22.3.1.1	Method 1: <i>N</i> -Alkylation: 3-Alkylselenazolium Salts	965
11.22.3.1.2	Method 2: Alkylation under Cyclization	966
11.22.3.1.2.1	Variation 1: Cyclization of Selenazol-2-amine with Dimethyl Acetylenedicarboxylate or Ethyl Propiolate	966
11.22.3.1.2.2	Variation 2: Cyclization of 2-Hydrazinoselenazoles with Ortho Esters	966
11.22.3.1.2.3	Variation 3: Cyclization of Selenazol-2-amines with α -Halo Carbonyl Compounds	967
11.22.3.2	Substitution of Existing Substituents	968
11.22.3.2.1	Of Hydrogen	968
11.22.3.2.1.1	Method 1: Nitration	968
11.22.3.2.1.2	Method 2: Sulfonation	968
11.22.3.2.1.3	Method 3: Halogenation	968
11.22.3.2.1.4	Method 4: Diazo Coupling	969
11.22.3.2.1.5	Method 5: Condensation of Selenazoles with Aldehydes or 4-Nitrosodialkylanilines	970
11.22.3.2.1.6	Method 6: Oxidation of Alkylidene Derivatives of 2-Hydrazinoselenazoles to Quinonoid Dyes	971

11.22.3.2.1.7	Method 7: Substitution of Hydrogen by Cyclization	973
11.22.3.2.2	Substitution of Carbon Functionalities	973
11.22.3.2.2.1	Method 1: Oxidation and Decarboxylation of 2,4-Dimethylselenazoles	973
11.22.3.2.2.2	Method 2: Acid Cleavage of Selenazol-2-yl Ketones to 2-Unsubstituted Selenazoles	973
11.22.3.2.3	Exchange of Heteroatoms	974
11.22.3.2.3.1	Method 1: Oxidation of 4-Aryl-2-hydrazinoselenazoles to 2-Unsubstituted Selenazoles	974
11.22.3.2.3.2	Method 2: Alkaline Splitting of (2,5-Diphenylselenazol-4-yl)phosphonium Perchlorate	974
11.22.3.2.3.3	Method 3: By Diazotization of Selenazol-2-amines	975
11.22.3.2.3.3.1	Variation 1: 4,5-Diphenylselenazol-2-ol	975
11.22.3.2.3.3.2	Variation 2: 2-Fluoroselenazoles and Selenazol-2-ols	975
11.22.3.2.3.4	Method 4: 2-Chloro-4,5-diphenylselenazole	975
11.22.3.2.3.5	Method 5: Nucleophilic Substitution of (5-Chloroselenazol-4-yl)triphenylphosphonium Perchlorates	976
11.22.3.2.3.6	Method 6: Aromatization by Addition of Iodomethane and Elimination of Trimethylamine	976
11.22.3.3	Modification of Substituents in the Side Chain	977
11.22.3.3.1	Method 1: By Oxidation of Substituted 2-Benzylselenazoles	977
11.22.3.3.2	Method 2: By Oxidation of 2-Hydrazinoselenazoles	978
11.22.3.3.3	Method 3: By Oxidation of 2-Aryl-4-(hydroxymethyl)selenazoles	979
11.22.3.3.4	Method 4: By Oxidation of 4-(Chloromethyl)selenazoles with Hexamethylenetetramine	979
11.22.3.3.5	Method 5: By Reaction of C—H Acidic Compounds with Aldehydes	980
11.22.3.3.6	Method 6: By Reaction with Ethyl Azidoacetate	980
11.22.3.3.7	Method 7: By Reaction of 2-Substituted 4-Formylselenazoles with Diazomethane	980
11.22.3.3.8	Method 8: By Reaction of 4-(Chloromethyl)- and 4-(Iodomethyl)selenazoles	981
11.22.3.3.9	Method 9: By Hydrolysis of 5-(Phthalimidoethyl)selenazol-2-amine	983
11.22.3.3.10	Method 10: By Reaction of Selenazol-2-ylacetonitriles with Sodium Nitrite	983
11.22.3.3.11	Method 11: By Hydrolysis of 2-(Benzoylamino)selenazoles	984
11.22.3.3.12	Method 12: By Hydrolysis of 2-(Isopropylidenehydrazino)selenazoles	985
11.22.3.4	Rearrangement Reactions in the Side Chain	985
11.22.3.4.1	Method 1: Beckmann Rearrangement of Aryl Selenazol-2-yl Ketoximes	985
11.23	Product Class 23: Annulated Selenazole Compounds W.-D. Pfeiffer	
11.23	Product Class 23: Annulated Selenazole Compounds	991
11.23.1	Synthesis by Ring-Closure Reactions	991
11.23.1.1	By Formation of Two Se—C Bonds	991
11.23.1.1.1	Fragments Arene—N—C and Se	991

11.23.1.1.1.1	Method 1: From <i>N</i> -Phenylbenzamide and Selenium	991
11.23.1.1.1.2	Method 2: From (2-Lithio-1 <i>H</i> -pyrrol-1-ylethynyl)lithium and Selenium ..	991
11.23.1.2	By Formation of One Se—C and One N—C Bond	992
11.23.1.2.1	Fragments N—Hetarene and Se—C	992
11.23.1.2.1.1	Method 1: From a Selenocarboxylate and an Arylamine	992
11.23.1.2.2	Fragments N—Arene—Se and C	993
11.23.1.2.2.1	Method 2: Synthesis by Ring Closure from 2-Seleno-Substituted Anilines and a Carboxylic Derivative	993
11.23.1.3	By Formation of One Se—C Bond	994
11.23.1.3.1	Fragment Arene—N—C—Se	994
11.23.1.3.1.1	Method 1: Benzoselenazoles by the Oxidation of <i>N</i> -Phenylselenoureas ..	994
11.23.1.4	By Formation of One N—C Bond	995
11.23.1.4.1	Fragment N—Arene—Se—C	995
11.23.1.4.1.1	Method 1: From 2-Aminoaryl Selenocyanates	995
11.23.1.4.2	Fragment N—C—Se—C—C	996
11.23.1.4.2.1	Method 1: Mesoionic Selenazolium Compounds from Diselenides	996
11.23.1.4.3	Fragment N—Hetarene—Se—C	997
11.23.1.4.3.1	Method 1: Thieno[3,2- <i>d</i>]selenazoles from Thienyl Selenoates	997
11.23.2	Synthesis by Substituent Modification	997
11.23.2.1	Method 1: <i>N</i> -Alkylation	997
11.23.2.2	Method 2: Reactions of Benzoselenazolium Salts	997
11.23.2.3	Method 3: Substitution of the 2-Methylsulfanyl Group of 3-Methyl-2- (methylsulfanyl)benzoselenazolium Iodide by Formic Hydrazide	998
11.23.2.4	Modification of Existing Substituents	998
11.23.2.4.1	Method 1: Condensation of 2-Methylbenzoselenazole with Aldehydes ..	998
11.23.2.4.2	Method 2: Oxidation of 2-Methyl-1,3-benzoselenazole	998
11.24	Product Class 24: 1,3-Selenaphospholes D. Gudat	
11.24	Product Class 24: 1,3-Selenaphospholes	1001
11.24.1	Synthesis by Ring-Closure Reactions	1001
11.24.1.1	By Formation of Two P—C Bonds	1001
11.24.1.1.1	Fragments C—Se—C—C and P	1001
11.24.1.1.1.1	Method 1: From Ketene <i>N,Se</i> -Acetals and Phosphorus Trihalides	1001
11.24.2	Synthesis by Ring Transformation	1002
11.24.2.1	Method 1: From 1,2,3-Selenadiazoles and Phosphaalkynes	1002

11.25	Product Class 25: Isotellurazoles, and Annulated Isotellurazole and Tellurazole Compounds W.-D. Pfeiffer	
11.25	Product Class 25: Isotellurazoles, and Annulated Isotellurazole and Tellurazole Compounds	1005
11.25.1	Product Subclass 1: Isotellurazoles (1,2-Tellurazoles)	1005
11.25.1.1	Synthesis by Ring-Closure Reactions	1005
11.25.1.1.1	By Formation of One Te—N, One Te—C, and One N—C Bond	1005
11.25.1.1.1.1	Method 1: From Acetylenic Ketones or Aldehydes and Hydroxylamine O-Sulfonic Acid	1005
11.25.1.1.2	By Formation One Te—N and One N—C Bond	1006
11.25.1.1.2.1	Fragments Te—C—C—C and N	1006
11.25.1.1.2.1.1	Method 1: Ring Closure of a 2-Tellurium-Functionalized Enal with a Nitrogen Source	1006
11.25.1.1.2.1.1.1	Variation 1: From 3-Oxoprop-1-ene-1-tellurenyl Bromides and Ammonia	1006
11.25.1.1.2.1.1.2	Variation 2: From 3-[Dibromo(methyl)- λ^4 -tellanyl]prop-2-enals and Ammonia	1007
11.25.1.1.2.1.1.3	Variation 3: From 3-[Dibromo(methyl)- λ^4 -tellanyl]prop-2-enals and Arylamines	1007
11.25.2	Product Subclass 2: 1,2-Benzisotellurazoles	1008
11.25.2.1	Synthesis by Ring-Closure Reactions	1008
11.25.2.1.1	By Formation of One Te—N and One N—C Bond	1008
11.25.2.1.1.1	Fragments Te—Arene—C and N	1008
11.25.2.1.1.1.1	Method 1: Ring Closure of a 2-Tellurium-Functionalized Arylaldehyde with a Nitrogen Source	1008
11.25.2.1.2	By Formation of One Te—N Bond	1009
11.25.2.1.2.1	Fragment Te—Arene—C—N	1009
11.25.2.1.2.1.1	Method 1: From 2-(Butyltellanyl)benzaldehyde	1009
11.25.2.1.2.1.2	Method 2: Cyclization of 2-(Iminomethyl)benzenetellurenyl Chlorides	1009
11.25.2.2	Modification of 1,2-Benzisotellurazole Iodide by Alkylation	1010
11.25.3	Product Subclass 3: Arenotellurazoles	1010
11.25.3.1	Synthesis by Ring-Closure Reactions	1010
11.25.3.1.1	By Formation of One Te—C Bond	1010
11.25.3.1.1.1	Fragment Te—Arene—N—C	1010
11.25.3.1.1.1.1	Method 1: Cyclization of 2-(Alkyltellanyl)- <i>N</i> -acylanilines	1010
11.25.3.1.1.1.1.1	Variation 1: From 1-(Ethyltellanyl)-2-nitrobenzene	1010
11.25.3.1.1.1.1.2	Variation 2: From Bis(2-aminophenyl)ditellurides	1011
11.25.3.1.1.1.2	Method 2: Cyclization of 2-Tellanyl- <i>N</i> -acylanilines	1012
11.25.3.1.1.1.2.1	Variation 1: From 1-Phenyl-2-[2-(trichloro- λ^4 -tellanyl)phenyl]diazene	1012
11.25.3.1.1.1.2.2	Variation 2: From 2-(<i>N</i> -Acetylaminoaryl)mercury Chlorides	1013

11.25.3.1.1.1.2.3	Variation 3: From Ditellurides via 1,2,5-Oxatellurazoles	1014
11.25.3.2	Synthesis by Modification	1016
11.25.3.2.1	Addition Reactions	1016
11.25.3.2.1.1	Method 1: Reaction with Halogens	1016
11.25.3.2.1.2	Method 2: Formation of Complexes	1016
11.25.3.2.1.3	Method 3: Protonation	1016
11.25.3.2.1.4	Method 4: Methylation of 2-Phenylbenzotellurazole	1017
11.25.3.2.1.5	Method 5: Reaction of the Methyl Group in 2-Methylbenzotellurazolium Salts	1017
 Keyword Index		1021
Author Index		1093
Abbreviations		1155