

**Volume 15:
Six-Membered Heterenes
with One Nitrogen or Phosphorus Atom**

	Preface	V
	Volume Editor's Preface	VII
	Table of Contents	XI
	Introduction	
	D. StC. Black	1
15.1	Product Class 1: Pyridines	
	D. Spitzner	11
15.2	Product Class 2: Pyridinones and Related Systems	
	P. A. Keller	285
15.3	Product Class 3: Quinolines	
	R. D. Larsen and D. Cai	389
15.4	Product Class 4: Quinolinones	
	R. D. Larsen	551
15.5	Product Class 5: Isoquinolines	
	M. Álvarez and J. A. Joule	661
15.6	Product Class 6: Isoquinolinones	
	M. Álvarez and J. A. Joule	839
15.7	Product Class 7: Quinolizinium Salts and Benzo Analogues	
	H. Ihmels	907
15.8	Product Class 8: Naphthyridines	
	P.-W. Phuan and M. C. Kozlowski	947
15.9	Product Class 9: Acridines	
	R. H. Prager and C. M. Williams	987
15.10	Product Class 10: Acridin-9(10<i>H</i>)-ones and Related Systems	
	R. H. Prager and C. M. Williams	1029
15.11	Product Class 11: Phenanthridines	
	P. A. Keller	1065
15.12	Product Class 12: Phenanthridinones and Related Systems	
	P. A. Keller	1089
15.13	Product Class 13: 1λ^3-Phosphinines	
	F. Mathey and P. Le Floch	1097

15.14	Product Class 14: $1\lambda^5$-Phosphinines	
	R. Streubel	1157
15.15	Product Class 15: Benzo-Fused And Other Annulated Phosphinines	
	U. Bergsträsser	1181
	Keyword Index	1191
	Author Index	1249
	Abbreviations	1315

Table of Contents

Introduction

D. StC. Black

Introduction	1
---------------------------	---

15.1 Product Class 1: Pyridines

D. Spitzner

15.1	Product Class 1: Pyridines	11
15.1.1	Product Subclass 1: Pyridines	11
15.1.1.1	Synthesis by Ring-Closure Reactions	13
15.1.1.1.1	By Formation of Two N—C and Two C—C Bonds	13
15.1.1.1.1.1	Fragments C—C, C—C, N, and C	13
15.1.1.1.1.1.1	Method 1: From 1,3-Diketones or β -Oxo Esters, Aldehydes, and Ammonia, with Subsequent Oxidation (Hantzsch Pyridine Synthesis)	13
15.1.1.1.1.1.1.1	Variation 1: An Abnormal Ring Closure	14
15.1.1.1.1.1.1.2	Variation 2: From β -Oxo Cyanides, Aldehydes, and Ammonium Salts	15
15.1.1.1.1.1.1.3	Variation 3: Alkyl- and Arylpyridines from Aldehydes and Ammonium Salts (Tschitschibabin Synthesis)	15
15.1.1.1.1.1.1.4	Variation 4: From Ketones, Iminium Compounds, and Ammonium Salts ...	16
15.1.1.1.2	By Formation of One N—C Bond and Three C—C Bonds	16
15.1.1.1.2.1	Fragments N—C, C—C, C, and C	16
15.1.1.1.2.1.1	Method 1: From Aldehydes, Nitriles, and Methylphosphonates	17
15.1.1.1.3	By Formation of Two N—C Bonds and One C—C Bond	18
15.1.1.1.3.1	Fragments C—C—C, C—C, and N	18
15.1.1.1.3.1.1	Method 1: From Enamines, α,β -Unsaturated Carbonyl Compounds, and Hydroxylamine or Ammonium Salts	18
15.1.1.1.3.1.1.1	Variation 1: From Enamines, α,β -Unsaturated Aldehydes, and Hydroxylamine	18
15.1.1.1.3.1.1.2	Variation 2: From Acyl Enamines, Ketones, and Ammonium Salts	18
15.1.1.1.3.1.1.3	Variation 3: From Ketones, α,β -Unsaturated Aldehydes or Ketones, and Ammonium Salts	19
15.1.1.1.3.1.2	Method 2: From Acylketene Dithioacetals, Ketones, and Ammonium Salts	21
15.1.1.1.3.1.3	Method 3: From Sulfonium Salts or Sulfur Ylides, α,β -Unsaturated Carbonyl Compounds, and Ammonium Salts	22
15.1.1.1.3.1.4	Method 4: From 1-(2-Oxoalkyl)pyridinium (or Quinolinium/Isoquinolinium) Compounds, α,β -Unsaturated Carbonyl Compounds, and Ammonium Salts	23
15.1.1.1.3.1.4.1	Variation 1: Cyclization of Ketones with Mannich Salts in the Presence of Ammonium Salts	24
15.1.1.1.3.1.5	Method 5: From Propynols, Enamines, and Ammonium Salts by a Four-Component, One-Pot Procedure	25

15.1.1.1.3.1.5.1	Variation 1: From Propynols, Ketones, and Ammonium Salts	26
15.1.1.1.4	By Formation of One N—C Bond and Two C—C Bonds	26
15.1.1.1.4.1	Fragments N—C, C—C, and C—C	26
15.1.1.1.4.1.1	Method 1: From Nitriles and Alkynes	26
15.1.1.1.4.1.1.1	Variation 1: From Nitriles and Alkynes by Catalyzed [2+2+2] Cycloaddition	27
15.1.1.1.4.1.2	Method 2: From Hydrazones and Alkynes	29
15.1.1.1.4.1.3	Method 3: From Imines and Aldehydes	29
15.1.1.1.4.1.4	Method 4: From 1-Substituted Phthalazines and Ynamines	30
15.1.1.1.4.2	Fragments N—C—C, C—C, and C	31
15.1.1.1.4.2.1	Method 1: From Malononitrile, Aldehydes, and Nucleophiles	31
15.1.1.1.4.2.1.1	Variation 1: From Malononitrile, Aldehydes, Ketones, and Ammonium Salts	31
15.1.1.1.4.2.1.2	Variation 2: From Malononitrile and 1,1-Diethoxy- <i>N,N</i> -dimethylalkanamines or Orthoformates	32
15.1.1.1.4.2.2	Method 2: From Lithiated Dimethylhydrazones, α,β -Unsaturated Ketones, and α -Oxo Cyanides	33
15.1.1.1.4.2.3	Method 3: From Enamines, Aldehydes, and 1,3-Dicarbonyl Compounds (Hantzsch Variation)	34
15.1.1.1.4.2.4	Method 4: From [(Diphenylphosphoryl)imino]ethanes and Aromatic Aldehydes	35
15.1.1.1.4.2.5	Method 5: From [(Triphenylphosphoranylidene)amino]alkenes, Isocyanates, and Enamines	35
15.1.1.1.4.3	Fragments C—C—C, N—C, and C	36
15.1.1.1.4.3.1	Method 1: From Nitriles and Methallyl Dianions	36
15.1.1.1.4.3.2	Method 2: From Acetonitrile and α,β -Unsaturated Carbonyl Compounds	36
15.1.1.1.4.3.3	Method 3: From Nitriles, α,β -Unsaturated Ketones, and Carbanions	37
15.1.1.1.4.3.4	Method 4: From Nitriles and Acylketene <i>N,S</i> -Acetals	37
15.1.1.1.4.3.5	Method 5: From Nitriles, Lithiated Diethyl Methylphosphonates, and α,β -Unsaturated Ketones	38
15.1.1.1.5	By Formation of Three C—C Bonds	38
15.1.1.1.5.1	Fragments C—N—C, C—C, and C	38
15.1.1.1.5.1.1	Method 1: Pyridin-3-ols from (2-Azaallenyl)pentacarbonylchromium Complexes, Alkynes, and Carbon Monoxide	38
15.1.1.1.6	By Formation of Two N—C Bonds	39
15.1.1.1.6.1	Fragments C—C—C—C and N	39
15.1.1.1.6.1.1	Method 1: From 1,5-Dioxo Compounds and Hydroxylamine	39
15.1.1.1.6.1.1.1	Variation 1: From Partially or Fully Protected 1,5-Dioxo Compounds and Hydroxylamine	40
15.1.1.1.6.1.1.2	Variation 2: From Penta-2,4-dienals and Hydroxylamine	41
15.1.1.1.6.1.2	Method 2: From 1,5-Dioxo Compounds and Ammonia or Ammonium Salts	42
15.1.1.1.6.1.2.1	Variation 1: From Hydroxy- or Halo-1,5-dioxo Compounds and Ammonia or Ammonium Salts	42
15.1.1.1.6.1.2.2	Variation 2: From α,β -Unsaturated 1,5-Dioxo Compounds (or Partially Protected Compounds) and Ammonia or Ammonium Salts	43
15.1.1.1.6.1.3	Method 3: From Pentamethinammonium Salts and Ammonium Salts	44

15.1.1.1.7	By Formation of One N—C and One C—C Bond	44
15.1.1.1.7.1	Fragments N—C—C—C and C—C	44
15.1.1.1.7.1.1	Method 1: From (Alkoxyethylene)malononitriles, Enamines, and Ammonia	44
15.1.1.1.7.1.1.1	Variation 1: From Arylidene malononitriles and Ketones	45
15.1.1.1.7.1.2	Method 2: From Malononitrile and Alkyl Halides under Lewis Acid Catalysis	46
15.1.1.1.7.1.2.1	Variation 1: From <i>N</i> -Alkylcyanoacetamides or <i>N,N</i> -Dialkylcyanoacetamides in the Presence of Phosphoryl Chloride	47
15.1.1.1.7.1.3	Method 3: From 3-Aminopropenals and Ketones or 1,3-Dicarbonyl Compounds	48
15.1.1.1.7.1.4	Method 4: From 1-Aza-1,3-dienes and Alkenes or Alkynes	48
15.1.1.1.7.1.4.1	Variation 1: From Alk-2-enal Acylhydrazones and Alkynes or <i>N</i> -Phenylmaleimide	49
15.1.1.1.7.1.4.2	Variation 2: From Alk-2-enal Dimethylhydrazones by $[4\pi + 2\pi]$ Cycloaddition	50
15.1.1.1.7.1.4.3	Variation 3: Cyclization of an Intermediate 1-(Dimethylamino)-1-aza-1,3,5-triene	51
15.1.1.1.7.1.4.4	Variation 4: From Alk-2-enal <i>O</i> -Alkyloximes and Acylalkynes	52
15.1.1.1.7.1.4.5	Variation 5: From Alk-2-enal <i>N</i> -Alkylimines and Enamines	52
15.1.1.1.7.1.4.6	Variation 6: From Alk-2-enal Oximes or <i>N</i> -Alkylimines and Carbonyl Compounds	53
15.1.1.1.7.1.4.7	Variation 7: Palladium-Catalyzed Iminoannulation of Alkynes	53
15.1.1.1.7.1.4.8	Variation 8: From 4-Amino-1-aza-1,3-dienes and Alkynes	54
15.1.1.1.7.1.4.9	Variation 9: From 3-[(Triphenylphosphoranylidene)amino]propenals and Acetylenic Esters	54
15.1.1.1.7.1.4.10	Variation 10: From Amidines (2-Amino-1-aza-1,3-dienes) and Electron-Poor Alkynes	55
15.1.1.1.7.1.4.11	Variation 11: From Coordinated 1-Aza-1,3-dienes (Fischer Carbene Complexes) and Alkynes	55
15.1.1.1.7.1.5	Method 5: From Acrylonitrile and 2,2-Dichlorinated Aldehydes	57
15.1.1.1.7.1.6	Method 6: From β -Oxo Carbonyl Compounds and Enaminonitriles	57
15.1.1.1.7.2	Fragments C—C—C—C and N—C	58
15.1.1.1.7.2.1	Method 1: From Alkylidenemalononitriles and Activated Nitriles	58
15.1.1.1.7.2.2	Method 2: From Lithiated α,β -Unsaturated Aldimines and Nitriles	59
15.1.1.1.7.2.3	Method 3: From Bicyclic Acetals and Nitriles	60
15.1.1.1.7.2.4	Method 4: From Hetero-[4 + 2] Cycloaddition of Dienes with Nitriles	60
15.1.1.1.7.2.4.1	Variation 1: From Push–Pull “Captodative” Activated 1,3-Dienes and Nitriles	61
15.1.1.1.7.2.4.2	Variation 2: From 1-Alkoxybuta-1,3-dienes and Activated Nitriles	61
15.1.1.1.7.2.4.3	Variation 3: Intramolecular Hetero-[4 + 2] Cycloaddition of Dienes with (Hydroxyimino)malonates	62
15.1.1.1.7.2.4.4	Variation 4: Hetero-[4 + 2] Cycloaddition of Dienes with <i>N</i> -(Sulfonyloxy)imines	62
15.1.1.1.7.3	Fragments N—C—C and C—C—C	63
15.1.1.1.7.3.1	Method 1: From α -Halogenated Cyanides and α,β -Unsaturated Aldehydes	63
15.1.1.1.7.3.1.1	Variation 1: From α -(1 <i>H</i> -1,2,3-Benzotriazol-1-yl)-Substituted Cyanides and α,β -Unsaturated Ketones	64

15.1.1.1.7.3.2	Method 2:	From Malonamide Derivatives and Unsaturated Compounds	64
15.1.1.1.7.3.2.1	Variation 1:	From Malononitrile and Unsaturated Compounds in the Presence of a Nucleophile	65
15.1.1.1.7.3.3	Method 3:	From Enamines and α,β -Unsaturated Cyanides	67
15.1.1.1.7.3.4	Method 4:	From Enamines or Imines and [3-(Dimethylamino)allylidene]-ammonium Salts	68
15.1.1.1.7.3.5	Method 5:	From Acyl Enamines and α,β -Unsaturated Carbonyl Compounds	68
15.1.1.1.7.3.5.1	Variation 1:	From Functionalized Enamines and Alkynes (Bohlmann–Rahtz Synthesis)	68
15.1.1.1.7.3.5.2	Variation 2:	From Functionalized Enamines and α,β -Unsaturated Carbonyl Compounds	69
15.1.1.1.7.3.5.3	Variation 3:	From Acyl Enamines and Ethene-1,1-diamines	71
15.1.1.1.7.3.5.4	Variation 4:	From Enaminonitriles and 1,3-Diketone Enol Ethers (Guareschi–Thorpe-Type Reaction)	72
15.1.1.1.7.3.5.5	Variation 5:	From <i>N,N</i> -Bis(silylated) Enamines and α,β -Unsaturated Carbonyl Compounds	73
15.1.1.1.7.3.5.6	Variation 6:	From 3-Aminobut-2-enoates or Acylmethyl Imines and 1,3,3-Triethoxyprop-1-ene	73
15.1.1.1.7.3.5.7	Variation 7:	From 3-Oxoalkanimidamides or Acyl Enamines and 1,1,3,3-Tetraalkoxypropanes	74
15.1.1.1.7.3.5.8	Variation 8:	From 3-Aminobut-2-enoates and Mannich Salts	74
15.1.1.1.7.3.5.9	Variation 9:	From Enaminonitriles and α,β -Unsaturated Carbonyl Compounds	75
15.1.1.1.7.3.6	Method 6:	From Enaminonitriles and Alkynyl Ortho Amides	75
15.1.1.1.7.3.7	Method 7:	From Lithiated Imines and α,β -Unsaturated Carbonyl Compounds	76
15.1.1.1.7.3.7.1	Variation 1:	From <i>N-tert</i> -Butyl Ketimines and α,β -Unsaturated Cyanides	76
15.1.1.1.7.3.8	Method 8:	From (Vinylimino)phosphoranes and α,β -Unsaturated Carbonyl Compounds (Aza-Wittig Reaction)	77
15.1.1.1.7.3.8.1	Variation 1:	From [(Diphenylphosphoryl)imino]alkanes and α,β -Unsaturated Aldehydes or Ketones	78
15.1.1.1.7.4	Fragments N—C—C—C and C		79
15.1.1.1.7.4.1	Method 1:	From Iminophosphoranes and Isocyanates by a Domino Aza-Wittig Electrocyclization Reaction	79
15.1.1.1.7.4.2	Method 2:	From Alkylidenemalononitriles	80
15.1.1.1.7.4.3	Method 3:	From 1-Amino-4-nitro-1,3-dienes and Orthoformates or Acetic Anhydride	80
15.1.1.1.8	By Formation of Two C—C Bonds		81
15.1.1.1.8.1	Fragments C—N—C—C and C—C		81
15.1.1.1.8.1.1	Method 1:	From <i>N</i> -Vinylcarbodiimides and Alkynes	81
15.1.1.1.8.1.1.1	Variation 1:	From <i>N</i> -Vinylcarbodiimides and Enamines	82
15.1.1.1.8.1.1.2	Variation 2:	From <i>N</i> -Vinylcarbodiimides and Ynamines	82
15.1.1.1.8.1.2	Method 2:	From 2-Azabuta-1,3-dienes and Alkynes	83
15.1.1.1.8.1.2.1	Variation 1:	From 2-Aza-1,3-dienes and Enamines	83

15.1.1.1.8.1.2.2	Variation 2: Intramolecular Reaction of Neutral 2-Aza-1,3-dienes with Alkynes or Alkenes	84
15.1.1.1.8.1.3	Method 3: From Vinyl Isocyanides and Alkynes	85
15.1.1.1.8.2	Fragments C—N—C and C—C—C	85
15.1.1.1.8.2.1	Method 1: From 2-Azaallyl Anions and 2-(Iodomethyl)-3-(trimethylsilyl)prop-1-ene	85
15.1.1.1.8.2.2	Method 2: From Nitrile Ylides and Methylene-cyclopropenes	85
15.1.1.1.8.2.3	Method 3: From 2-Azapropenylium Salts and Enamines	86
15.1.1.1.8.3	Fragments C—C—N—C—C and C	86
15.1.1.1.8.3.1	Method 1: From 3-Azapenta-1,3-dienes by Carbonylation	86
15.1.1.1.8.3.2	Method 2: From 3-Azapenta-1,3-dienes and Aldehydes or Imines	87
15.1.1.1.9	By Formation of One N—C Bond	87
15.1.1.1.9.1	Fragment N—C—C—C—C	87
15.1.1.1.9.1.1	Method 1: Cyclization of 2-Substituted Pentanedinitriles with Nucleophiles	87
15.1.1.1.9.1.1.1	Variation 1: Cyclization of 1,1,3,3-Tetracyanopropenide Salts with Nucleophiles	88
15.1.1.1.9.1.1.2	Variation 2: Cyclization of Pentanedinitriles with Carbon Nucleophiles	89
15.1.1.1.9.1.2	Method 2: From Halogenated 5-Oxoalkanenitriles (or Their Acetals)	90
15.1.1.1.9.1.2.1	Variation 1: Cyclization of 3-Hydroxy-5,5-bis(methylsulfanyl)pent-4-enitriles with Acid	90
15.1.1.1.9.1.2.2	Variation 2: Cyclization of 4-Oxopentane-1,1,2,2-tetracarbonitrile with Nucleophiles	91
15.1.1.1.9.1.3	Method 3: From Push–Pull Alka-2,4-dienenitriles	91
15.1.1.1.9.1.4	Method 4: From Alka-2,4-dienal Oximes (1-Hydroxy-1-azahexa-1,3,5-trienes)	92
15.1.1.1.9.1.4.1	Variation 1: Selenium-Induced Cyclization of Alk-4-enone Oximes	94
15.1.1.1.9.1.4.2	Variation 2: Cyclization of 5-Aminoalk-4-enal Oximes	94
15.1.1.1.9.1.5	Method 5: From Pentane-1,5-dione Mono(dimethylhydrazone) Equivalents	95
15.1.1.1.9.1.5.1	Variation 1: From Pentane-1,5-dione Mono- or Dioximes	95
15.1.1.1.9.1.5.2	Variation 2: From Oxo Imidates	96
15.1.1.1.9.1.6	Method 6: From Tri-O-acetylated Sugar Diamides	97
15.1.1.1.9.1.7	Method 7: From 5-(Hydroxyimino)alkanenitriles	97
15.1.1.1.9.1.8	Method 8: From 4-Amino-1-aza-1,3,5-trienes (3-Amino-2,4-dienone Imines)	98
15.1.1.1.9.1.8.1	Variation 1: Thermal Condensation of Alka-2,4-dienal Dimethylhydrazones [1-(Dimethylamino)-1-aza-1,3,5-trienes]	98
15.1.1.1.9.1.9	Method 9: Cyclization of Iminoalkenynes in Polyphosphoric Acid	99
15.1.1.1.10	By Formation of One C—C Bond	99
15.1.1.1.10.1	Fragment C—N—C—C—C	99
15.1.1.1.10.1.1	Method 1: From <i>N</i> -(Buta-1,3-dienyl)carbodiimides	99
15.1.1.1.10.1.2	Method 2: From <i>N</i> -Acylbuta-1,3-dien-1-amines	100
15.1.1.1.10.1.3	Method 3: From 2-Azahexa-1,3,5-trienes	100
15.1.1.1.10.2	Fragment C—C—N—C—C	100

15.1.1.1.10.2.1	Method 1:	From <i>O</i> -Allyloximes by Thermal Rearrangement, Followed by Cyclization	100
15.1.1.1.10.2.2	Method 2:	By Amino-Claisen Rearrangement	101
15.1.1.1.10.2.3	Method 3:	From 3-Azahexa-1,3,5-trienes	101
15.1.1.1.10.2.3.1	Variation 1:	Model for Prebiotic Pyridine Synthesis	102
15.1.1.1.10.2.4	Method 4:	From 3-Azahexa-1,3-dien-5-yne	102
15.1.1.1.10.2.5	Method 5:	From {[2-(Pyrrolidin-1-yl)prop-1-enyl]imino}malononitriles	103
15.1.1.1.10.2.6	Method 6:	Oxidative Cyclization of Hydroxy Enaminones	103
15.1.1.2	Synthesis by Ring Transformation		103
15.1.1.2.1	By Ring Enlargement		103
15.1.1.2.1.1	Of Three-Membered Carbocyclic or Heterocyclic Compounds		103
15.1.1.2.1.1.1	Method 1:	From 2,2-Dichlorocyclopropanecarbaldehydes	103
15.1.1.2.1.1.1.1	Variation 1:	From Tris(isopropylsulfanyl)cyclopropenylium Perchlorate and α -Lithiated Isocyanides	104
15.1.1.2.1.1.2	Method 2:	From 2 <i>H</i> -Azirines	105
15.1.1.2.1.1.3	Method 3:	From Oxaziridines	105
15.1.1.2.1.2	Of Four-Membered Heterocyclic Compounds		106
15.1.1.2.1.2.1	Method 1:	From 2,3-Dihydroazetes	106
15.1.1.2.1.2.1.1	Variation 1:	From 2,3-Dihydroazete 1-Oxides	106
15.1.1.2.1.2.1.2	Variation 2:	From Dewar Pyridines	107
15.1.1.2.1.3	Of Five-Membered Carbocyclic or Heterocyclic Compounds		107
15.1.1.2.1.3.1	Method 1:	From Cyclopentadienones and Nitriles	107
15.1.1.2.1.3.2	Method 2:	From Furans or Dihydrofurans	108
15.1.1.2.1.3.2.1	Variation 1:	From Furfurylamines	108
15.1.1.2.1.3.2.2	Variation 2:	From 2-Acylfurans and Ammonium Salts	110
15.1.1.2.1.3.3	Method 3:	From Pyrroles	110
15.1.1.2.1.3.4	Method 4:	From Isoxazoles	111
15.1.1.2.1.3.5	Method 5:	From Oxazoles	111
15.1.1.2.1.3.5.1	Variation 1:	From Dihydrooxazolones	113
15.1.1.2.1.3.5.2	Variation 2:	From Oxazoles and Malononitrile	113
15.1.1.2.2	By Formal Exchange of Ring Members with Retention of the Ring Size		114
15.1.1.2.2.1	From Oxygen Heterocycles and a Nitrogen Source		114
15.1.1.2.2.1.1	Method 1:	From Pirylium Salts and Ammonium Salts	114
15.1.1.2.2.1.2	Method 2:	From Substituted 4 <i>H</i> -Pyrans and Ammonium Salts	115
15.1.1.2.2.1.3	Method 3:	From 2 <i>H</i> -Pyran-2-ones or 4 <i>H</i> -Pyran-4-ones and Ammonium Salts	115
15.1.1.2.2.1.4	Method 4:	From 2-Alkoxy-3,4-dihydro-2 <i>H</i> -pyrans and Hydroxylamine	116
15.1.1.2.2.2	From Sulfur Heterocycles		117
15.1.1.2.2.2.1	Method 1:	From 2-Azido-2 <i>H</i> -thiopyrans	117
15.1.1.2.2.3	From Pyridinium Compounds or Pyridinones		117
15.1.1.2.2.3.1	Method 1:	From Pyridinium Compounds by an S_N (ANRORC) Mechanism	117
15.1.1.2.2.3.2	Method 2:	From Pyridin-2(1 <i>H</i>)-ones, Ketones or Enamines, and Ammonia	118
15.1.1.2.2.4	From 1, <i>n</i> -Oxazines		119

15.1.1.2.2.4.1	Method 1:	From 6-(2-Oxoalkyl)-6 <i>H</i> -1,2-oxazines by Base-Catalyzed Rearrangement	119
15.1.1.2.2.4.2	Method 2:	Cycloaddition of 6-Alkoxy-6 <i>H</i> -1,2-oxazines with Alkynes	120
15.1.1.2.2.4.2.1	Variation 1:	From 6 <i>H</i> -1,3-Oxazin-6-ones by [4 + 2] Cycloaddition	120
15.1.1.2.2.4.2.2	Variation 2:	From 2 <i>H</i> -1,3-Oxazin-2-ones by [4 + 2] Cycloaddition	121
15.1.1.2.2.4.2.3	Variation 3:	From 2 <i>H</i> -1,4-Oxazin-2-ones by [4 + 2] Cycloaddition	121
15.1.1.2.2.5		From 1, <i>n</i> -Diazines	122
15.1.1.2.2.5.1	Method 1:	From Pyridazines by [4 + 2] Cycloaddition	122
15.1.1.2.2.5.2	Method 2:	From Pyrimidines or Pyrimidinium Compounds	123
15.1.1.2.2.5.2.1	Variation 1:	By [4 + 2] Cycloaddition	123
15.1.1.2.2.5.2.2	Variation 2:	By an S _N (ANRORC) Mechanism	124
15.1.1.2.2.5.3	Method 3:	From Pyrimidinones or Pyrimidinethiones	125
15.1.1.2.2.5.4	Method 4:	From Pyrazines or Pyrazin-2(1 <i>H</i>)-ones by [4 + 2] Cycloaddition	127
15.1.1.2.2.6		From 1, <i>n,m</i> -Triazines	127
15.1.1.2.2.6.1	Method 1:	From 1,2,3-Triazines by [4 + 2] Cycloaddition	127
15.1.1.2.2.6.2	Method 2:	From 1,2,4-Triazines by [4 + 2] Cycloaddition	128
15.1.1.2.3		By Ring Contraction	130
15.1.1.2.3.1		Of Seven-Membered Heterocycles	130
15.1.1.2.3.1.1	Method 1:	From 3 <i>H</i> -Azepines	130
15.1.1.2.3.1.2	Method 2:	From 1, <i>n</i> -Oxazepines	130
15.1.1.2.3.2		Of Eight-Membered Heterocycles	130
15.1.1.2.3.2.1	Method 1:	From Azocines	130
15.1.1.2.3.2.2	Method 2:	From 1,2-Diazocines	131
15.1.1.3		Aromatization	131
15.1.1.3.1		By Isomerization	131
15.1.1.3.1.1	Method 1:	Isomerization of 1-Acyl-4-benzylidene-1,4-dihydropyridines and 1-Amino-4-methylene-1,4-dihydropyridines	131
15.1.1.3.2		By Dehydrogenation	132
15.1.1.3.2.1	Method 1:	Oxidation (Dehydrogenation) of 1,2-, 1,4-, 3,4-, and 2,5-Dihydropyridines	132
15.1.1.3.2.1.1	Variation 1:	Oxidation of Hantzsch Esters	133
15.1.1.3.2.2	Method 2:	Dehydrogenation of Tetrahydropyridines	134
15.1.1.3.3		By Elimination from Dihydropyridines or Tetrahydropyridines	135
15.1.1.3.3.1	Method 1:	From 1-(Alkoxy-carbonyl)-1,4- or 1-(Alkoxy-carbonyl)-1,2-dihydropyridines	135
15.1.1.3.3.1.1	Variation 1:	From 1,4-Bis(trimethylsilyl)-1,4-dihydropyridine and Carbonyl Compounds	136
15.1.1.3.3.2	Method 2:	From Tetrahydropyridines	137
15.1.1.3.3.2.1	Variation 1:	From 2,3,4,5-Tetrahydropyridines by Chlorination–Elimination	138
15.1.1.3.3.3	Method 3:	From 2-Azabicyclo[2.2.2]octa-2,5-dienes	138
15.1.1.3.3.3.1	Variation 1:	From Dewar Pyridines	139
15.1.1.3.3.4	Method 4:	From Piperidine-2,6-diones	139
15.1.1.4		Synthesis by Substituent Modification	140

15.1.1.4.1	Substitution of Existing Substituents	140
15.1.1.4.1.1	Of Hydrogen	140
15.1.1.4.1.1.1	By Deuterium	140
15.1.1.4.1.1.1.1	Method 1: Base-Induced Hydrogen–Deuterium Exchange	140
15.1.1.4.1.1.1.2	Method 2: Hydrogen–Deuterium Exchange Using Heterogeneous Catalysis	140
15.1.1.4.1.1.2	By Metals	141
15.1.1.4.1.1.2.1	Method 1: By Alkali Metals	141
15.1.1.4.1.1.2.2	Method 2: By Mercury	142
15.1.1.4.1.1.2.3	Method 3: By Silicon	142
15.1.1.4.1.1.2.4	Method 4: By Tin	143
15.1.1.4.1.1.3	By a Carbon Functionality	144
15.1.1.4.1.1.3.1	Method 1: Radical Aminocarbonylation, Alkoxy carbonylation (Minisci Method), and Acylation	144
15.1.1.4.1.1.3.2	Method 2: Acylation via Metalation	144
15.1.1.4.1.1.3.3	Method 3: Alkylation and Arylation via Metalation	145
15.1.1.4.1.1.3.3.1	Variation 1: Hydroxyalkylation	145
15.1.1.4.1.1.3.3.2	Variation 2: Reaction with Organomagnesium Compounds (Deprotonation vs Addition)	146
15.1.1.4.1.1.3.3.3	Variation 3: Radical Alkylations and Arylations	146
15.1.1.4.1.1.3.3.4	Variation 4: Dihalomethylation of Nitropyridines (Vicarious Nucleophilic Substitution)	147
15.1.1.4.1.1.3.4	Method 4: Intramolecular Ring Closure via Hetarynes	148
15.1.1.4.1.1.4	By a Halogen	148
15.1.1.4.1.1.4.1	Method 1: Direct Halogenation	148
15.1.1.4.1.1.4.2	Method 2: Halogenation via Lithiation	150
15.1.1.4.1.1.5	By Oxygen	151
15.1.1.4.1.1.5.1	Method 1: Oxidation with Acetyl Hypofluorite	151
15.1.1.4.1.1.5.2	Method 2: Oxidation via Boronates	151
15.1.1.4.1.1.6	By Sulfur or Selenium	152
15.1.1.4.1.1.6.1	Method 1: Sulfonylation	152
15.1.1.4.1.1.6.2	Method 2: Alkyl- or Arylthiolation	153
15.1.1.4.1.1.6.3	Method 3: Arylselenation	153
15.1.1.4.1.1.7	By Nitrogen	153
15.1.1.4.1.1.7.1	Method 1: Nitration	153
15.1.1.4.1.1.7.2	Method 2: Direct Amination or Alkylamination with Alkali Amides (Tschitschibabin Reaction)	154
15.1.1.4.1.1.7.2.1	Variation 1: Amination of Nitropyridines (Vicarious Nucleophilic Substitution)	154
15.1.1.4.1.1.7.2.2	Variation 2: Acylamination in the Presence of Fluorine	155
15.1.1.4.1.1.7.3	Method 3: Alkylamination via Hetarynes	156
15.1.1.4.1.1.8	By Phosphorus	156
15.1.1.4.1.1.8.1	Method 1: Phosphorylation	156

15.1.1.4.1.2	Of Metals	157
15.1.1.4.1.2.1	By a Carbon Functionality	157
15.1.1.4.1.2.1.1	Method 1: Replacement of Silicon	157
15.1.1.4.1.2.1.2	Method 2: Replacement of Tin	158
15.1.1.4.1.2.1.3	Method 3: Replacement of Boron	159
15.1.1.4.1.3	Of Carbon Functionalities	159
15.1.1.4.1.3.1	By Hydrogen	159
15.1.1.4.1.3.1.1	Method 1: Decarboxylation	159
15.1.1.4.1.3.2	By Another Carbon Functionality	160
15.1.1.4.1.3.2.1	Method 1: The Hammick Reaction	160
15.1.1.4.1.3.3	By a Halogen	160
15.1.1.4.1.3.3.1	Method 1: Borodin–Hundsdiecker-Type Halogenation	160
15.1.1.4.1.3.4	By Nitrogen	161
15.1.1.4.1.3.4.1	Method 1: Curtius Rearrangement	161
15.1.1.4.1.4	Of Halogens	162
15.1.1.4.1.4.1	By Deuterium or Hydrogen	162
15.1.1.4.1.4.1.1	Method 1: Reduction	162
15.1.1.4.1.4.1.2	Method 2: Deuteration via Metalation	162
15.1.1.4.1.4.2	By Metals	162
15.1.1.4.1.4.2.1	Method 1: By Lithium or Magnesium	162
15.1.1.4.1.4.2.2	Method 2: By Mercury, Arsenic, Silicon, Tin, or Boron	163
15.1.1.4.1.4.3	By a Carbon Functionality	164
15.1.1.4.1.4.3.1	Method 1: Cyanation	164
15.1.1.4.1.4.3.2	Method 2: Palladium-Catalyzed Aminocarbonylation, Formylation, or Alkoxycarbonylation	165
15.1.1.4.1.4.3.3	Method 3: Palladium-Catalyzed Coupling with Alkynes and Alkenes	166
15.1.1.4.1.4.3.4	Method 4: Metal-Catalyzed Aryl–Hetaryl or Hetaryl–Hetaryl Coupling ..	167
15.1.1.4.1.4.3.4.1	Variation 1: Palladium-Catalyzed Coupling with a 2-Pyridylboronate Generated In Situ	168
15.1.1.4.1.4.3.5	Method 5: Alkylation	169
15.1.1.4.1.4.3.5.1	Variation 1: Via Photochemical Substitution	170
15.1.1.4.1.4.4	By a Different Halogen	170
15.1.1.4.1.4.4.1	Method 1: Halogen Exchange under Lewis Acid Catalysis	170
15.1.1.4.1.4.5	By Oxygen	171
15.1.1.4.1.4.5.1	Method 1: Reaction with Oxygen Nucleophiles	171
15.1.1.4.1.4.6	By Sulfur	172
15.1.1.4.1.4.6.1	Method 1: Reaction with Sulfur Nucleophiles	172
15.1.1.4.1.4.6.2	Method 2: Introduction of a Sulfinyl Group via Metalation	173
15.1.1.4.1.4.7	By Selenium or Tellurium	173
15.1.1.4.1.4.7.1	Method 1: Reaction with Selenium or Tellurium Nucleophiles	173

15.1.1.4.1.4.8	By Nitrogen	174
15.1.1.4.1.4.8.1	Method 1: Reaction with Nitrogen Nucleophiles	174
15.1.1.4.1.4.8.1.1	Variation 1: Palladium-Catalyzed Bromine–Nitrogen Exchange	174
15.1.1.4.1.4.8.1.2	Variation 2: Alkylamination under High Pressure	175
15.1.1.4.1.4.9	By Phosphorus	175
15.1.1.4.1.4.9.1	Method 1: Direct Phosphorylation	175
15.1.1.4.1.4.9.2	Method 2: Phosphorylation via Metalation	176
15.1.1.4.1.4.9.2.1	Variation 1: Photochemical Phosphorylation	176
15.1.1.4.1.5	Of Oxygen or Sulfur	177
15.1.1.4.1.5.1	By Hydrogen	177
15.1.1.4.1.5.1.1	Method 1: Desulfurization	177
15.1.1.4.1.5.1.2	Method 2: Deoxygenation	177
15.1.1.4.1.5.2	By a Carbon Functionality	178
15.1.1.4.1.5.2.1	Method 1: Displacement with Carbon Nucleophiles	178
15.1.1.4.1.5.3	By a Halogen	180
15.1.1.4.1.5.3.1	Method 1: Halogenation with Phosphoryl Halides	180
15.1.1.4.1.5.4	By Nitrogen	180
15.1.1.4.1.5.4.1	Method 1: Oxygen–Nitrogen Exchange at Elevated Temperatures	180
15.1.1.4.1.6	Of Nitrogen	181
15.1.1.4.1.6.1	By a Halogen	181
15.1.1.4.1.6.1.1	Method 1: Sandmeyer and Schiemann Reactions	181
15.1.1.4.1.6.2	By Oxygen or Sulfur	182
15.1.1.4.1.6.2.1	Method 1: 1,3-Dipolar Addition	182
15.1.1.4.1.6.2.2	Method 2: Exchange via Diazotization	182
15.1.1.4.2	Addition Reactions	183
15.1.1.4.3	Rearrangement of Substituents	183
15.1.1.4.3.1	Method 1: Isomerization	183
15.1.1.4.3.2	Method 2: Rearrangement of Halogens	183
15.1.1.4.4	Modification of Substituents	184
15.1.1.4.4.1	From Annulated Pyridines by Selective Degradation	184
15.1.1.4.4.1.1	Method 1: From Quinolines or Isoquinolines	184
15.1.1.4.4.1.1.1	Variation 1: Oxidation of Quinolines	184
15.1.1.4.4.1.1.2	Variation 2: Oxidation of Isoquinolines	185
15.1.1.4.4.1.1.3	Variation 3: Partial Hydrogenation of Quinolines	185
15.1.1.4.4.1.1.4	Variation 4: Partial Hydrogenation or Reduction of Isoquinolines	186
15.1.1.4.4.1.1.5	Variation 5: Ionic Hydrogenation of Quinolines or Isoquinolines	186
15.1.1.4.4.1.2	Method 2: From [1,2,3]Triazolo[1,5- <i>a</i>]pyridines	187
15.1.1.4.4.1.2.1	Variation 1: From 1 <i>H</i> -1,2,3-Triazolo[4,5- <i>c</i>]pyridin-1-amine	188
15.1.1.4.4.1.3	Method 3: From Quinolizinium Salts	189
15.1.1.4.4.1.4	Method 4: From Furo[2,3- <i>c</i>]pyridines	189
15.1.1.4.4.2	From Pyridine 1-Oxides by Deoxygenation	190

15.1.1.4.4.2.1	Method 1:	Reduction	190
15.1.1.4.4.2.2	Method 2:	Reaction with Acid Chlorides or Phosphoryl Chloride	191
15.1.1.4.4.2.3	Method 3:	Reaction with Acid Anhydrides [Deoxygenation and Side-Chain Substitution (The Boekelheide Reaction)]	192
15.1.1.4.4.2.3.1	Variation 1:	Deoxygenation and Side-Chain Halogenation	192
15.1.1.4.4.2.4	Method 4:	Addition–Elimination Reactions	193
15.1.1.4.4.2.4.1	Variation 1:	Decarboxylation–Substitution Reactions	194
15.1.1.4.4.3		From Pyridinium Compounds	194
15.1.1.4.4.3.1	Method 1:	Addition–Elimination Reactions	194
15.1.1.4.4.4		By Addition to Existing Substituents	195
15.1.1.4.4.4.1	Method 1:	Aromatization of Pyridinones and Pyridinethiones by Alkylation, Acylation, or Silylation	195
15.1.1.4.4.4.2	Method 2:	Oxidative Aromatization of Pyridinethiones	197
15.1.1.4.4.5		By Modification of Side Chains	197
15.1.1.4.4.5.1	Method 1:	Oxidation of (Hydroxymethyl)pyridines	197
15.1.1.4.4.5.2	Method 2:	Deprotonation of Alkylpyridines	197
15.1.1.4.4.5.2.1	Variation 1:	Deuteration of Alkyl Groups	198
15.1.1.4.4.5.2.2	Variation 2:	Deprotonation with Strong Bases, Followed by Alkylation or Silylation	198
15.1.1.4.4.5.3	Method 3:	Side-Chain Deoxygenation	199
15.1.1.4.4.5.4	Method 4:	Formation of <i>o</i> -Quinodimethanes by Dehalogenation	199
15.1.2		Product Subclass 2: Pyridine 1-Oxides	200
15.1.2.1		Synthesis by Ring-Closure Reactions	201
15.1.2.1.1		By Formation of Two N–C Bonds and One C–C Bond	201
15.1.2.1.1.1		Fragments C–C–C, C–C, and N	201
15.1.2.1.1.1.1	Method 1:	From Vinylogous Amidinium Salts, Ketones, and Hydroxylamine	201
15.1.2.1.2		By Formation of Two N–C Bonds	202
15.1.2.1.2.1		Fragments C–C–C–C–C and N	202
15.1.2.1.2.1.1	Method 1:	From 1,5-Dioxopent-2-enes and Hydroxylamine	202
15.1.2.1.2.1.1.1	Variation 1:	From Acyl Dienamines or Masked 1,5-Dioxo Compounds and Hydroxylamine	202
15.1.2.1.2.1.1.2	Variation 2:	From 1,1-Dicyanobutadienes and Hydroxylamine	203
15.1.2.1.2.1.1.3	Variation 3:	From 5,5-Dichloropentadienals and Hydroxylamine	204
15.1.2.1.3		By Formation of One N–C and One C–C Bond	204
15.1.2.1.3.1		Fragments C–C–C–C and N–C	204
15.1.2.1.3.1.1	Method 1:	From Ethylidenemalononitriles and <i>N</i> -Hydroxybenzimidoyl Chlorides	204
15.1.2.1.4		By Formation of One N–C Bond	205
15.1.2.1.4.1		Fragment N–C–C–C–C	205
15.1.2.1.4.1.1	Method 1:	Cyclization of Pentadienal Oximes	205
15.1.2.1.4.1.2	Method 2:	Cyclization of Enynal Oximes	205

15.1.2.2	Synthesis by Ring Transformation	206
15.1.2.2.1	By Ring Enlargement	206
15.1.2.2.1.1	Method 1: From Isoxazoles	206
15.1.2.2.1.1.1	Variation 1: From 4,5-Dihydroisoxazoles	206
15.1.2.2.2	By Formal Exchange of Ring Members with Retention of the Ring Size	207
15.1.2.2.2.1	Method 1: From Pyrylium Salts and Hydroxylamine	207
15.1.2.2.2.2	Method 2: From Pyridinium Salts and Hydroxylamine	207
15.1.2.2.2.3	Method 3: From 3-Acyl-6-alkoxy-5,6-dihydro-4 <i>H</i> -1,2-oxazines	208
15.1.2.3	Synthesis by Substituent Modification	208
15.1.2.3.1	Substitution of Existing Substituents	208
15.1.2.3.1.1	Of Hydrogen	208
15.1.2.3.1.1.1	By Metals	208
15.1.2.3.1.1.1.1	Method 1: Intermediate Lithiation	208
15.1.2.3.1.1.2	By a Carbon Functionality	209
15.1.2.3.1.1.2.1	Method 1: Mannich Reaction	209
15.1.2.3.1.1.2.2	Method 2: Alkylation and Acylation via Metalation	209
15.1.2.3.1.1.3	By a Halogen	210
15.1.2.3.1.1.3.1	Method 1: Bromination	210
15.1.2.3.1.1.3.1.1	Variation 1: Bromination via Metalation	211
15.1.2.3.1.1.4	By Nitrogen	211
15.1.2.3.1.1.4.1	Method 1: Nitration	211
15.1.2.3.1.2	Of Carbon Functionalities	212
15.1.2.3.1.2.1	By Hydrogen	212
15.1.2.3.1.2.1.1	Method 1: Decarboxylation and Deacylation	212
15.1.2.3.1.2.2	By Nitrogen	212
15.1.2.3.1.2.2.1	Method 1: Curtius Rearrangement	212
15.1.2.3.1.3	Of a Halogen	213
15.1.2.3.1.3.1	By Hydrogen	213
15.1.2.3.1.3.1.1	Method 1: Protonation via Metalation	213
15.1.2.3.1.3.2	By a Carbon Functionality	213
15.1.2.3.1.3.2.1	Method 1: Metal-Catalyzed Aryl–Hetaryl or Hetaryl–Hetaryl Coupling ..	213
15.1.2.3.1.3.2.1.1	Variation 1: Stille Coupling	214
15.1.2.3.1.3.2.1.2	Variation 2: Suzuki–Miyaura-Type Coupling	214
15.1.2.3.1.3.2.1.3	Variation 3: Zinc-Promoted Coupling	215
15.1.2.3.1.3.3	By Oxygen	215
15.1.2.3.1.3.3.1	Method 1: Reaction with Oxygen Nucleophiles	215
15.1.2.3.1.3.4	By Nitrogen	216
15.1.2.3.1.3.4.1	Method 1: Reaction with Amino Compounds	216
15.1.2.3.1.4	Of Nitrogen	216

15.1.2.3.1.4.1	By a Halogen	216
15.1.2.3.1.4.1.1	Method 1: Replacement of a Nitro Group by Reaction with Acetyl Halides	216
15.1.2.3.1.4.2	By Oxygen	217
15.1.2.3.1.4.2.1	Method 1: Replacement of a Nitro Group by Reaction with Oxygen Nucleophiles	217
15.1.2.3.1.4.3	By Another Nitrogen	218
15.1.2.3.1.4.3.1	Method 1: Replacement of a Nitro Group by Reaction with Ammonia ..	218
15.1.2.3.2	Addition Reactions	218
15.1.2.3.2.1	Method 1: N-Oxidation of Pyridines	218
15.1.2.3.3	Modification of Substituents	219
15.1.2.3.3.1	Method 1: Alkylation and Acylation of Alkyl Substituents	219
15.1.2.3.3.2	Method 2: Oxidation of Side Chains	220
15.1.2.3.3.3	Method 3: Reduction of Acyl Substituents	220
15.1.3	Product Subclass 3: Pyridinium Salts	221
15.1.3.1	Synthesis by Ring-Closure Reactions	222
15.1.3.1.1	By Formation of Two N—C Bonds and One C—C Bond	222
15.1.3.1.1.1	Fragments C—C—C, C—C, and N	222
15.1.3.1.1.1.1	Method 1: From α,β -Unsaturated Ketones, Amines, and a Michael Donor	222
15.1.3.1.1.1.2	Method 2: From α -Bromo Ketones and Primary Amines	223
15.1.3.1.1.2	Fragments C—C—C—C, C, and N	224
15.1.3.1.1.2.1	Method 1: From an α,β -Unsaturated Aldehyde, a Primary Amine, and an Aldehyde	224
15.1.3.1.2	By Formation of Two N—C Bonds	224
15.1.3.1.2.1	Fragments C—C—C—C—C and N	224
15.1.3.1.2.1.1	Method 1: From Protected or Free 1,5-Dioxo Compounds and Primary Amines	224
15.1.3.1.2.1.2	Method 2: 1-Aminopyridinium Compounds from Oxonium Salts and Hydrazides	225
15.1.3.1.2.1.3	Method 3: From 2 <i>H</i> -1,2-Thiazine-6-carbaldehyde 1,1-Dioxides and Primary Amines	226
15.1.3.1.2.1.4	Method 4: From Acyl Enynes and Primary Amines	226
15.1.3.1.3	By Formation of One N—C and One C—C Bond	227
15.1.3.1.3.1	Fragments N—C—C—C and C—C	227
15.1.3.1.3.1.1	Method 1: From [β -(Monoalkylamino)vinyl]carbene—Chromium Complexes and Alkynes	227
15.1.3.1.3.1.2	Method 2: Cyclization of Enamino Iminium Salts with 1,3-Dicarbonyl Compounds	227
15.1.3.1.3.2	Fragments C—C—C—C and N—C	228
15.1.3.1.3.2.1	Method 1: From β,γ -Unsaturated Ketones and Imines	228
15.1.3.1.3.2.2	Method 2: From 1,3-Diketones and Formanilides	228

15.1.3.1.3.3	Fragments N—C—C and C—C—C	229
15.1.3.1.3.3.1	Method 1: From α -(Aminomethylene) Carbonyl Compounds and Methyl Imines	229
15.1.3.1.3.3.1.1	Variation 1: From 1,3-Dicarbonyl Compounds and Methyl Imines	230
15.1.3.1.3.3.2	Method 2: From <i>N</i> -Alkyl-2-cyanoacetamides in the Presence of Phosphoryl Chloride	230
15.1.3.1.4	By Formation of One N—C Bond	231
15.1.3.1.4.1	Fragment N—C—C—C—C	231
15.1.3.1.4.1.1	Method 1: Cyclization of 1-Aza-1,3,5-trienes	231
15.1.3.1.5	By Formation of One C—C Bond	231
15.1.3.1.5.1	Fragment C—C—N—C—C—C	231
15.1.3.1.5.1.1	Method 1: Cyclization of 3-Alkyl-1,5-dioxo-3-azahexanes	231
15.1.3.1.5.1.2	Method 2: Cyclization of Diethyl 2-[(Alkylamino)methylene]succinates ..	232
15.1.3.2	Synthesis by Ring Transformation	233
15.1.3.2.1	By Ring Enlargement	233
15.1.3.2.1.1	Method 1: From Furans	233
15.1.3.2.1.1.1	Variation 1: From 5-(Hydroxymethyl)furan-2-carbaldehyde and Amino Compounds	234
15.1.3.2.2	By Formal Exchange of Ring Members with Retention of the Ring Size	234
15.1.3.2.2.1	From Pirylium Compounds	234
15.1.3.2.2.1.1	Method 1: From Pirylium Compounds and Primary Amines	234
15.1.3.2.2.1.2	Method 2: Formation of 1-Aminopyridinium Compounds from Pirylium Salts and Hydrazines	235
15.1.3.2.2.1.3	Method 3: From Pirylium Compounds and Methyl Imines	236
15.1.3.2.2.2	From 4 <i>H</i> -Pyran-4-ones	237
15.1.3.2.2.2.1	Method 1: From 4 <i>H</i> -Pyran-4-ones and Primary Amines	237
15.1.3.2.2.3	From Other Pyridinium Compounds	237
15.1.3.2.2.3.1	Method 1: From Pyridinium Compounds and Primary Amines or Hydrazines by an S_N (ANRORC) Mechanism	237
15.1.3.3	Aromatization	238
15.1.3.3.1	Method 1: Oxidation of 1-Alkyl- or 1-Aryl-1,4-dihydropyridines	238
15.1.3.3.2	Method 2: Oxidation of 1-Substituted Tetrahydropyridines	239
15.1.3.3.3	Method 3: Elimination from 4-Substituted Dihydropyridines	240
15.1.3.4	Synthesis by Substituent Modification	241
15.1.3.4.1	Substitution of Existing Substituents	241
15.1.3.4.1.1	Of Hydrogen	241
15.1.3.4.1.1.1	By Metals	241
15.1.3.4.1.1.1.1	Method 1: Lithiation	241
15.1.3.4.1.1.2	By a Carbon Functionality	241
15.1.3.4.1.1.2.1	Method 1: Alkylation and Arylation	241

15.1.3.4.1.1.3	By Phosphorus	242
15.1.3.4.1.1.3.1	Method 1: Phosphorylation	242
15.1.3.4.1.2	Of Halogens	243
15.1.3.4.1.2.1	By a Carbon Functionality	243
15.1.3.4.1.2.1.1	Method 1: Reaction with Enamines	243
15.1.3.4.1.2.1.2	Method 2: Stille Coupling	244
15.1.3.4.1.2.2	By Sulfur	245
15.1.3.4.1.2.2.1	Method 1: Reaction with Sulfur Nucleophiles	245
15.1.3.4.1.2.3	By Nitrogen	245
15.1.3.4.1.2.3.1	Method 1: Reaction with Nitrogen Nucleophiles	245
15.1.3.4.1.3	Of Oxygen	246
15.1.3.4.1.3.1	By a Halogen	246
15.1.3.4.1.3.1.1	Method 1: Chlorination of N-Substituted Pyridinones	246
15.1.3.4.2	Addition Reactions	246
15.1.3.4.2.1	Method 1: Protonation of 4-Ilidene-1,4-dihydropyridines	246
15.1.3.4.2.2	Method 2: N-Silylation of Pyridines	247
15.1.3.4.2.3	Method 3: N-Alkylation of Pyridines	248
15.1.3.4.2.3.1	Variation 1: N-Halomethylation with Formaldehyde and Thionyl Halides ..	249
15.1.3.4.2.3.2	Variation 2: N-Arylation with Electrophilic Aryl or Hetaryl Groups	249
15.1.3.4.2.3.3	Variation 3: N-Acylation with Acyl Halides	250
15.1.3.4.2.4	Method 4: N-Fluorination of Pyridines	250
15.1.3.4.2.5	Method 5: N-Sulfonylation of Pyridines	250
15.1.3.4.2.6	Method 6: N-Amination and N-Nitration of Pyridines	251
15.1.3.4.2.7	Method 7: N-Phosphorylation of Pyridines	252
15.1.3.4.3	Modification of Substituents	252
15.1.3.4.3.1	Method 1: Alkylation of Alkyl Substituents	252
15.1.3.4.3.2	Method 2: Reduction of Acyl Substituents	253
15.1.3.4.3.3	Method 3: O-Alkylation or O-Acylation of Pyridine 1-Oxides	253
15.1.3.4.3.4	Method 4: O-Alkylation of N-Substituted Pyridinones	254
15.1.3.4.3.5	Method 5: S-Alkylation or S-Acylation of N-Substituted Pyridinethiones ..	255
15.2	Product Class 2: Pyridinones and Related Systems	
	P. A. Keller	
15.2	Product Class 2: Pyridinones and Related Systems	285
15.2.1	Product Subclass 1: Pyridin-2(1H)-ones	285
15.2.1.1	Synthesis by Ring-Closure Reactions	285
15.2.1.1.1	By Formation of Two N—C and Two C—C Bonds	286
15.2.1.1.1.1	Fragments C—C, C—C, C, and N	286
15.2.1.1.1.1.1	Method 1: From an Aldehyde, a Ketone, an Active Methylene Ester, and an Ammonium Salt	286

15.2.1.1.2	By Formation of One N—C and Two C—C Bonds	287
15.2.1.1.2.1	Fragments N—C, C—C, and C—C	287
15.2.1.1.2.1.1	Method 1: Cyclization Using Isocyanates and Disubstituted Alkynes	287
15.2.1.1.2.1.1.1	Variation 1: Cyclization Using Isocyanates and Diketene	289
15.2.1.1.3	By Formation of One N—C and One C—C Bond	290
15.2.1.1.3.1	Fragments N—C—C—C and C—C	290
15.2.1.1.3.1.1	Method 1: Self Condensation of β -Oxoamides	290
15.2.1.1.3.1.1.1	Variation 1: Thermal Autocondensation of Oxoamides	290
15.2.1.1.3.2	Fragments C—C—C—C and N—C	291
15.2.1.1.3.2.1	Method 1: Addition of Activated Methylene Compounds to Imidates	291
15.2.1.1.3.2.2	Method 2: Addition of Imines to α,β -Unsaturated Carbonyls and Subsequent Cyclization	292
15.2.1.1.3.2.3	Method 3: Photoreaction of Arenecarbothioamides with α,β -Unsaturated Carboxylic Acids	293
15.2.1.1.3.2.4	Method 4: [4+2] Cycloaddition of Cobaltacyclopentadienes and Isocyanates	294
15.2.1.1.3.3	Fragments N—C—C and C—C—C	294
15.2.1.1.3.3.1	Method 1: Addition of Activated Methylene Compounds to Ketones	294
15.2.1.1.3.3.2	Method 2: Addition of Activated Methylene Compounds to Aldehydes ..	295
15.2.1.1.3.3.3	Method 3: Addition of Activated Methylene Compounds to α,β -Unsaturated Ketones with Elimination of Ethoxide	297
15.2.1.1.3.3.4	Method 4: Addition of Activated Methylene Compounds to α,β -Unsaturated Carbonyl Compounds	298
15.2.1.1.3.3.5	Method 5: Addition of Activated Methylene Compounds to Resin-Bound α,β -Unsaturated Ketones	299
15.2.1.1.3.3.6	Method 6: Addition of Activated Methylene Compounds to Ketene Dithioacetal Compounds	300
15.2.1.1.3.3.7	Method 7: Addition of Activated Methylene Compounds to Ethyl Propynoate	301
15.2.1.1.3.3.8	Method 8: Addition of an Activated Methylene Compound to an Acylketene	301
15.2.1.1.3.3.9	Method 9: Resin-Supported Synthesis of Enaminones	301
15.2.1.1.3.3.10	Method 10: Addition of Activated Methylene Compounds to α,β -Unsaturated Ketones with Sequential Aromatization	303
15.2.1.1.3.3.11	Method 11: Sequential Nucleophilic Substitution/Addition to Malonyl Chloride	304
15.2.1.1.4	By Formation of Two C—C Bonds	304
15.2.1.1.4.1	Fragments C—N—C—C and C—C	304
15.2.1.1.4.1.1	Method 1: [4+2] Cycloadditions Using Acyl Isocyanates	304
15.2.1.1.4.1.2	Method 2: 1,3-Dipolar Cycloadditions	305
15.2.1.1.4.2	Fragments C—C—N—C—C and C	306
15.2.1.1.4.2.1	Method 1: The Use of Vilsmeier Reagents	306
15.2.1.1.4.2.1.1	Variation 1: Dual-Purpose Substrate and Vilsmeier Reagent	307

15.2.1.1.5	By Formation of One N—C Bond	308
15.2.1.1.5.1	Fragment N—C—C—C—C	308
15.2.1.1.5.1.1	Method 1: Acid-Catalyzed Cyclization of Oxoamides	308
15.2.1.1.5.1.1.1	Variation 1: Hydrolysis of 5-Amino-3-oxopent-4-enitriles	309
15.2.1.1.5.1.2	Method 2: Resin-Supported Ring-Closing Reactions of Diesters with Amines	309
15.2.1.1.5.1.3	Method 3: Zinc Reduction of Nitroso-Diels–Alder Adducts	311
15.2.1.1.5.1.4	Method 4: Cyclization of Ketene Dithioacetals	312
15.2.1.1.6	By Formation of One C—C Bond	312
15.2.1.1.6.1	Fragment C—N—C—C—C	312
15.2.1.1.6.1.1	Method 1: Intramolecular Cyclization of an In Situ Generated Diene Isocyanate	312
15.2.1.1.6.1.2	Method 2: Intramolecular Cyclization of Alka-1,3-dienyl Isocyanates	314
15.2.1.1.6.1.3	Method 3: Insertion of Amines into Diketones	315
15.2.1.1.6.2	Fragment C—C—N—C—C	316
15.2.1.1.6.2.1	Method 1: Thermal Cyclization of Ketene Dithioacetals	316
15.2.1.2	Aromatization	317
15.2.1.2.1	Method 1: By Utilization of an Exocyclic Alkene	317
15.2.1.2.2	Method 2: By Elimination of Two Hydrogen Atoms Using Nitrous Acid ..	318
15.2.1.2.2.1	Variation 1: By Oxidation of 3,4-Dihydropyridin-2(1 <i>H</i>)-ones Using Nitrosylsulfuric Acid	318
15.2.1.2.2.2	Variation 2: By Oxidation of 6-Hydroxytetrahydropyridin-2(1 <i>H</i>)-ones Using Thionyl Chloride and Pyridine	319
15.2.1.2.2.3	Variation 3: By Oxidation Using 2,3-Dichloro-5,6-dicyanobenzo-1,4-quinone	320
15.2.1.2.2.4	Variation 4: By Dehydrogenation under Alcoholic Basic Conditions	321
15.2.1.2.2.5	Variation 5: By Controlled Use of Nitrous Acid	321
15.2.1.3	Synthesis by Substituent Modification	321
15.2.1.3.1	Method 1: By Intramolecular Oxygen-Transfer Reactions	321
15.2.1.3.1.1	Variation 1: By Alkyl Elimination from 2-Ethoxypyridinium Oxide	322
15.2.1.3.2	Method 2: By Oxidation of Pyridines	323
15.2.1.3.2.1	Variation 1: By Oxidation of 1-Aminopyridinium Salts	324
15.2.1.3.2.2	Variation 2: By Oxidation of 1-Alkylpyridinium Salts	325
15.2.1.3.3	Method 3: By Oxidation of 2-Methylpyridinium Salts	326
15.2.1.3.4	Method 4: By Displacement from Pyridinium Salts	327
15.2.1.3.4.1	Variation 1: By Methoxy Substitution of 2-Chloropyridines	328
15.2.1.3.4.2	Variation 2: By Nucleophilic Substitution of 2-(Methylsulfanyl)pyridines ..	329
15.2.1.3.4.3	Variation 3: By Substitution and Subsequent Rearrangement	329
15.2.1.3.4.4	Variation 4: By Resin-Supported Conversion of 2-Halopyridines	330
15.2.1.3.4.5	Variation 5: By Oxidative Demethylation of Methylpyridinium Cations	331
15.2.1.3.5	Method 5: By Rearrangement of 2-Allyloxypyridines	332
15.2.1.3.6	Method 6: By Oxidation of Pyridine-2(1 <i>H</i>)-thiones	334
15.2.2	Product Subclass 2: Pyridin-4(1<i>H</i>)-ones	335
15.2.2.1	Synthesis by Ring-Closure Reactions	335

15.2.2.1.1	By Formation of Two N—C Bonds and One C—C Bond	335
15.2.2.1.1.1	Fragments C—C—C, C, and N	335
15.2.2.1.1.1.1	Method 1: In Situ Acylation of Enaminone Anions with Cyclization	335
15.2.2.1.2	By Formation of One N—C and Two C—C Bonds	337
15.2.2.1.2.1	Fragments C—C—C, N—C, and C	337
15.2.2.1.2.1.1	Method 1: Addition of an <i>N</i> -Silyl-1-azaallyl Anion to Alkoxyalkenes	337
15.2.2.1.3	By Formation of Two N—C Bonds	337
15.2.2.1.3.1	Fragments C—C—C—C and N	337
15.2.2.1.3.1.1	Method 1: Nitrogen Exchange in Lactones	337
15.2.2.1.3.1.2	Method 2: Ring Closure of Triketones	338
15.2.2.1.3.1.3	Method 3: Cyclization of 5-Methoxypent-4-en-1-ones Followed by Amination with Hydroxylamine Hydrochloride	339
15.2.2.1.4	By Formation of One N—C and One C—C Bond	341
15.2.2.1.4.1	Fragments C—C—C and N—C	341
15.2.2.1.4.1.1	Method 1: Addition of β -Cyano Ketones to Cyclic Imines	341
15.2.2.1.4.1.2	Method 2: Solid-Phase Synthesis of Dihydropyridinones	341
15.2.2.1.4.2	Fragments N—C—C and C—C—C	343
15.2.2.1.4.2.1	Method 1: Thermal Cyclization of Enaminones with Ketenes	343
15.2.2.1.4.2.2	Method 2: Claisen Addition of Pyridylacetonitriles to Alk-2-ynoates	344
15.2.2.1.5	By Formation of Two C—C Bonds	346
15.2.2.1.5.1	Fragments C—C—N—C—C and C	346
15.2.2.1.5.1.1	Method 1: Carbonylation of 2-Aza-1,3-dienes	346
15.2.2.1.6	By Formation of One N—C Bond	346
15.2.2.1.6.1	Fragment N—C—C—C—C	346
15.2.2.1.6.1.1	Method 1: Cyclization with Nitrogen Nucleophiles onto Cyano Electrophiles	346
15.2.2.1.6.1.2	Method 2: Metal Carbonyl Induced Cleavage of Isoxazoles	347
15.2.2.1.7	By Formation of One C—C Bond	348
15.2.2.1.7.1	Fragment C—C—N—C—C—C	348
15.2.2.1.7.1.1	Method 1: Addition of Amines to α,β -Unsaturated Esters	348
15.2.2.2	Synthesis by Ring Transformation	349
15.2.2.2.1	Method 1: By Heteroatom Exchange in Pyran-4-ones Using Amines	349
15.2.2.2.1.1	Variation 1: By Heteroatom Exchange Using Pyranones and Ammonia	351
15.2.2.2.1.2	Variation 2: By Heteroatom Exchange Using Hydroxylactones	352
15.2.2.2.2	Method 2: By Cyclic Nitrone Cycloaddition to Acetylenes	353
15.2.2.3	Aromatization	353
15.2.2.3.1	Method 1: By Michael Addition to Cyclic Diones	353
15.2.2.3.2	Method 2: By Formation of Diselenides Followed by Oxidation	354
15.2.2.3.3	Method 3: By the Generation of Imines from Dihydropyridin-4(1 <i>H</i>)-ones	355
15.2.2.4	Synthesis by Substituent Modification	355

15.2.2.4.1	Method 1:	By Substitution and Subsequent Rearrangement	355
15.2.2.4.2	Method 2:	By Rearrangement of 2-Substituted 4-Methoxypyridines	356
15.2.2.4.3	Method 3:	From Pyridin-2(1 <i>H</i>)-ones by Functional Group Transformations	357
15.2.2.4.4	Method 4:	By Oxidative Demethylation of Methylpyridinium Cations	358
15.2.2.4.5	Method 5:	Via Metalation of Dichloropyridines	359
15.2.2.4.6	Method 6:	By Oxidative Addition to Halopyridines	359
15.2.2.4.7	Method 7:	By Substitution of 4-Nitropyridine 1-Oxides	360
15.2.2.4.7.1	Variation 1:	By Methoxy Substitution of Nitropyridines	361
15.2.3	Product Subclass 3: Pyridine-2(1<i>H</i>)-thiones		362
15.2.3.1	Synthesis by Ring-Closure Reactions		362
15.2.3.1.1	By Formation of One N—C and One C—C Bond		363
15.2.3.1.1.1	Fragments C—C—C—C—C and N—C		363
15.2.3.1.1.1.1	Method 1:	By Reaction of the Glutacondialdehyde Anion with Isothiocyanates	363
15.2.3.1.1.2	Fragments N—C—C and C—C—C		363
15.2.3.1.1.2.1	Method 1:	Addition of Activated Methylene Compounds to α,β -Unsaturated Ketones in the Absence of Base	363
15.2.3.1.1.2.1.1	Variation 1:	Addition of Activated Methylene Compounds to α,β -Unsaturated Ketones in the Presence of Base	364
15.2.3.1.1.2.2	Method 2:	Addition of Activated Methylene Compounds to α,β -Unsaturated Aldehydes	365
15.2.3.1.1.2.3	Method 3:	Addition of Thioamide—Cyano Activated Methylene Groups to β -Dicarbonyls	365
15.2.3.1.1.2.4	Method 4:	Addition of Thioamide—Amide Activated Methylene Groups to β -Diketones	366
15.2.3.1.1.2.5	Method 5:	Addition of Thioamide—Cyano Activated Methylene Groups to α,β -Unsaturated Dicyanides	366
15.2.3.1.1.2.6	Method 6:	Addition of Thioamide—Cyano Activated Methylene Groups to α,β -Unsaturated α -Cyano Esters	367
15.2.3.2	Synthesis by Substituent Modification		368
15.2.3.2.1	Method 1:	Substitution of 2-Halopyridines with Hydrogen Sulfide or Thiourea	368
15.2.3.2.2	Method 2:	Reactions of Dihydropyridin-2(1 <i>H</i>)-ones with Phosphorus Pentasulfide	369
15.2.3.2.3	Method 3:	Rearrangement of Thio Salts of 1-Oxidopyridine-2(1 <i>H</i>)-thione	369
15.2.3.2.4	Method 4:	By Reaction of 1-Alkylpyridinium Salts with Sodium Sulfide	370
15.2.4	Product Subclass 4: Pyridine-4(1<i>H</i>)-thiones		371
15.2.4.1	Synthesis by Ring Transformation		371
15.2.4.1.1	Method 1:	By Heteroatom Exchange in Pyran-4-thiones Using Amines	371
15.2.4.1.1.1	Variation 1:	By Methylamine Exchange with 4 <i>H</i> -Pyran-4-thiones	372
15.2.4.2	Synthesis by Substituent Modification		373
15.2.4.2.1	Method 1:	Reaction of Pyridin-4(1 <i>H</i>)-ones with Phosphorus Pentasulfide	373
15.2.5	Product Subclass 5: Pyridine-2(1<i>H</i>)-selones and Pyridine-2(1<i>H</i>)-tellones		374

15.2.5.1	Synthesis by Ring-Closure Reactions	374
15.2.5.1.1	By Formation of One N—C and One C—C Bond	374
15.2.5.1.1.1	Fragments C—C—C—C and N—C	374
15.2.5.1.1.1	Method 1: Reaction of the Glutacondialdehyde Anion with Isoselenocyanates	374
15.2.5.1.1.2	Fragments N—C—C and C—C—C	375
15.2.5.1.1.2.1	Method 1: Addition of Cyanoselenoacetamide to α,β -Unsaturated Ketones	375
15.2.5.1.1.2.2	Method 2: Addition of Cyanoselenoacetamide to β -Diketones	378
15.2.5.1.1.2.3	Method 3: Addition of Selenoamide—Cyano Activated Methylene Groups to α,β -Unsaturated Dicyanides	378
15.2.5.1.2	By Formation of One N—C Bond	379
15.2.5.1.2.1	Fragment N—C—C—C—C	379
15.2.5.1.2.1.1	Method 1: Addition of Hydrogen Selenide to 1-Aryl-4,4-dicyanobutan- 1-ones	379
15.2.5.2	Synthesis by Substituent Modification	380
15.2.5.2.1	Method 1: By Substitution of 2-Halopyridines with Selenium Reagents	380
15.2.6	Product Subclass 6: Pyridine-4(1H)-selones and Pyridine-4(1H)-tellones	382
15.2.6.1	Synthesis by Substituent Modification	383
15.2.6.1.1	Method 1: By Heteroatom Exchange from Pyridine-4(1H)-thiones	383
15.3	Product Class 3: Quinolines R. D. Larsen and D. Cai	
15.3	Product Class 3: Quinolines	389
15.3.1	Product Subclass 1: Quinolines	389
15.3.1.1	Synthesis by Ring-Closure Reactions	391
15.3.1.1.1	By Annulation to an Arene	391
15.3.1.1.1.1	By Formation of One N—C and Two C—C Bonds	391
15.3.1.1.1.1.1	With Formation of the 1—2, 2—3, and 4—4a Bonds	391
15.3.1.1.1.1.1.1	Method 1: From Arylamines	391
15.3.1.1.1.1.1.1.1	Variation 1: Reaction with an In Situ Generated Enal or Enone (The Doebner—von Miller Reaction)	391
15.3.1.1.1.1.1.1.2	Variation 2: Reaction with Pyruvic Acid and an Aldehyde (The Doebner Reaction)	392
15.3.1.1.1.1.1.1.3	Variation 3: Metal-Mediated Three-Component Coupling	393
15.3.1.1.1.1.1.2	Method 2: From Nitroarenes	394
15.3.1.1.1.1.2	With Formation of the 1—2, 3—4, and 4—4a Bonds	394
15.3.1.1.1.1.2.1	Method 1: From Arylamines	394
15.3.1.1.1.1.3	With Formation of the 1—8a Bond and Two C—C Bonds	395
15.3.1.1.1.1.3.1	Method 1: From Aryl Ketoximes	395

15.3.1.1.1.2	By Formation of Two N—C Bonds	396
15.3.1.1.1.2.1	Method 1: From 1-(2-Haloaryl)allyl Acetates	396
15.3.1.1.1.3	By Formation of One N—C and One C—C Bond	396
15.3.1.1.1.3.1	With Formation of the 1—2 and 2—3 Bonds	396
15.3.1.1.1.3.1.1	Method 1: From (2-Alkenylaryl)amines	396
15.3.1.1.1.3.1.2	Method 2: From (2-Acylaryl)amines	397
15.3.1.1.1.3.2	With Formation of the 1—2 and 3—4 Bonds	398
15.3.1.1.1.3.2.1	Method 1: From 2-Acylarylamines	398
15.3.1.1.1.3.2.1.1	Variation 1: Friedländer Synthesis	398
15.3.1.1.1.3.2.1.2	Variation 2: Oximino Derivatives of Aldehydes and Ketones	400
15.3.1.1.1.3.2.1.3	Variation 3: Reaction with Alkynes or Allenes	401
15.3.1.1.1.3.2.1.4	Variation 4: Reaction with Vinyl Sulfides	402
15.3.1.1.1.3.2.2	Method 2: From Isatins (The Pfitzinger Reaction)	402
15.3.1.1.1.3.2.3	Method 3: From <i>N</i> -(2-Aminobenzylidene)- <i>p</i> -toluidines (The Borsche Modification of the Friedländer Synthesis)	404
15.3.1.1.1.3.2.4	Method 4: From 2-(Perfluoroalkyl)arylamines	405
15.3.1.1.1.3.3	With Formation of the 1—2 and 4—4a Bonds	406
15.3.1.1.1.3.3.1	Method 1: From Arylamines	406
15.3.1.1.1.3.3.1.1	Variation 1: Reaction with Glycerol (The Skraup Reaction)	406
15.3.1.1.1.3.3.1.2	Variation 2: Reaction with Propenal Derivatives; Modified Skraup and Doebner–von Miller Reactions	408
15.3.1.1.1.3.3.1.3	Variation 3: Reaction with α,β -Unsaturated Ketones	410
15.3.1.1.1.3.3.1.4	Variation 4: Reaction with β -Heterosubstituted α,β -Unsaturated Aldehydes and Ketones	411
15.3.1.1.1.3.3.1.5	Variation 5: Ruthenium-Catalyzed Cyclization with Propane-1,3-diol or 1-Aminopropan-3-ol	413
15.3.1.1.1.3.3.1.6	Variation 6: From Cyclopropyl Building Blocks	414
15.3.1.1.1.3.3.1.7	Variation 7: Reaction with Malonic Acid	415
15.3.1.1.1.3.3.2	Method 2: From Nitroarenes	415
15.3.1.1.1.3.3.2.1	Variation 1: Reaction with Allyl Aryl Sulfones	415
15.3.1.1.1.3.3.2.2	Variation 2: Smiles Rearrangement of 3-(2-Nitrobenzoyl)pentane- 2,4-dione	416
15.3.1.1.1.3.3.3	Method 3: From (2-Haloaryl)amines	417
15.3.1.1.1.3.3.4	Method 4: From (2-Sulfanyaryl)amines	417
15.3.1.1.1.3.4	With Formation of the 1—8a and 4—4a Bonds	418
15.3.1.1.1.3.4.1	Method 1: From Quinones and Azadienes	418
15.3.1.1.1.4	By Formation of Two C—C Bonds	419
15.3.1.1.1.4.1	With Formation of the 2—3 and 4—4a Bonds	419
15.3.1.1.1.4.1.1	Method 1: From Arylidene <i>N</i> -Arylamines	419
15.3.1.1.1.4.1.1.1	Variation 1: Reaction with Alkenes	419
15.3.1.1.1.4.1.1.2	Variation 2: Reaction with Vinyl Ethers	420
15.3.1.1.1.4.1.1.3	Variation 3: Reaction with Alkynes	420
15.3.1.1.1.4.1.1.4	Variation 4: Cross Coupling with Allylamines	421
15.3.1.1.1.4.1.2	Method 2: From <i>N</i> -Arylimidates	422

15.3.1.1.1.4.1.3	Method 3:	From Aryl Isocyanides	423
15.3.1.1.1.4.1.4	Method 4:	From <i>N</i> -Arylnitrilium Salts	424
15.3.1.1.1.4.1.5	Method 5:	From (Anilinomethyl)phosphonates	425
15.3.1.1.1.4.1.6	Method 6:	From 3-(2-Acylaminophenylsulfanyl)cyclohex-2-enones	425
15.3.1.1.1.4.2		With Formation of the 3—4 and 4—4a Bonds	426
15.3.1.1.1.4.2.1	Method 1:	From Anilides	426
15.3.1.1.1.4.2.2	Method 2:	From <i>N</i> -Arylimines (Enamines)	426
15.3.1.1.1.5		By Formation of One N—C Bond	428
15.3.1.1.1.5.1		With Formation of the 1—2 Bond	428
15.3.1.1.1.5.1.1	Method 1:	From 2-(3-Hydroxyprop-1-ynyl)acetanilides	428
15.3.1.1.1.5.1.2	Method 2:	From β -(2-Aminoaryl)- α,β -ynones	429
15.3.1.1.1.5.1.3	Method 3:	From β -(2-Aminoaryl)- α,β -enones	430
15.3.1.1.1.5.1.4	Method 4:	From 2-Alkylarylamines	430
15.3.1.1.1.5.1.5	Method 5:	From β -(2-Nitroaryl)- α,β -enones	431
15.3.1.1.1.5.1.6	Method 6:	From β -(2-Nitroaryl)acrylates or (2-Nitroarylidene)acetates ..	432
15.3.1.1.1.5.1.7	Method 7:	From (2-Nitroaryl)dihydrocinnamoyl Derivatives	433
15.3.1.1.1.5.1.8	Method 8:	From 2-Azidophenyl α,β -Unsaturated Carbonyl Compounds ..	433
15.3.1.1.1.5.2		With Formation of the 1—8a Bond	434
15.3.1.1.1.5.2.1	Method 1:	From 3-Arylallylamines	434
15.3.1.1.1.5.2.2	Method 2:	From Oximes of β -Aryl- α,β -enones and β -Arylalkanones	434
15.3.1.1.1.6		By Formation of One C—C Bond	435
15.3.1.1.1.6.1		With Formation of the 2—3 Bond	435
15.3.1.1.1.6.1.1	Method 1:	From 2-Alkenylarylamines	435
15.3.1.1.1.6.1.1.1	Variation 1:	Activation of Azomethine Derivatives	435
15.3.1.1.1.6.1.1.2	Variation 2:	Activation of Amide Derivatives	436
15.3.1.1.1.6.1.1.3	Variation 3:	Cyclization via Isocyanides	437
15.3.1.1.1.6.1.2	Method 2:	From 2-Alkynylanilides	438
15.3.1.1.1.6.1.3	Method 3:	From 2-Alkynylphenyl Isocyanides	438
15.3.1.1.1.6.2		With Formation of the 3—4 Bond	439
15.3.1.1.1.6.2.1	Method 1:	From (2-Alkylphenyl)aminoethenes	439
15.3.1.1.1.6.2.2	Method 2:	From (2-Acylphenyl)aminoethenes	439
15.3.1.1.1.6.3		With Formation of the 4—4a Bond	440
15.3.1.1.1.6.3.1	Method 1:	From β -(Arylamino)- α,β -unsaturated Ketones and Aldehydes; the Beyer–Combes Synthesis	440
15.3.1.1.1.6.3.2	Method 2:	3-(Arylamino)alk-2-enamines	443
15.3.1.1.1.6.3.3	Method 3:	From Malonanilides and β -(Arylamino)methylenemalonates ..	443
15.3.1.1.1.6.3.4	Method 4:	From 3-Arylamino Ketones and Aldehydes	444
15.3.1.1.1.6.3.5	Method 5:	From α,β -Unsaturated <i>N</i> -Arylimines	444
15.3.1.1.1.6.3.6	Method 6:	From 3-Aryl-2-oxo-1,3-diazabicyclo[2.2.0]hex-5-enes	445
15.3.1.1.1.6.3.7	Method 7:	From <i>N</i> -Allyl-2-iodoanilines	446
15.3.1.1.2		By Annulation to the Heterocyclic Ring	446
15.3.1.1.2.1		By Formation of Two C—C Bonds	446
15.3.1.1.2.1.1		With Formation of the 5—6 and 7—8 Bonds	446

15.3.1.1.2.1.1	Method 1: From Dimethyl Pyridine-2,3-Dicarboxylate	446
15.3.1.1.2.1.2	With Formation of the 4–4a and 8–8a Bonds	447
15.3.1.1.2.1.1	Method 1: From 2,3-Dihalopyridines	447
15.3.1.1.2.2	By Formation of One C–C Bond	447
15.3.1.1.2.2.1	With Formation of the 5–6 Bond	447
15.3.1.1.2.2.1.1	Method 1: From Ethyl 3-(1-Ethoxyethoxy)-2-({3-[(phenylsulfonyl)methyl]-2-pyridyl)methyl}butanoate	447
15.3.1.1.2.2.2	With Formation of the 8–8a Bond	448
15.3.1.1.2.2.2.1	Method 1: From Azastilbenes	448
15.3.1.2	Synthesis by Ring Transformation	449
15.3.1.2.1	By Ring Enlargement	449
15.3.1.2.1.1	Method 1: From Indoles	449
15.3.1.2.1.2	Method 2: From 2,1-Benzisoxazole	449
15.3.1.2.1.3	Method 3: From Indazoles	450
15.3.1.2.2	Formal Exchange of Ring Members with Retention of the Ring Size	451
15.3.1.2.2.1	Method 1: From Benzotriazines	451
15.3.1.2.2.2	Method 2: From Benzoxazines	451
15.3.1.2.3	By Ring Contraction	452
15.3.1.3	Aromatization	452
15.3.1.3.1	Method 1: By Dehydrogenation	452
15.3.1.3.1.1	Variation 1: Air Oxidation	452
15.3.1.3.1.2	Variation 2: Oxidation with Quinone Reagents	453
15.3.1.3.1.3	Variation 3: Oxidation with Iodine Reagents	454
15.3.1.3.1.4	Variation 4: Metal-Mediated Oxidations	455
15.3.1.3.1.5	Variation 5: Dehydrogenation	457
15.3.1.3.2	Method 2: By Elimination Reactions	457
15.3.1.4	Synthesis by Substituent Modification	459
15.3.1.4.1	Substitution of Existing Substituents	459
15.3.1.4.1.1	Of Hydrogen	459
15.3.1.4.1.1.1	Method 1: Metalation	459
15.3.1.4.1.1.2	Method 2: Acylation	459
15.3.1.4.1.1.2.1	Variation 1: Reaction with Acyl Radicals	459
15.3.1.4.1.1.2.2	Variation 2: Electrophilic Acylation	461
15.3.1.4.1.1.3	Method 3: Alkylation and Arylation	462
15.3.1.4.1.1.3.1	Variation 1: Addition of Alkyl Radicals	462
15.3.1.4.1.1.3.2	Variation 2: Addition of Organometallic Compounds	463
15.3.1.4.1.1.3.3	Variation 3: Nucleophilic Addition	464
15.3.1.4.1.1.4	Method 4: Halogenation	465
15.3.1.4.1.1.5	Method 5: Sulfonylation	467
15.3.1.4.1.1.6	Method 6: Nitration	468
15.3.1.4.1.1.7	Method 7: Oxidation	469
15.3.1.4.1.2	Of Metals	470

15.3.1.4.1.2.1	Method 1:	Metal-Mediated Reductions of Heteroatoms	470
15.3.1.4.1.2.2	Method 2:	Substitution of Metalated Quinolines	472
15.3.1.4.1.2.3	Method 3:	Metal-Catalyzed Cross Coupling	474
15.3.1.4.1.3		Of Carbon Functionalities	478
15.3.1.4.1.4		Of Heteroatoms	479
15.3.1.4.1.4.1	Method 1:	From Quinoline 1-Oxides	479
15.3.1.4.1.4.1.1	Variation 1:	Deoxygenation to Quinolines	479
15.3.1.4.1.4.1.2	Variation 2:	Substitution with Concomitant Deoxygenation	480
15.3.1.4.1.4.2	Method 2:	Substitution of Heteroatoms in the 2- or 4-Position	483
15.3.1.4.1.4.2.1	Variation 1:	Halogenation of Quinolines	483
15.3.1.4.1.4.2.2	Variation 2:	Halogen Exchange	484
15.3.1.4.1.4.2.3	Variation 3:	Substitution with Carbon Nucleophiles	485
15.3.1.4.1.4.2.4	Variation 4:	Displacement with Oxygen Nucleophiles	485
15.3.1.4.1.4.2.5	Variation 5:	Displacement with Sulfur Nucleophiles	486
15.3.1.4.1.4.2.6	Variation 6:	Displacement with Nitrogen Nucleophiles	488
15.3.1.4.1.4.3	Method 3:	General Displacement Reactions	488
15.3.1.4.2		Rearrangement of Substituents	490
15.3.1.4.3		Modification of Substituents	490
15.3.1.4.3.1		Of C-Carbon Functionalities	490
15.3.1.4.3.1.1		Modification of Alkyl Groups	490
15.3.1.4.3.1.1.1	Method 1:	Alkylation	490
15.3.1.4.3.1.1.2	Method 2:	Halogenation of Alkyl Groups	492
15.3.1.4.3.1.1.3	Method 3:	Preparation of Oxidized Carbon Functionalities	493
15.3.1.4.3.1.1.3.1	Variation 1:	Oxidation of Methyl-Substituted Quinolines	493
15.3.1.4.3.2		Of C-Heteroatom Functionalities	495
15.3.1.4.3.2.1	Method 1:	Reactions of C-Oxygen Groups	495
15.3.1.4.3.2.1.1	Variation 1:	Alkylation of Quinolines	495
15.3.1.4.3.2.1.2	Variation 2:	Oxidation of Heterosubstituted Quinolines to the Quinolinediones	495
15.3.1.4.3.2.2	Method 2:	Reactions of C-Sulfur Groups	496
15.3.2		Product Subclass 2: Quinoline 1-Oxides	496
15.3.2.1		Synthesis by Ring-Closure Reactions	497
15.3.2.1.1		By Annulation to an Arene	497
15.3.2.1.1.1		By Formation of One N—C and One C—C Bond	497
15.3.2.1.1.1.1		With Formation of the 1—2 and 3—4 Bonds	497
15.3.2.1.1.1.1.1	Method 1:	From 2-Nitrobenzaldehydes	497
15.3.2.1.1.1.1.2	Method 2:	From 2-Nitrobenzyl Sulfones	498
15.3.2.1.1.2		By Formation of One N—C Bond	498
15.3.2.1.1.2.1		With Formation of the 1—2 Bond	498
15.3.2.1.1.2.1.1	Method 1:	From 2'-Nitrocinnamoyl Derivatives	498
15.3.2.1.1.2.1.2	Method 2:	From (2-Nitrobenzoyl)acetates	499

15.3.2.1.1.2.1.3	Method 3: From 1-(2-Nitrophenyl)allylic Alcohols (Baylis–Hillman Adducts)	500
15.3.2.1.1.2.1.4	Method 4: From 1-(2-Nitrophenyl)propenes	500
15.3.2.1.1.2.2	With Formation of the 1–8a Bond	501
15.3.2.1.1.2.2.1	Method 1: From Oximes and Alkoxyamines	501
15.3.2.2	Synthesis by Ring Transformation	502
15.3.2.3	Aromatization	503
15.3.2.4	Synthesis by Substituent Modification	503
15.3.2.4.1	Substitution of Existing Substituents	503
15.3.2.4.1.1	Of Hydrogen	503
15.3.2.4.1.1.1	Method 1: Metalation	503
15.3.2.4.1.1.2	Method 2: Alkylation	503
15.3.2.4.1.1.3	Method 3: Halogenation	504
15.3.2.4.1.1.4	Method 4: Nitration	504
15.3.2.4.1.1.5	Method 5: Oxidation	505
15.3.2.4.1.2	Of Heteroatoms	506
15.3.2.4.1.2.1	Method 1: Substitution of 4-Heterosubstituted Quinoline 1-Oxides	506
15.3.2.4.2	Addition Reactions	508
15.3.2.4.2.1	Method 1: N-Oxidation	508
15.3.2.4.3	Modification of Substituents	509
15.3.2.4.3.1	Of <i>N</i> -Heteroatom Functionalities	509
15.3.2.4.3.1.1	Method 1: O-Alkylation of Quinoline 1-Oxide	509
15.3.3	Product Subclass 3: 1-Alkyl- and 1-Arylquinolinium Salts	510
15.3.3.1	Synthesis by Ring-Closure Reactions	511
15.3.3.1.1	By Annulation to an Arene	511
15.3.3.1.1.1	By Formation of One N–C and Two C–C Bonds	511
15.3.3.1.1.1.1	Method 1: From 1-Aryl- or 1-Alkylanilines	511
15.3.3.1.1.2	By Formation of One N–C and One C–C Bond	512
15.3.3.1.1.2.1	With Formation of the 1–2 and 4–4a Bonds	512
15.3.3.1.1.2.1.1	Method 1: From <i>N</i> -Aryl- or <i>N</i> -Alkylarylamines	512
15.3.3.1.1.3	By Formation of Two C–C Bonds	513
15.3.3.1.1.3.1	With Formation of the 2–3 and 4–4a Bonds	513
15.3.3.1.1.3.1.1	Method 1: From <i>N</i> -Alkylanilides	513
15.3.3.1.1.3.1.2	Method 2: From <i>N</i> -Alkyl- <i>N</i> -(benzotriazol-1-ylmethyl)arylamines	513
15.3.3.1.1.3.2	With Formation of the 3–4 and 4–4a Bonds	514
15.3.3.1.1.3.2.1	Method 1: From <i>N</i> -Arylenamines	514
15.3.3.1.1.4	By Formation of One C–C Bond	515
15.3.3.1.1.4.1	Method 1: From β -Arylamino Ketones	515

15.3.3.2	Synthesis by Ring Transformation	515
15.3.3.3	Aromatization	516
15.3.3.4	Synthesis by Substituent Modification	517
15.3.3.4.1	Substitution of Existing Substituents	517
15.3.3.4.1.1	Of Metal	517
15.3.3.4.1.2	Of Carbon Functionalities	517
15.3.3.4.1.3	Of Heteroatoms	517
15.3.3.4.1.3.1	Method 1: From Quinoline 1-Oxides	517
15.3.3.4.1.3.2	Method 2: From 2- or 4-Heterosubstituted Quinolinium Intermediates ..	517
15.3.3.4.2	Addition Reactions	518
15.3.3.4.2.1	Method 1: N-Alkylation	518
15.3.3.4.3	Modification of Substituents	519
15.3.3.4.3.1	Of C-Carbon Functionalities	519
15.3.3.4.3.1.1	Method 1: Alkylation Reactions	519
15.4	Product Class 4: Quinolinones R. D. Larsen	
15.4	Product Class 4: Quinolinones	551
15.4.1	Product Subclass 1: Quinolin-2(1H)-ones	551
15.4.1.1	Synthesis by Ring-Closure Reactions	553
15.4.1.1.1	By Annulation to an Arene	553
15.4.1.1.1.1	By Formation of One N—C and One C—C Bond	553
15.4.1.1.1.1.1	With Formation of the 1—2 and 2—3 Bonds	553
15.4.1.1.1.1.1.1	Method 1: From 2-Aminostyrenes	553
15.4.1.1.1.1.1.2	Method 2: From (2-Acylaryl)amines	553
15.4.1.1.1.1.2	With Formation of the 1—2 and 3—4 Bonds	554
15.4.1.1.1.1.2.1	Method 1: From (2-Acylaryl)amines	554
15.4.1.1.1.1.2.1.1	Variation 1: Reaction with Acetates: The Friedländer Quinolinone Synthesis	554
15.4.1.1.1.1.2.1.2	Variation 2: Reaction with Diethoxyphosphorylketenes	555
15.4.1.1.1.1.2.2	Method 2: From Anils of 2-Aminobenzaldehydes	555
15.4.1.1.1.1.2.3	Method 3: From Anthranilates or 2-Aminobenzonitriles	556
15.4.1.1.1.1.2.4	Method 4: From 2-Tolylformamides	557
15.4.1.1.1.1.3	With Formation of the 1—2 and 4—4a Bonds	557
15.4.1.1.1.1.3.1	Method 1: From Arylamines	557
15.4.1.1.1.1.3.2	Method 2: From (2-Haloaryl)amines	558
15.4.1.1.1.1.4	With Formation of the 1—8a and 3—4 Bonds	559
15.4.1.1.1.1.4.1	Method 1: From 2-Halobenzaldehydes	559
15.4.1.1.1.2	By Formation of Two C—C Bonds	559

15.4.1.1.1.2.1	With Formation of the 2–3 and 4–4a Bonds	559
15.4.1.1.1.2.1.1	Method 1: From Aryl Isocyanates	559
15.4.1.1.1.2.1.1.1	Variation 1: Reaction with Alkynes	559
15.4.1.1.1.2.1.1.2	Variation 2: Reaction with Triphenylphosphonium Malonate	560
15.4.1.1.1.2.2	With Formation of the 3–4 and 4–4a Bonds	560
15.4.1.1.1.2.2.1	Method 1: From <i>N</i> -Acylarylamines	560
15.4.1.1.1.3	By Formation of One N–C Bond	561
15.4.1.1.1.3.1	With Formation of the 1–2 Bond	561
15.4.1.1.1.3.1.1	Method 1: From 2-Aminocinnamates	561
15.4.1.1.1.3.1.2	Method 2: From a 3-(2-Acylaminophenyl)-3-hydroxypropanoate	561
15.4.1.1.1.3.1.3	Method 3: From 3-{2-[(Alkoxy carbonyl)amino]phenyl}-3-hydroxy-acrylates	562
15.4.1.1.1.3.1.4	Method 4: From 2-Nitrocinnamates	563
15.4.1.1.1.3.1.5	Method 5: From Methyl 2-[(Acetyloxy)(2-nitrophenyl)methyl]acrylate	563
15.4.1.1.1.3.1.6	Method 6: From 2-Nitrobenzoylacetates	564
15.4.1.1.1.4	By Formation of One C–C Bond	565
15.4.1.1.1.4.1	With Formation of the 2–3 Bond	565
15.4.1.1.1.4.1.1	Method 1: From 2-Alkenylaryl Isocyanates	565
15.4.1.1.1.4.1.2	Method 2: From (2-Acylphenyl)carbamates	565
15.4.1.1.1.4.2	With Formation of the 3–4 Bond	566
15.4.1.1.1.4.2.1	Method 1: From 2-Acylanilides: Camps Synthesis	566
15.4.1.1.1.4.2.2	Method 2: From <i>N</i> -Acylanthranilates or 2-(Cyanoaryl)acylamides	567
15.4.1.1.1.4.2.3	Method 3: From 2-Alkynylmalonanilides	568
15.4.1.1.1.4.2.4	Method 4: From 2-(α -Oxoacylamino)benzylphosphonium Salts	569
15.4.1.1.1.4.3	With Formation of the 4–4a Bond	570
15.4.1.1.1.4.3.1	Method 1: From Acylacetanilides: Knorr Synthesis	570
15.4.1.1.1.4.3.2	Method 2: From Malonanilides or Malondianilides	571
15.4.1.1.1.4.3.3	Method 3: From Acrylanilides	572
15.4.1.1.1.4.3.4	Method 4: From <i>N</i> -Aryl-3-(arylsulfinyl)propanamides or <i>N</i> -Aryl-3-(arylsulfonyl)propanamides	573
15.4.1.1.1.4.3.5	Method 5: From Alkynylanilides	574
15.4.1.1.1.4.3.6	Method 6: From <i>N</i> -(2-Bromophenyl)acrylamides	574
15.4.1.1.2	By Annulation to the Heterocyclic Ring	575
15.4.1.2	Synthesis by Ring Transformation	575
15.4.1.2.1	By Ring Enlargement	575
15.4.1.2.1.1	Method 1: From Isatin	575
15.4.1.2.1.2	Method 2: From Indoles	576
15.4.1.2.2	Formal Exchange of Ring Members with Retention of the Ring Size	577
15.4.1.2.2.1	Method 1: From Isoic Anhydrides (Benzoxazinones)	577
15.4.1.2.3	By Ring Contraction	578
15.4.1.3	Aromatization	578

15.4.1.3.1	Method 1: By Dehydrogenation	578
15.4.1.3.2	Method 2: Elimination Reactions	579
15.4.1.4	Synthesis by Substituent Modification	579
15.4.1.4.1	Substitution of Existing Substituents	579
15.4.1.4.1.1	Of <i>N</i> -Hydrogen	579
15.4.1.4.1.2	Of <i>C</i> -Hydrogen	580
15.4.1.4.1.2.1	Method 1: By Metals	580
15.4.1.4.1.2.2	Method 2: By Alkyl Groups	580
15.4.1.4.1.2.3	Method 3: By Halogens	581
15.4.1.4.1.2.4	Method 4: By Hydroxy Groups	581
15.4.1.4.1.2.5	Method 5: Sulfanylation	582
15.4.1.4.1.2.6	Method 6: Nitration	582
15.4.1.4.1.3	Of Metals	583
15.4.1.4.1.4	Of Carbon	584
15.4.1.4.1.5	Of Heteroatoms	584
15.4.1.4.1.5.1	Method 1: From Quinoline <i>N</i> -Oxides	584
15.4.1.4.1.5.2	Method 2: Reactions at the 2-Position To Generate Quinolinones	585
15.4.1.4.1.5.3	Method 3: Substitution of 4-Heterosubstituted Quinolin-2-ones	585
15.4.1.4.2	Modification of Substituents	586
15.4.1.4.2.1	Of <i>C</i> -Carbon Functionalities	586
15.4.1.4.2.2	Of <i>C</i> -Heteroatom Functionalities	586
15.4.2	Product Subclass 2: Quinolin-4(1<i>H</i>)-ones	587
15.4.2.1	Synthesis by Ring-Closure Reactions	588
15.4.2.1.1	By Annulation to an Arene	588
15.4.2.1.1.1	By Formation of One <i>N</i> — <i>C</i> and Two <i>C</i> — <i>C</i> Bonds	588
15.4.2.1.1.1.1	With Formation of the 1—2, 3—4, and 4—4a Bonds	588
15.4.2.1.1.1.1.1	Method 1: From 2-Haloarylamines	588
15.4.2.1.1.2	By Formation of One <i>N</i> — <i>C</i> and One <i>C</i> — <i>C</i> Bond	588
15.4.2.1.1.2.1	With Formation of the 1—2 and 2—3 Bonds	588
15.4.2.1.1.2.1.1	Method 1: From (2-Acylaryl)amines	588
15.4.2.1.1.2.2	With Formation of the 1—2 and 3—4 Bonds	589
15.4.2.1.1.2.2.1	Method 1: From 2-Aminobenzaldehydes	589
15.4.2.1.1.2.2.2	Method 2: From Anthranilates	590
15.4.2.1.1.2.2.2.1	Variation 1: Reaction with Ketones (The Niementowski Reaction)	590
15.4.2.1.1.2.2.2.2	Variation 2: Reaction with Ketene Equivalents	591
15.4.2.1.1.2.3	With Formation of the 1—2 and 4—4a Bonds	592
15.4.2.1.1.2.3.1	Method 1: From Arylamines	592
15.4.2.1.1.3	By Formation of Two <i>C</i> — <i>C</i> Bonds	593
15.4.2.1.1.3.1	With Formation of the 2—3 and 3—4 Bonds	593

15.4.2.1.1.3.1.1	Method 1: From 2-Isocyanobenzoates	593
15.4.2.1.1.3.1.2	Method 2: From 2-Phthalimidobenzoic Acid	593
15.4.2.1.1.3.2	With Formation of the 2–3 and 4–4a Bonds	594
15.4.2.1.1.3.2.1	Method 1: From Formanilides	594
15.4.2.1.1.3.3	With Formation of the 3–4 and 4–4a Bonds	595
15.4.2.1.1.3.3.1	Method 1: From Anilides or <i>N</i> -Arylenamines	595
15.4.2.1.1.3.3.2	Method 2: From <i>N</i> -(2-Haloaryl)enamines	595
15.4.2.1.1.4	By Formation of One N–C Bond	596
15.4.2.1.1.4.1	With Formation of the 1–2 Bond	596
15.4.2.1.1.4.1.1	Method 1: From (2-Aminobenzoyl)alkanes or (2-Aminobenzoyl)alkenes	596
15.4.2.1.1.4.1.2	Method 2: From (2-Nitrobenzoyl)alkanes or (2-Nitrobenzoyl)alkenes	597
15.4.2.1.1.4.2	With Formation of the 1–8a Bond	598
15.4.2.1.1.4.2.1	Method 1: From 3-Amino-1-(2-halophenyl)- or 3-Amino-1-(2-nitrophenyl)alkenes: Cycloaracylation Process (The Grohe Method)	598
15.4.2.1.1.5	Formation of One C–C Bond	599
15.4.2.1.1.5.1	With Formation of the 2–3 Bond	599
15.4.2.1.1.5.1.1	Method 1: From 2-Acylanilides	599
15.4.2.1.1.5.2	With Formation of the 3–4 Bond	599
15.4.2.1.1.5.2.1	Method 1: From <i>N</i> -[2-(Alkoxy carbonyl)aryl]enamines	599
15.4.2.1.1.5.2.2	Method 2: From <i>N</i> -(2-Trifluoromethylphenyl)ketimines	600
15.4.2.1.1.5.3	With Formation of the 4–4a Bond	601
15.4.2.1.1.5.3.1	Method 1: From β -(Arylamino)acrylates: The Conrad–Limpach Synthesis (Gould–Jacobs Modification)	601
15.4.2.1.1.5.3.2	Method 2: From <i>N</i> -Arylketenimines	603
15.4.2.1.1.5.3.3	Method 3: From <i>N</i> -Aryl Heterocycles	603
15.4.2.2	Synthesis by Ring Transformation	604
15.4.2.2.1	By Ring Enlargement	604
15.4.2.2.2	Formal Exchange of Ring Members with Retention of the Ring Size	604
15.4.2.3	Aromatization	605
15.4.2.3.1	Method 1: Dehydrogenation of Dihydroquinolin-4(1 <i>H</i>)-ones	605
15.4.2.3.2	Method 2: Elimination Reactions	606
15.4.2.4	Synthesis by Substituent Modification	607
15.4.2.4.1	Substitution of Existing Substituents	607
15.4.2.4.1.1	Of <i>N</i> -Hydrogen	607
15.4.2.4.1.2	Of <i>C</i> -Hydrogen	608
15.4.2.4.1.2.1	Method 1: By Metals	608
15.4.2.4.1.2.2	Method 2: By Alkyl Groups	608
15.4.2.4.1.2.3	Method 3: By Halogens	609
15.4.2.4.1.2.4	Method 4: Hydroxylation	609
15.4.2.4.1.2.5	Method 5: Nitration	610

15.4.2.4.1.3	Of Metals	610
15.4.2.4.1.4	Of Carbon Functionalities	611
15.4.2.4.1.5	Of Heteroatoms	612
15.4.2.4.1.5.1	Method 1: Hydrolysis of 4-Heterosubstituted Quinolines	612
15.4.2.4.1.5.2	Method 2: Substitution of 7-Haloquinolin-4(1 <i>H</i>)-ones	613
15.4.2.4.2	Modification of Substituents	614
15.4.2.4.2.1	Of C-Heteroatom Functionalities	614
15.4.2.4.2.1.1	Method 1: Of C-Oxygen Quinolines	614
15.4.2.4.2.1.1.1	Variation 1: Dealkylation of 4-Alkoxyquinolines	614
15.4.2.4.2.1.1.2	Variation 2: Oxidation of 5,8-Dihydroxyquinolin-4(1 <i>H</i>)-one to Quinoline-4,5,8(1 <i>H</i>)-trione	614
15.4.3	Product Subclass 3: Quinoline-2(1<i>H</i>)-thiones and Quinoline-4(1<i>H</i>)-thiones	614
15.4.3.1	Synthesis by Ring-Closure Reactions	615
15.4.3.1.1	By Annulation to an Arene	615
15.4.3.1.1.1	By Formation of One N—C and One C—C Bond	615
15.4.3.1.1.1.1	With Formation of the 1—2 and 2—3 Bonds	615
15.4.3.1.1.1.1.1	Method 1: From (2-Alkenylaryl)amines	615
15.4.3.1.1.2	By Formation of One C—C Bond	616
15.4.3.1.1.2.1	With Formation of the 4—4a Bond	616
15.4.3.1.1.2.1.1	Method 1: From Diethyl (Anilinothioyl)malonates	616
15.4.3.2	Synthesis by Ring Transformation	616
15.4.3.2.1	Formal Exchange of Ring Members with Retention of the Ring Size	616
15.4.3.2.2	By Ring Contraction	616
15.4.3.3	Synthesis by Substituent Modification	617
15.4.3.3.1	Substitution of Existing Substituents	617
15.4.3.3.1.1	Of Metals	617
15.4.3.3.1.2	Of Heteroatoms	617
15.4.3.3.1.2.1	Method 1: Conversion of 2- or 4-Heterosubstituted Quinolines into Quinolinethiones	617
15.4.4	Product Subclass 4: Amine Derivatives of Quinolines	618
15.4.4.1	Synthesis by Ring-Closure Reactions	619
15.4.4.1.1	By Annulation to an Arene	619
15.4.4.1.1.1	By Formation of One N—C and Two C—C Bonds	619
15.4.4.1.1.1.1	With Formation of the 1—2, 3—4, and 4—4a Bonds	619
15.4.4.1.1.1.1.1	Method 1: From <i>N</i> -Aryl Isocyanates, <i>N</i> -Arylcarbodiimides, or <i>N</i> -Arylthioureas	619
15.4.4.1.1.2	By Formation of One N—C and One C—C Bond	619
15.4.4.1.1.2.1	With Formation of the 1—2 and 2—3 Bonds	619

15.4.4.1.1.2.1.1	Method 1:	From 2-Acylaryl Azides	619
15.4.4.1.1.2.1.2	Method 2:	From (2-Alkynylaryl)amines	620
15.4.4.1.1.2.2	With Formation of the 1–2 and 3–4 Bonds		621
15.4.4.1.1.2.2.1	Method 1:	From (2-Acylaryl)amines	621
15.4.4.1.1.2.2.1.1	Variation 1:	Reaction with Alkylcarbonitriles	621
15.4.4.1.1.2.2.1.2	Variation 2:	Reaction with Amide Derivatives	621
15.4.4.1.1.2.2.2	Method 2:	From 2-(Perfluoroalkyl)anilines	622
15.4.4.1.1.2.2.3	Method 3:	From Anthranilates	623
15.4.4.1.1.2.2.4	Method 4:	From (2-Cyanoaryl)amines	623
15.4.4.1.1.2.2.5	Method 5:	From (2-Acylaryl)azides	624
15.4.4.1.1.2.3	With Formation of the 1–2 and 4–4a Bonds		625
15.4.4.1.1.2.3.1	Method 1:	From Arylamines	625
15.4.4.1.1.2.3.2	Method 2:	From <i>N</i> -Arylhydroxylamines	625
15.4.4.1.1.3	By Formation of Two C–C Bonds		626
15.4.4.1.1.3.1	With Formation of the 2–3 and 4–4a Bonds		626
15.4.4.1.1.3.1.1	Method 1:	From <i>N</i> -Arylketenimines	626
15.4.4.1.1.3.2	With Formation of the 3–4 and 4–4a Bonds		626
15.4.4.1.1.3.2.1	Method 1:	From Anilides	626
15.4.4.1.1.3.2.2	Method 2:	From <i>N</i> -Arylimidoylphosphoranes	627
15.4.4.1.1.4	By Formation of One N–C Bond		627
15.4.4.1.1.4.1	With Formation of the 1–2 Bond		627
15.4.4.1.1.4.1.1	Method 1:	From 2-Nitrocinnamionitriles	627
15.4.4.1.1.4.1.2	Method 2:	From 3-(2-Aminophenyl)-3-hydroxypropanenitrile	628
15.4.4.1.1.4.2	With Formation of the 1–8a Bond		628
15.4.4.1.1.4.2.1	Method 1:	From 3-Amino-3-arylalk-2-enamines	628
15.4.4.1.1.4.2.2	Method 2:	From 3-(2-Haloaryl)prop-3-en-1-amines	629
15.4.4.1.1.5	By Formation of One C–C Bond		629
15.4.4.1.1.5.1	With Formation of the 2–3 Bond		629
15.4.4.1.1.5.1.1	Method 1:	From <i>N</i> -(2-Alkenylphenyl)- <i>N'</i> -arylcarbodiimides	629
15.4.4.1.1.5.1.2	Method 2:	From <i>N</i> -(2-Vinylaryl)ureas	630
15.4.4.1.1.5.1.3	Method 3:	From 2-Alkynylphenyl Isocyanates	631
15.4.4.1.1.5.2	With Formation of the 3–4 Bond		631
15.4.4.1.1.5.2.1	Method 1:	From <i>N</i> -(2-Acylaryl)imidates/Amidines	631
15.4.4.1.1.5.2.2	Method 2:	From <i>N</i> -Acetylanthranilates	632
15.4.4.1.1.5.2.3	Method 3:	From 2-(Ethylideneamino)benzonitriles	632
15.4.4.1.1.5.2.4	Method 4:	From <i>N</i> -(2-Perfluoroalkyl)phenylketimines	633
15.4.4.1.1.5.3	With Formation of the 4–4a Bond		634
15.4.4.1.1.5.3.1	Method 1:	From Bis(arylamino) Enones and Acrylates	634
15.4.4.1.1.5.3.2	Method 2:	From 3-(Arylamino)acrylonitriles or 3-(Arylamino)acrylamides	635
15.4.4.1.1.5.3.3	Method 3:	From Propynimidamides	635
15.4.4.1.2	By Annulation to the Heterocyclic Ring		636

15.4.4.1.2.1	By Formation of Two C—C Bonds	636
15.4.4.1.2.1.1	Method 1: From 2-Amino-6-methylpyridine-3,4,5-tricarbonitrile	636
15.4.4.2	Synthesis by Ring Transformation	637
15.4.4.2.1	By Ring Enlargement	637
15.4.4.2.2	Formal Exchange of Ring Members with Retention of the Ring Size	637
15.4.4.2.3	By Ring Contraction	638
15.4.4.3	Synthesis by Substituent Modification	638
15.4.4.3.1	Substitution of Existing Substituents	638
15.4.4.3.1.1	Of Hydrogen	638
15.4.4.3.1.1.1	Method 1: Amination (The Chichibabin Reaction)	638
15.4.4.3.1.2	Of Metals	639
15.4.4.3.1.3	Of Carbon Functionalities	639
15.4.4.3.1.4	Of Heteroatoms	640
15.4.4.3.1.4.1	Method 1: From Quinoline 1-Oxides	640
15.4.4.3.1.4.2	Method 2: Substitution of Heteroatoms in the 2- or 4-Position	640
15.4.4.3.2	Rearrangement of Substituents	642
15.5	Product Class 5: Isoquinolines M. Álvarez and J. A. Joule	
15.5	Product Class 5: Isoquinolines	661
15.5.1	Product Subclass 1: Isoquinolines	661
15.5.1.1	Synthesis by Ring-Closure Reactions	664
15.5.1.1.1	By Annulation to an Arene	665
15.5.1.1.1.1	By Formation of Two N—C Bonds	665
15.5.1.1.1.1.1	With Formation of the 1—2 and 2—3 Bonds	665
15.5.1.1.1.1.1.1	Method 1: From 1,5-Dicarbonyl Compounds	665
15.5.1.1.1.1.1.1.1	Variation 1: From 2-Formylbenzylcarbonyl Compounds	665
15.5.1.1.1.1.1.1.2	Variation 2: From 1,5-Diketones	666
15.5.1.1.1.1.1.1.3	Variation 3: From 2-[(Cyclohexylimino)methyl]benzyl Ketones	667
15.5.1.1.1.2	By Formation of One N—C and One C—C Bond	668
15.5.1.1.1.2.1	With Formation of the 2—3 and 1—8a Bonds	668
15.5.1.1.1.2.1.1	Method 1: From Arylacetic Acids and Derivatives	668
15.5.1.1.1.2.1.1.1	Variation 1: From Arylacetic Esters	668
15.5.1.1.1.2.1.1.2	Variation 2: From Arylacetamides	668
15.5.1.1.1.2.1.2	Method 2: From 1-Aryl-2-bromoalkenes and a Nitrile	670
15.5.1.1.1.2.1.3	Method 3: From 3-Arylprop-1-enes and a Nitrile	670
15.5.1.1.1.2.1.4	Method 4: From Benzyl Ketones	671
15.5.1.1.1.2.1.4.1	Variation 1: By Gatterman Reaction	671
15.5.1.1.1.2.1.4.2	Variation 2: Via <i>ortho</i> Lithiation	672

15.5.1.1.1.2.2	With Formation of the 2—3 and 4—4a Bonds	672
15.5.1.1.1.2.2.1	Method 1: From 2-Iodobenzylamine	672
15.5.1.1.1.2.2.2	Method 2: From <i>N</i> -Acetyl-2-iodobenzylamine	673
15.5.1.1.1.2.2.3	Method 3: From 1-(2-Iodoaryl)alkan-1-imines	673
15.5.1.1.1.2.2.3.1	Variation 1: From <i>N</i> - <i>tert</i> -Butyl-2-iodoarylmethanimines and Alkynes	673
15.5.1.1.1.2.2.3.2	Variation 2: From 1-(2-Iodophenyl)alkan-1-imines and Allenes	674
15.5.1.1.1.2.2.4	Method 4: From 1-Arylalkan-1-imines	675
15.5.1.1.1.2.3	With Formation of the 1—2 and 1—8a Bonds	676
15.5.1.1.1.2.3.1	Method 1: From Arylacetonitriles	676
15.5.1.1.1.2.3.1.1	Variation 1: Synthesis of 1,3-Dichloroisoquinolines Using Phosgene	676
15.5.1.1.1.2.3.1.2	Variation 2: Synthesis of 3-Chloroisoquinolines Using a Vilsmeier Reaction	676
15.5.1.1.1.2.3.2	Method 2: Synthesis of 1,2,3,4-Tetrahydroisoquinolines: The Pictet–Spengler Synthesis	677
15.5.1.1.1.2.3.2.1	Variation 1: Pictet–Spengler Synthesis Using a Superacid	679
15.5.1.1.1.2.3.2.2	Variation 2: Using an Acyl Chloride	680
15.5.1.1.1.2.3.2.3	Variation 3: Using 2-Aryl- <i>N</i> -(arylsulfonyl)ethylamines	680
15.5.1.1.1.2.3.2.4	Variation 4: Using 2-Aryl- <i>N</i> -(phenylsulfonyl)ethylamines with Ethyl Chloro(methylsulfanyl)- or Chloro(phenylselanyl)acetates	681
15.5.1.1.1.2.3.2.5	Variation 5: Using <i>N</i> -Acyl- or <i>N</i> -(Alkoxy carbonyl)-2-arylethylamines	682
15.5.1.1.1.2.3.2.6	Variation 6: Using Carbonyl Component Synthons	684
15.5.1.1.1.2.3.2.7	Variation 7: Generation of Intermediates Using Aza-Wittig Reactions	686
15.5.1.1.1.2.3.3	Method 3: From 2-Arylethylamines, Carboxylic Acids, and Polyphosphoric Acid	688
15.5.1.1.1.2.3.4	Method 4: Stereo- and Enantioselective Pictet–Spengler-Type Syntheses of 1,2,3,4-Tetrahydroisoquinolines	688
15.5.1.1.1.2.4	With Formation of the 1—2 and 3—4 Bonds	691
15.5.1.1.1.2.4.1	Method 1: From 1,2-Diacylarenes	692
15.5.1.1.1.2.4.1.1	Variation 1: Using Diethyl [(Triphenylphosphoranylidene)amino]methyl- phosphonate	692
15.5.1.1.1.2.4.1.2	Variation 2: From 1,2-Diacylarenes and an Iminophosphorane Generated In Situ	692
15.5.1.1.1.2.4.2	Method 2: From <i>ortho</i> -Substituted Arenecarbonitriles and Arenecarboxamides	693
15.5.1.1.1.2.4.2.1	Variation 1: From (2-Cyanophenyl)acetonitrile	693
15.5.1.1.1.2.4.2.2	Variation 2: From 2-Methylbenzonitriles	693
15.5.1.1.1.2.4.2.3	Variation 3: From 3-(2-Cyanophenyl)acrylamide	694
15.5.1.1.1.2.4.2.4	Variation 4: From 2-Methylbenzamides	694
15.5.1.1.1.3	By Formation of Two C—C Bonds	694
15.5.1.1.1.3.1	With Formation of the 1—8a and 4—4a Bonds	694
15.5.1.1.1.3.1.1	Method 1: From 2-Aza-1,3-dienes	694
15.5.1.1.1.3.1.2	Method 2: From Heterocyclic 2-Aza-1,3-dienes	695
15.5.1.1.1.4	By Formation of One N—C Bond	696
15.5.1.1.1.4.1	With Formation of the 1—2 Bond	696
15.5.1.1.1.4.1.1	Method 1: From (2-Formylphenyl)acetonitriles	696

15.5.1.1.1.4.1.2	Method 2:	From 3-(2-Acylphenyl)-2-azidoacrylates with Phosphorus(III) Reagents: Intramolecular Aza-Wittig Reactions	696
15.5.1.1.1.4.1.3	Method 3:	From 2-(2-Acylaryl)ethylamines	697
15.5.1.1.1.4.2		With Formation of the 2–3 Bond	697
15.5.1.1.1.4.2.1	Method 1:	From 1-[2-[(<i>tert</i> -Butylimino)methyl]phenyl]alk-1-yne	697
15.5.1.1.1.4.2.1.1	Variation 1:	With Proton, Iodine, Sulfur, or Selenium Electrophiles	697
15.5.1.1.1.4.2.1.2	Variation 2:	With Carbon Monoxide and an Electrophile	698
15.5.1.1.1.4.2.1.3	Variation 3:	Vinylation by the Heck Reaction	699
15.5.1.1.1.4.2.1.4	Variation 4:	With Addition of Aryl or Alkyl Halides	699
15.5.1.1.1.4.2.2	Method 2:	Palladium(0)-Catalyzed Cyclization of 2-Allylbenzaldehyde <i>O</i> -(Pentafluorobenzoyl)oximes	700
15.5.1.1.1.4.2.3	Method 3:	Electrocyclization of 2-Vinylbenzaldehyde Oximes	700
15.5.1.1.1.4.2.4	Method 4:	From <i>N</i> -Acetyl-2-(2-acylaryl)ethylamines	701
15.5.1.1.1.4.2.5	Method 5:	From 2-(1-Cyano-2-phenylvinyl)benzoxonitrile	702
15.5.1.1.1.4.2.6	Method 6:	From 2-(2,2-Difluorovinyl)benzoxonitriles	702
15.5.1.1.1.4.2.7	Method 7:	From (2-Cyanophenyl)acetoxonitriles	703
15.5.1.1.1.4.2.8	Method 8:	From 2-(Cyanomethyl)benzoic Acids	703
15.5.1.1.1.4.2.9	Method 9:	From 2-Vinylbenzaldehyde Imines	704
15.5.1.1.1.4.2.10	Method 10:	From Sulfinamides	705
15.5.1.1.1.4.2.11	Method 11:	From Chiral 2-[(<i>Tosylamino</i>)methyl]phenylacetaldehyde Acetals	705
15.5.1.1.1.4.2.12	Method 12:	From a Protected 2-(2-Oxoalkyl)benzoxonitrile	706
15.5.1.1.1.5		By Formation of One C–C Bond	706
15.5.1.1.1.5.1		With Formation of the 4–4a Bond	706
15.5.1.1.1.5.1.1	Method 1:	From 2,2-Dialkoxy- <i>N</i> -benzylideneethylamine: The Pomeranz–Fritsch Synthesis	706
15.5.1.1.1.5.1.1.1	Variation 1:	From 1-Arylalkylamines and Glyoxal Monoacetal	708
15.5.1.1.1.5.1.1.2	Variation 2:	Pomeranz–Fritsch Syntheses Using Alternative Acids for the Cyclizations	709
15.5.1.1.1.5.1.1.3	Variation 3:	Pomeranz–Fritsch Synthesis Using Ethyl Chloroformate/Trialkyl Phosphite Followed by Titanium(IV) Chloride	709
15.5.1.1.1.5.1.1.4	Variation 4:	From 2,2-Dialkoxy- <i>N</i> -benzylethylamines in the Presence of an Aldehyde	711
15.5.1.1.1.5.1.1.5	Variation 5:	From 2,2-Dialkoxy- <i>N</i> -(1-Arylalkyl)ethylamines Using Chlorosulfonic Acid	712
15.5.1.1.1.5.1.1.6	Variation 6:	From 2,2-Dialkoxy- <i>N</i> -benzyl- <i>N</i> -tosylethylamines	712
15.5.1.1.1.5.1.1.7	Variation 7:	From 2,2-Dialkoxy- <i>N</i> -(1-arylalkyl)- <i>N</i> -(benzylsulfonyl)ethylamines	714
15.5.1.1.1.5.1.1.8	Variation 8:	From <i>N</i> -Acetyl-2,2-dialkoxy- <i>N</i> -benzylethylamines	715
15.5.1.1.1.5.1.1.9	Variation 9:	From (Benzylamino)acetates	715
15.5.1.1.1.5.1.2	Method 2:	Stereo- and Enantioselective Syntheses of 1,2,3,4-Tetrahydroisoquinolines with Formation of the 4–4a Bond	716
15.5.1.1.1.5.2		With Formation of the 1–8a Bond	716
15.5.1.1.1.5.2.1	Method 1:	From <i>N</i> -Acyl-2-arylethylamines: The Bischler–Napieralski Synthesis	716

15.5.1.1.1.5.2.1.1	Variation 1:	From <i>N</i> -Acyl-2-aryl-2-hydroxyethylamines: Pictet–Gams Modification of the Bischler–Napieralski Synthesis	720
15.5.1.1.1.5.2.1.2	Variation 2:	From 5-Aryl-4,5-dihydrooxazoles	721
15.5.1.1.1.5.2.1.3	Variation 3:	From 3-Arylpropanone Oximes	722
15.5.1.1.1.5.2.1.4	Variation 4:	From <i>N</i> -Acyl-2-arylvinylamines	723
15.5.1.1.1.5.2.1.5	Variation 5:	Photolysis of <i>N</i> -Acyl-2-arylvinylamines	724
15.5.1.1.1.5.2.1.6	Variation 6:	From 4-Arylbut-3-en-2-one Oximes	724
15.5.1.1.1.5.2.1.7	Variation 7:	From 2-Aryl- <i>N</i> -(thioacyl)ethylamines	725
15.5.1.1.1.5.2.2	Method 2:	From <i>N</i> -Substituted <i>N'</i> -(1,2-Diarylvinyl)carbodiimides	725
15.5.1.1.1.5.2.3	Method 3:	From 2-Arylethyl Isocyanides	726
15.5.1.1.1.5.2.4	Method 4:	Synthesis of Enantiomerically Pure 1,2,3,4-Tetrahydroisoquinolines via Bischler–Napieralski Closure	727
15.5.1.1.2		By Annulation to a Pyridine or Reduced Pyridine	728
15.5.1.1.2.1		With Formation of the 4a–5 and 8–8a Bonds	728
15.5.1.1.2.1.1	Method 1:	From a 3,4-Didehydropyridine	728
15.5.1.1.2.1.2	Method 2:	From 3,4-Bis(chloromethyl)pyridines	728
15.5.1.1.2.2		With Formation of the 6–7 and 8–8a Bonds	729
15.5.1.1.2.2.1	Method 1:	From a 4-Vinyl-1,2,3,6-tetrahydropyridine	729
15.5.1.1.2.3		With Formation of the 5–6 and 7–8 Bonds	730
15.5.1.1.2.3.1	Method 1:	From a Piperidin-4-one-Derived Enamine	730
15.5.1.2		Synthesis by Ring Transformation	731
15.5.1.2.1	Method 1:	Dehydrogenation of 1,2,3,4-Tetrahydroisoquinolines	731
15.5.1.2.1.1	Variation 1:	Using Sulfur	731
15.5.1.2.1.2	Variation 2:	Using Manganese(IV) Oxide	731
15.5.1.2.1.3	Variation 3:	Using Chloranil	732
15.5.1.2.1.4	Variation 4:	Using Diphenylselenium Bis(trifluoroacetate)	733
15.5.1.2.1.5	Variation 5:	Using Mercury(II) Acetate	733
15.5.1.2.1.6	Variation 6:	Using Iodine	733
15.5.1.2.1.7	Variation 7:	Using a Noble-Metal Catalyst	734
15.5.1.2.1.8	Variation 8:	From an <i>N</i> -Sulfonyl-1,2,3,4-tetrahydroisoquinoline	735
15.5.1.2.2	Method 2:	Dehydrogenation of 3,4-Dihydroisoquinolines	735
15.5.1.2.2.1	Variation 1:	Using Sulfur	735
15.5.1.2.2.2	Variation 2:	Using Manganese(IV) Oxide	736
15.5.1.2.2.3	Variation 3:	Using <i>N</i> -Bromosuccinimide	736
15.5.1.2.2.4	Variation 4:	Using Diphenyl Disulfide	737
15.5.1.2.2.5	Variation 5:	Using Palladium	737
15.5.1.2.2.6	Variation 6:	Special Situations	738
15.5.1.2.3	Method 3:	Dehydrogenation of 1,2-Dihydroisoquinolines	739
15.5.1.2.4	Method 4:	Synthesis from 2-Benzopyrylium Salts	739
15.5.1.2.5	Method 5:	Synthesis from 1 <i>H</i> -Indene-1,2(3 <i>H</i>)-dione Monooximes	740
15.5.1.2.6	Method 6:	From an Isoquinoline-1,3(2 <i>H</i> ,4 <i>H</i>)-dione	740
15.5.1.2.7	Method 7:	From 1,4-Benzothiazepines	741
15.5.1.2.8	Method 8:	From Furo[3,4- <i>c</i>]pyridines	741
15.5.1.2.9	Method 9:	From [1,2,3]Triazolo[5,1- <i>a</i>]isoquinolines	743
15.5.1.2.10	Method 10:	From Zirconacyclopentadienes	743

15.5.1.3	Synthesis by Substituent Modification	744
15.5.1.3.1	Substitution of Existing Substituents	744
15.5.1.3.1.1	Of Hydrogen	744
15.5.1.3.1.1.1	Method 1: Protonation/Deuteration	744
15.5.1.3.1.1.2	Method 2: Metalation	745
15.5.1.3.1.1.3	Method 3: Acylation	745
15.5.1.3.1.1.3.1	Variation 1: Reaction with Acyl Radicals	745
15.5.1.3.1.1.3.2	Variation 2: Electrophilic Acylation	746
15.5.1.3.1.1.4	Method 4: Alkylation and Arylation	746
15.5.1.3.1.1.4.1	Variation 1: Reaction with Radicals	746
15.5.1.3.1.1.4.2	Variation 2: Electrophilic Alkylation	747
15.5.1.3.1.1.4.3	Variation 3: Nucleophilic Alkylation/Arylation	748
15.5.1.3.1.1.4.4	Variation 4: By Rearrangement	752
15.5.1.3.1.1.5	Method 5: Halogenation	752
15.5.1.3.1.1.6	Method 6: Sulfonylation and Seleninylation	754
15.5.1.3.1.1.7	Method 7: Nitration and Nitrosation	755
15.5.1.3.1.1.7.1	Variation 1: Electrophilic Substitution	755
15.5.1.3.1.1.7.2	Variation 2: Nucleophilic Substitution	757
15.5.1.3.1.1.8	Method 8: Diazo Coupling	758
15.5.1.3.1.1.9	Method 9: Amination	758
15.5.1.3.1.1.10	Method 10: Hydroxylation	759
15.5.1.3.1.1.11	Method 11: Oxidation	760
15.5.1.3.1.1.12	Method 12: Using Reissert Compounds	764
15.5.1.3.1.1.12.1	Variation 1: Synthesis of 1-Alkylisoquinolines	766
15.5.1.3.1.1.12.2	Variation 2: Synthesis of 1-(Hydroxyalkyl)isoquinolines	769
15.5.1.3.1.1.12.3	Variation 3: Synthesis of Isoquinoline-1-carbonitriles	770
15.5.1.3.1.1.12.4	Variation 4: Synthesis of 1-Hetarylisoquinolines	771
15.5.1.3.1.1.12.5	Variation 5: Synthesis of 1-Benzoylisoquinoline	771
15.5.1.3.1.1.12.6	Variation 6: Synthesis of 1-Pyrrol-2-ylisoquinolines	772
15.5.1.3.1.1.12.7	Variation 7: Synthesis of Phenyl 1-Allyl-1,2-dihydroisoquinoline-2-carboxylates	772
15.5.1.3.1.1.13	Method 13: Using <i>N</i> -Oxides	773
15.5.1.3.1.1.13.1	Variation 1: Synthesis of Isoquinolines	773
15.5.1.3.1.1.13.2	Variation 2: Synthesis of 1-Haloisoquinolines	773
15.5.1.3.1.1.13.3	Variation 3: Synthesis of 1-Acetoxyisoquinolines and Isoquinolin-1(2 <i>H</i>)-ones	774
15.5.1.3.1.1.13.4	Variation 4: Synthesis of Isoquinoline-1-carbonitriles	775
15.5.1.3.1.1.13.5	Variation 5: Introduction of Carbon Acids at C1	776
15.5.1.3.1.1.13.6	Variation 6: Reaction with Dichloroketene	778
15.5.1.3.1.1.13.7	Variation 7: Reaction with Diketene	779
15.5.1.3.1.1.13.8	Variation 8: Conversion into 1-Benzoylisoquinolines	779
15.5.1.3.1.1.13.9	Variation 9: Introduction of 1-Nitrogen Substituents	780
15.5.1.3.1.1.13.10	Variation 10: Using 2-(Acylimino)isoquinolinium Salts	782
15.5.1.3.1.1.13.11	Variation 11: Using 2-(2,6-Dimethyl-4-oxo-1,4-dihydro-1-pyridyl)isoquinolinium Salts	782
15.5.1.3.1.1.13.12	Variation 12: Using Isoquinolinium Salts and Ethyl (Trimethylsilyl)acetate or (Trimethylsilyl)acetonitrile	783

15.5.1.3.1.2	Of Metals	783
15.5.1.3.1.2.1	Method 1: Substitution of Metalated Isoquinolines	783
15.5.1.3.1.2.2	Method 2: Metal-Catalyzed Cross Coupling	785
15.5.1.3.1.3	Of Carbon Functionalities	786
15.5.1.3.1.4	Of Halogens	788
15.5.1.3.1.4.1	Method 1: Nucleophilic Substitutions	788
15.5.1.3.1.4.2	Method 2: Hydrogenolysis of C-Halo Groups	794
15.5.1.3.1.4.3	Method 3: Exchange of C-Halo Groups	795
15.5.1.3.1.4.4	Method 4: Metal-Catalyzed Cross Coupling	795
15.5.1.3.1.5	Of Oxygen	800
15.5.1.3.1.5.1	Method 1: Modification of Isoquinolin-1(2 <i>H</i>)-ones	800
15.5.1.3.1.5.2	Method 2: Of Hydroxy or Alkoxy Groups	802
15.5.1.3.1.5.3	Method 3: Of 4-Toluenesulfonates	803
15.5.1.3.1.6	Of Sulfur	803
15.5.1.3.1.6.1	Method 1: Of Sulfonic Acids	803
15.5.1.3.1.7	Of Nitrogen	804
15.5.1.3.2	Modification of Substituents	805
15.5.1.3.2.1	Of C-Carbon Substituents	805
15.5.1.3.2.1.1	Method 1: Carboxy and Related Functional Groups	805
15.5.1.3.2.1.2	Method 2: Aldehyde, Ketone, and Related Groups	806
15.5.1.3.2.1.3	Method 3: Alkyl and Hydroxyalkyl Groups	808
15.5.1.3.2.2	Of C-Oxygen Substituents	813
15.5.1.3.2.3	Of C-Nitrogen Substituents	814
15.5.2	Product Subclass 2: Isoquinoline 2-Oxides	815
15.5.2.1	Synthesis by Ring-Closure Reactions	815
15.5.2.1.1	By Annulation to an Arene	815
15.5.2.1.1.1	By Formation of One N—C Bond	815
15.5.2.1.1.1.1	With Formation of the 2—3 Bond	815
15.5.2.1.1.1.1.1	Method 1: From 2-Alk-1-ynylbenzaldehyde Oximes	815
15.5.2.2	Synthesis by Ring Transformation	816
15.5.2.3	Synthesis by Substituent Modification	817
15.5.2.3.1	Substitution of Existing Substituents	817
15.5.2.3.1.1	Of Hydrogen	817
15.5.2.3.1.1.1	Method 1: Nitration	817
15.5.2.3.1.2	Of C-Halogen Substituents	817
15.5.2.3.1.3	Of N-Oxygen Substituents	817
15.5.3	Product Subclass 3: Isoquinolinium Salts	817
15.5.3.1	Synthesis by Ring-Closure Reactions	818
15.5.3.1.1	By Annulation to an Arene	818

15.5.3.1.1.1	By Formation of Two N—C Bonds	818
15.5.3.1.1.1.1	With Formation of the 1—2 and 2—3 Bonds	818
15.5.3.1.1.1.1.1	Method 1: From 1,5-Diketones	818
15.5.3.1.1.2	By Formation of One N—C and One C—C Bond	819
15.5.3.1.1.2.1	With Formation of the 2—3 and 4—4a Bonds	819
15.5.3.1.1.3	By Formation of One N—C Bond	820
15.5.3.1.1.3.1	With Formation of the 1—2 Bond	820
15.5.3.1.1.4	By Formation of One C—C Bond	820
15.5.3.1.1.4.1	With Formation of the 1—8a Bond	820
15.5.3.2	Synthesis by Ring Transformation	820
15.5.3.2.1	Method 1: From Isoquinolines	820
15.5.3.2.2	Method 2: From 2-Benzopyrylium Salts	823
15.5.3.2.3	Method 3: From 2-Alkyl-1,2-dihydro- and 2-Alkyl-1,2,3,4-tetrahydro- isoquinolines	824
15.5.3.2.4	Method 4: From 5 <i>H</i> -2,3-Benzodiazepines	824
15.5.3.2.5	Method 5: From 1a,6a-Dihydroindeno[1,2- <i>b</i>]azirin-6(1 <i>H</i>)-ones	825
15.5.3.3	Synthesis by Substituent Modification	825
15.5.3.3.1	Substitution of Existing Substituents	825
15.5.3.3.1.1	Of Hydrogen	825
15.5.3.3.1.1.1	Method 1: Oxidation	825
15.5.3.3.2	Modification of Substituents	825
15.5.3.3.2.1	Of C-Carbon Substituents	825
15.5.3.3.2.2	Of C-Oxygen Substituents	826
15.5.3.3.2.3	Of N-Carbon Substituents	826
15.5.3.3.2.4	Of N-Nitrogen Substituents	827
15.6	Product Class 6: Isoquinolinones M. Álvarez and J. A. Joule	
15.6	Product Class 6: Isoquinolinones	839
15.6.1	Product Subclass 1: Isoquinolin-1(2<i>H</i>)-ones	841
15.6.1.1	Synthesis by Ring-Closure Reactions	841
15.6.1.1.1	By Annulation to an Arene	841
15.6.1.1.1.1	By Formation of One N—C and One C—C Bond	841
15.6.1.1.1.1.1	With Formation of the 1—2 and 3—4 Bonds	841
15.6.1.1.1.1.1.1	Method 1: From 2-Acylbenzoates	841
15.6.1.1.1.1.1.2	Method 2: From <i>N,N</i> -Diethyl-2-methylbenzamides and Imines	842
15.6.1.1.1.1.1.3	Method 3: Self-Condensation of Potassium 2-(Cyanomethyl)benzoate ..	843
15.6.1.1.1.1.2	With Formation of the 2—3 and 3—4 Bonds	844
15.6.1.1.1.1.2.1	Method 1: From (2-Alkylaryl)carboxamides	844

15.6.1.1.1.2.1.1	Variation 1:	From 2-Methylbenzamides	844
15.6.1.1.1.2.1.2	Variation 2:	From <i>O</i> -Methyl-2-alkylbenzohydroxamates	845
15.6.1.1.1.2.2	Method 2:	From 2-Methylisophthalonitrile	845
15.6.1.1.1.2.3	Method 3:	From 2-Alkynylbenzonitriles	846
15.6.1.1.1.2.4	Method 4:	From 2-Methylbenzoic Acid Hydrazides	846
15.6.1.1.1.3	With Formation of the 2–3 and 4–4a Bonds		847
15.6.1.1.1.3.1	Method 1:	From <i>N</i> -Methylbenzamide	847
15.6.1.1.1.4	By Formation of One N–C Bond		848
15.6.1.1.1.4.1	With Formation of the 1–2 Bond		848
15.6.1.1.1.4.1.1	Method 1:	From 2-(Cyanomethyl)benzoates	848
15.6.1.1.1.4.1.2	Method 2:	From Methyl 2-(Oxazol-5-yl)benzoates	848
15.6.1.1.1.4.1.3	Method 3:	From 2-[2-(Hydroxyimino)ethyl]benzoic Acid	849
15.6.1.1.1.4.1.4	Method 4:	From Chiral 2-(2-Aminoethyl)benzamides	849
15.6.1.1.1.4.1.5	Method 5:	From Alkyl 2-(2-Aminoethyl)benzoates	850
15.6.1.1.1.4.2	With Formation of the 2–3 Bond		850
15.6.1.1.1.4.2.1	Method 1:	From 2-(1-Cyano-2-phenylvinyl)benzonitrile	850
15.6.1.1.1.4.2.2	Method 2:	From Azidocinnamates with Phosphorus(III) Reagents: Intramolecular Aza-Wittig Reactions	851
15.6.1.1.1.4.2.3	Method 3:	From 2-[2-(Dimethylamino)vinyl]isophthalonitrile	852
15.6.1.1.1.4.2.4	Method 4:	From 2-(2-Ethoxyvinyl)benzonitrile	852
15.6.1.1.1.4.2.5	Method 5:	From Benzamides	853
15.6.1.1.1.4.2.5.1	Variation 1:	From 2-Halobenzamides	853
15.6.1.1.1.4.2.5.2	Variation 2:	From 2-Cyanobenzyl Ketones	853
15.6.1.1.1.4.2.6	Method 6:	From 2-Bromobenzamide	854
15.6.1.1.1.4.2.7	Method 7:	From 2-Chloro-5-nitrobenzonitrile	854
15.6.1.1.1.4.2.8	Method 8:	From 2-(2-Phenylvinyl)benzamide and <i>N</i> -(Phenylselanyl)succinimide	855
15.6.1.1.1.4.2.9	Method 9:	From 2-(2-Hydroxyethyl)benzamides	855
15.6.1.1.1.5	By Formation of One C–C Bond		856
15.6.1.1.1.5.1	With Formation of the 1–8a Bond		856
15.6.1.1.1.5.1.1	Method 1:	From the Acyl Azide of a Cinnamic Acid	856
15.6.1.1.1.5.1.2	Method 2:	From 2-Phenylvinyl Isothiocyanates	858
15.6.1.1.1.5.1.3	Method 3:	From 2-Arylethanamine Carbamates	858
15.6.1.1.1.5.2	With Formation of the 3–4 Bond		858
15.6.1.1.1.5.2.1	Method 1:	From Phosphorylated 2-Aroylbenzamides	858
15.6.1.1.1.5.2.2	Method 2:	From <i>N</i> -Acylbenzamides	859
15.6.1.1.1.5.2.3	Method 3:	From <i>N</i> -Alkyl- <i>N</i> -(silylmethyl)benzamides	860
15.6.1.1.1.5.3	With Formation of the 4–4a Bond		860
15.6.1.1.1.5.3.1	Method 1:	From Allenes	860
15.6.1.1.1.5.3.2	Method 2:	From <i>N</i> -(2,2-Dimethoxyethyl)benzamides	861
15.6.1.1.2	By Annulation to a Pyridine		861
15.6.1.1.2.1	By Formation of Two C–C Bonds		861
15.6.1.1.2.1.1	With Formation of the 4a–5 and 8–8a Bonds		861

15.6.1.1.2.1.1.1	Method 1:	From 2-Oxo-1,2-dihydropyridine-4-carbonitriles	861
15.6.1.1.2.1.1.1.1	Variation 1:	With 2,3-Dimethylbuta-1,3-diene	861
15.6.1.1.2.1.1.1.2	Variation 2:	With 2-/3-Methoxybuta-1,3-dienes	862
15.6.1.2	Synthesis by Ring Transformation		862
15.6.1.2.1	Method 1:	From Isoquinolines	862
15.6.1.2.2	Method 2:	From Isoquinoline 2-Oxides	863
15.6.1.2.3	Method 3:	From Isoquinolinium Salts	864
15.6.1.2.4	Method 4:	From Isoquinolin-1-amines	865
15.6.1.2.5	Method 5:	From Partially Reduced Isoquinolin-1(2 <i>H</i>)-ones	865
15.6.1.2.6	Method 6:	From Homophthalic Anhydrides and Homophthalimides	866
15.6.1.2.6.1	Variation 1:	Conversion of Homophthalic Anhydrides into Imides Followed by Partial Reduction	866
15.6.1.2.6.2	Variation 2:	Condensation of Homophthalic Anhydride with Trimethyl Orthoformate Followed by Reaction with Ammonia or Amines	867
15.6.1.2.6.3	Variation 3:	From Arylidenehomophthalimides	867
15.6.1.2.6.4	Variation 4:	From Homophthalic Anhydride and Aromatic Aldehydes	868
15.6.1.2.7	Method 7:	From 1 <i>H</i> -2-Benzopyran-1-ones	869
15.6.1.2.8	Method 8:	From Isobenzofuran-1(3 <i>H</i>)-ones	869
15.6.1.2.8.1	Variation 1:	By Lithiation of Isobenzofuran-1(3 <i>H</i>)-ones	869
15.6.1.2.8.2	Variation 2:	From 3-(Hydroxymethyl)isobenzofuran-1(3 <i>H</i>)-ones	870
15.6.1.2.9	Method 9:	From 1-Haloisoquinolines	871
15.6.1.2.10	Method 10:	From 2-(Carboxymethyl)benzoic Acids	872
15.6.1.2.11	Method 11:	Synthesis from Indanones	872
15.6.1.2.12	Method 12:	From Isoindolones	873
15.6.1.2.13	Method 13:	Synthesis from 1-(Allyloxy)isoquinoline	873
15.6.1.2.14	Method 14:	Synthesis from a Naphthalene	874
15.6.1.2.15	Method 15:	From Benzopyrancarbonitriles	874
15.6.1.3	Synthesis by Substituent Modification		875
15.6.1.3.1	Substitution of Existing Substituents		875
15.6.1.3.1.1	Of Hydrogen		875
15.6.1.3.1.1.1	Method 1:	Protonation/Deuteration	875
15.6.1.3.1.1.2	Method 2:	Acylation	875
15.6.1.3.1.1.3	Method 3:	Alkylation at Carbon	876
15.6.1.3.1.1.3.1	Variation 1:	Electrophilic Alkylation	876
15.6.1.3.1.1.3.2	Variation 2:	Photochemical Alkylation	878
15.6.1.3.1.1.4	Method 4:	Alkylation at Nitrogen	879
15.6.1.3.1.1.4.1	Variation 1:	Via Anions	879
15.6.1.3.1.1.4.2	Variation 2:	Via 1-(Trimethylsiloxy)isoquinolines	880
15.6.1.3.1.1.5	Method 5:	Halogenation	880
15.6.1.3.1.1.6	Method 6:	Nitration	881
15.6.1.3.1.1.7	Method 7:	Oxidation	882
15.6.1.3.1.2	Of Metals		883
15.6.1.3.1.3	Of Carbon Functionalities		883
15.6.1.3.1.4	Of Halogens		884
15.6.1.3.1.4.1	Method 1:	Using Copper Salts	884

15.6.1.3.1.4.2	Method 2: Metal-Catalyzed Cross Coupling	884
15.6.1.3.1.5	Of Oxygen	885
15.6.1.3.1.5.1	Method 1: Reduction of 2-Hydroxyisoquinolin-1(2 <i>H</i>)-ones	885
15.6.1.3.1.5.2	Method 2: Conversion into Isoquinolin-1(2 <i>H</i>)-thiones	885
15.6.1.3.2	Modification of Substituents	886
15.6.1.3.2.1	Of C-Carbon Substituents	886
15.6.1.3.2.1.1	Method 1: Modification of Carboxyl and Related Functional Groups	886
15.6.1.3.2.1.2	Method 2: Modification of Carbonyl and Related Groups	886
15.6.1.3.2.1.3	Method 3: Modification of Alkyl and Hydroxyalkyl Groups	886
15.6.1.3.2.2	Of C-Oxygen Substituents	887
15.6.1.3.2.2.1	Method 1: Modification of a 4-Hydroxy Substituent	887
15.6.1.3.2.3	Of C-Nitrogen Substituents	887
15.6.1.3.2.3.1	Method 1: Modification of Nitro and Amino Substituents	887
15.6.2	Product Subclass 2: Isoquinolin-3-ones and Isoquinolin-3-ols	888
15.6.2.1	Synthesis by Ring-Closure Reactions	888
15.6.2.1.1	By Annulation to an Arene	888
15.6.2.1.1.1	By Formation of Two N—C Bonds	888
15.6.2.1.1.1.1	With Formation of the 1—2 and 2—3 Bonds	888
15.6.2.1.1.1.1.1	Method 1: From 1,5-Dicarbonyl Compounds	888
15.6.2.1.1.1.1.1.1	Variation 1: From Alkyl (2-Acylphenyl)acetates	888
15.6.2.1.1.1.1.1.2	Variation 2: From 2-(2-Formylphenyl)- <i>N,N</i> -dimethylacetamides	889
15.6.2.1.1.1.1.1.3	Variation 3: From Homophthalic Acids	889
15.6.2.1.1.2	By Formation of One N—C and One C—C Bond	890
15.6.2.1.1.2.1	With Formation of the 1—8a and 2—3 Bonds	890
15.6.2.1.1.2.1.1	Method 1: From Phenylacetyl Chlorides and Imidoyl Chlorides	890
15.6.2.1.1.2.2	With Formation of the 1—8a and 1—2 Bonds	890
15.6.2.1.1.2.2.1	Method 1: From Phenylacetamides and Hexamethylenetetramine	890
15.6.2.1.1.3	By Formation of Two C—C Bonds	891
15.6.2.1.1.3.1	With Formation of the 4—4a and 1—8a Bonds	891
15.6.2.1.1.3.1.1	Method 1: From Benzo-1,4-quinones	891
15.6.2.1.1.4	By Formation of One N—C Bond	892
15.6.2.1.1.4.1	With Formation of the 2—3 Bond	892
15.6.2.1.1.4.1.1	Method 1: From (2-Cyanoaryl)acetyl Chlorides	892
15.6.2.1.1.4.1.2	Method 2: From Methyl (2-Cyanoaryl)acetates	893
15.6.2.1.1.5	By Formation of One C—C Bond	893
15.6.2.1.1.5.1	With Formation of the 4—4a Bond	893
15.6.2.1.1.5.1.1	Method 1: From <i>N</i> -Benzyl-2,2-diethoxyacetamides	893
15.6.2.1.1.5.1.2	Method 2: From <i>N</i> -Benzyl-2-hydroxy-2-phenylacetamide	894
15.6.2.1.1.5.1.3	Method 3: From <i>N</i> -Benzyl- <i>N</i> - <i>tert</i> -butyl-2-(2-sulfinylphenyl)acetamides	895

15.6.2.1.2	From Acyclic Precursors	895
15.6.2.1.2.1	By Formation of Two C—C bonds	895
15.6.2.1.2.1.1	With Formation of the 4a—8a and 7—8 Bonds	895
15.6.2.1.2.1.1.1	Method 1: From <i>N</i> -Prop-2-ynylhexadienylamides	895
15.6.2.2	Synthesis by Ring Transformation	896
15.6.2.2.1	Method 1: From Isoquinolin-3-amines	896
15.6.2.2.2	Method 2: From Benzo[<i>c</i>]pyrylium Salts	896
15.6.2.2.3	Method 3: By Dehydrogenation of 1,4-Dihydroisoquinolin-3(2 <i>H</i>)-ones	897
15.6.2.2.3.1	Variation 1: Without Incorporating Other Substituents	897
15.6.2.2.3.2	Variation 2: With Incorporation of a 4-Alkyl Group	898
15.6.2.2.3.3	Variation 3: With Incorporation of a 4-Halogen	899
15.6.2.2.3.4	Variation 4: With Incorporation of a 4-Oxygen	900
15.6.2.3	Synthesis by Substituent Modification	900
15.6.2.3.1	Substitution of Existing Substituents	900
15.6.2.3.1.1	Of Hydrogen	900
15.6.2.3.1.1.1	Method 1: Halogenation	900
15.6.2.3.1.1.2	Method 2: Nitration and Nitrosation	901
15.6.2.3.2	Modification of Substituents	901
15.6.2.3.2.1	Of C-Oxygen Substituents	901
15.6.2.3.2.2	Of C-Nitrogen Substituents	902
15.7	Product Class 7: Quinolizinium Salts and Benzo Analogues H. Ihmels	
15.7	Product Class 7: Quinolizinium Salts and Benzo Analogues	907
15.7.1	Product Subclass 1: Quinolizinium Salts	909
15.7.1.1	Synthesis by Ring-Closure Reactions	909
15.7.1.1.1	By Annulation to an Arene	909
15.7.1.1.1.1	By Formation of One N—C and One C—C Bond	909
15.7.1.1.1.1.1	Method 1: From 2-Pyridylacetonitrile and β -Dicarbonyl Derivatives	909
15.7.1.1.1.1.2	Method 2: From Ethyl 2-Pyridylacetate and 3-Ethoxyprop-2-en-1-one Derivatives	909
15.7.1.1.1.2	By Formation of Two C—C Bonds	910
15.7.1.1.1.2.1	Method 1: From 1,2-Dialkylpyridinium Salts and 1,2-Diketones	910
15.7.1.1.1.3	By Formation of One N—C Bond	911
15.7.1.1.1.3.1	Method 1: Cyclization of Hydroxy-Substituted 1-Oxo-4-(2-pyridyl)butane Derivatives	911
15.7.1.1.1.3.1.1	Variation 1: Cyclization of 4-Alkoxy-1-(2-pyridyl)pent-3-en-2-ol Derivatives	911
15.7.1.1.1.3.1.2	Variation 2: Cyclization of 4,4-Dialkoxy-1-(2-pyridyl)butan-2-ol Derivatives	912
15.7.1.1.1.3.1.3	Variation 3: From 8-[(Dimethylamino)methylene]-5 <i>H</i> -pyrano[4,3- <i>b</i>]pyridine-5,7(8 <i>H</i>)-dione	913

15.7.1.1.1.3.2	Method 2: Cyclization of 4,4-Bis(methylsulfanyl)-1-(2-pyridyl)but-3-en-2-ol Derivatives	914
15.7.1.1.1.4	By Formation of One C—C Bond	915
15.7.1.1.1.4.1	Method 1: Cyclization of 2-(1,3-Dioxolan-2-yl)-1-(2-oxopropyl)pyridinium Derivatives	915
15.7.1.1.1.4.2	Method 2: Cyclization of 2-Acetyl-1-(2-oxoethyl)pyridinium Derivatives ..	915
15.7.1.2	Aromatization	916
15.7.1.2.1	Method 1: Of 1-Oxo-1,2,3,4-tetrahydroquinolizinium Derivatives	916
15.7.1.2.2	Method 2: Of 2 <i>H</i> -Quinolizin-2-one	917
15.7.1.2.3	Method 3: Of Dihydroquinolizine and Tetrahydroquinolizinium Derivatives	917
15.7.1.3	Synthesis by Substituent Modification	919
15.7.1.3.1	Substitution of Existing Substituents	919
15.7.1.3.1.1	Of Hydrogen	919
15.7.1.3.1.1.1	Method 1: Introduction of Bromine	919
15.7.1.3.1.1.2	Method 2: Introduction of Nitrogen Substituents	919
15.7.1.3.1.2	Of Carbon Functionalities	920
15.7.1.3.1.2.1	Method 1: Cleavage of C—C Bonds	920
15.7.1.3.1.3	Of Heteroatoms	920
15.7.1.3.1.3.1	Method 1: Stille Coupling of Bromoquinolizinium Derivatives	920
15.7.1.3.1.3.2	Method 2: Replacement of a Diazonium Group by Bromine	921
15.7.1.3.1.3.3	Method 3: Replacement of Halogens	922
15.7.1.3.2	Modification of Substituents	922
15.7.1.3.2.1	Method 1: Oxidation of Methyl Groups	922
15.7.1.3.2.2	Method 2: Oxidation of Benzo[<i>b</i>]quinolizinium Derivatives	923
15.7.1.3.2.3	Method 3: Reduction of Methoxycarbonyl Groups	923
15.7.1.3.2.4	Method 4: Knoevenagel-Type Condensation of Methylquinolizinium Derivatives	923
15.7.2	Product Subclass 2: Benzo[<i>b</i>]quinolizinium Salts	924
15.7.2.1	Synthesis by Ring-Closure Reactions	924
15.7.2.1.1	By Annulation to an Arene	924
15.7.2.1.1.1	By Formation of One N—C and One C—C Bond	924
15.7.2.1.1.1.1	Method 1: From 2-Benzylpyridine Derivatives and Acylating Reagents ...	924
15.7.2.1.1.1.2	Method 2: From 2-Benzylpyridine Derivatives and Alkynes	924
15.7.2.1.1.2	By Formation of One N—C Bond	925
15.7.2.1.1.2.1	Method 1: Cyclization of 2-(2-Pyridylmethyl)benzaldehyde Oxime Derivatives	925
15.7.2.1.1.2.2	Method 2: Cyclization of 2-(2-Pyridylmethyl)benzotrile Derivatives ...	926
15.7.2.1.1.2.3	Method 3: Cyclization of [2-(Chloromethyl)phenyl](2-pyridyl)methanol Derivatives	927
15.7.2.1.1.3	By Formation of One C—C Bond	928

15.7.2.1.1.3.1	Method 1:	Cyclization of 1-Benzyl-2-formylpyridinium Derivatives	928
15.7.2.2	Aromatization		930
15.7.2.2.1	Method 1:	Of 7,8-Dioxo-7,8-dihydrobenzo[<i>b</i>]quinolizinium Derivatives	930
15.7.2.2.2	Method 2:	Of 1-Oxo-1,2,3,4-tetrahydrobenzo[<i>b</i>]quinolizinium Derivatives	930
15.7.2.2.3	Method 3:	Of 6 <i>H</i> -Benzo[<i>b</i>]quinolizine-6-carbonitrile Derivatives	931
15.7.2.2.4	Method 4:	Of 7,8,9,10-Tetrabromo-7,8,9,10-tetrahydrobenzo[<i>b</i>]quinolizinium Derivatives	932
15.7.2.3	Synthesis by Substituent Modification		932
15.7.2.3.1	Substitution of Existing Substituents		932
15.7.2.3.1.1	Of Hydrogen		932
15.7.2.3.1.1.1	Method 1:	Introduction of Bromine	932
15.7.2.3.1.1.2	Method 2:	Introduction of Nitrogen Substituents	933
15.7.2.3.1.1.3	Method 3:	Introduction of Sulfur Substituents	934
15.7.3	Product Subclass 3: Benzo[<i>c</i>]quinolizinium Salts		934
15.7.3.1	Synthesis by Ring-Closure Reactions		934
15.7.3.1.1	By Annulation to an Arene		934
15.7.3.1.1.1	By Formation of One N—C and One C—C Bond		934
15.7.3.1.1.1.1	Method 1:	Condensation between Pyrylium Salts and 2-Aminobenzaldehyde	934
15.7.3.1.1.2	By Formation of One N—C Bond		935
15.7.3.1.1.2.1	Method 1:	Cyclization of 2-[2-(2-Chlorophenyl)vinyl]pyridine Derivatives	935
15.7.3.1.1.2.2	Method 2:	Cyclization of 1-(2-Chlorophenyl)-2-(2-pyridyl)ethanone Derivatives	936
15.7.3.2	Aromatization		936
15.7.3.2.1	Method 1:	Of 4-Oxo-1,2,3,4-tetrahydropyridobenzo[<i>c</i>]quinolizinium Derivatives	936
15.7.3.3	Synthesis by Substituent Modification		937
15.7.3.3.1	Substitution of Existing Substituents		937
15.7.3.3.1.1	Of Hydrogen		937
15.7.3.3.1.1.1	Method 1:	Introduction of Nitrogen Substituents	937
15.7.3.3.1.1.2	Method 2:	Introduction of Sulfur Substituents	938
15.7.3.3.2	Modification of Substituents		938
15.7.3.3.2.1	Method 1:	Knoevenagel-Type Condensation of Methylbenzo[<i>c</i>]quinolizinium Derivatives	938
15.7.4	Product Subclass 4: Benzo[<i>a</i>]quinolizinium Salts		938
15.7.4.1	Synthesis by Ring-Closure Reactions		938
15.7.4.1.1	By Annulation to an Arene		938
15.7.4.1.1.1	By Formation of One N—C and One C—C Bond		938

15.7.4.1.1.1	Method 1: From N-Alkylated 1-Alkylisoquinolinium Salts and 1,2-Diketones	938
15.7.4.1.1.2	By Formation of Two C—C Bonds	939
15.7.4.1.1.2.1	Method 1: From Isoquinolinium Salts and Mesityl Oxide	939
15.7.4.1.1.3	By Formation of One C—C Bond	939
15.7.4.1.1.3.1	Method 1: Cyclization of 1-(2-Oxoalkyl)-2-phenylpyridinium Derivatives ·	939
15.7.4.1.1.3.2	Method 2: Cyclization of 1-Styrylpyridinium Derivatives	940
15.7.4.2	Synthesis by Ring Transformation	941
15.7.4.2.1	Method 1: Ring Contraction of Pyrido[2,1- <i>b</i>][1,3]benzothiazepinium Derivatives	941
15.7.4.3	Aromatization	941
15.7.4.3.1	Method 1: Of 1-Oxo-1,2,3,4-tetrahydrobenzo[<i>a</i>]quinolizinium Derivatives	941
15.7.4.4	Synthesis by Substituent Modification	942
15.7.4.4.1	Substitution of Existing Substituents	942
15.7.4.4.1.1	Of Hydrogen	942
15.7.4.4.1.1.1	Method 1: Introduction of Nitrogen Substituents	942
15.8	Product Class 8: Naphthyridines	
	P.-W. Phuan and M. C. Kozłowski	
15.8	Product Class 8: Naphthyridines	947
15.8.1	Synthesis by Ring-Closure Reactions	949
15.8.1.1	By the Direct Formation of Naphthyridine Rings	949
15.8.1.1.1	By Formation of Four N—C Bonds	949
15.8.1.1.1.1	Method 1: Friedel–Crafts Tetraacylation of 2-Methylpropene	949
15.8.1.1.2	By Formation of Two N—C Bonds	951
15.8.1.1.2.1	Method 1: Multiple Iminoalkylation of Alkenes with the Vilsmeier Reagent, Followed by Cyclization	951
15.8.1.1.2.2	Method 2: Conversion of a Pyranilideneiminium Salt	952
15.8.1.2	By Annulation to the Heterocyclic Ring	952
15.8.1.2.1	By Formation of Two N—C Bonds	952
15.8.1.2.1.1	Method 1: The Sonogashira Reaction	952
15.8.1.2.1.1.1	Variation 1: Electrophilic Ring Closure of Iminoalkynes	955
15.8.1.2.2	By Formation of One N—C and One C—C Bond	956
15.8.1.2.2.1	Method 1: The Skraup Reaction	956
15.8.1.2.2.2	Method 2: Ethoxymethylenemalonic Ester Condensation	959
15.8.1.2.2.3	Method 3: The Heck Reaction	964
15.8.1.2.2.4	Method 4: The Suzuki and Stille Reactions	966
15.8.1.2.2.4.1	Variation 1: Synthesis of Thieno-Fused Naphthyridines via a Suzuki Coupling	966

15.8.1.2.2.4.2	Variation 2:	Synthesis of Benzo-Fused Naphthyridines via a Suzuki Coupling	967
15.8.1.2.2.4.3	Variation 3:	Synthesis of Thieno-Fused Naphthyridines via a Stille Coupling	968
15.8.1.2.2.4.4	Variation 4:	Synthesis of Benzo-Fused Naphthyridines via a Stille Coupling	970
15.8.1.2.2.5	Method 5:	The Friedländer Reaction	970
15.8.1.2.2.5.1	Variation 1:	A Highly Regioselective Reaction Using β -Oxo Phosphonates	974
15.8.1.2.2.6	Method 6:	The Niementowski Condensation	975
15.8.1.2.3	By Formation of One N—C Bond		975
15.8.1.2.3.1	By Homologation of <i>o</i> -Methylpyridinecarbonitriles, Followed by Cyclization ..		975
15.8.1.2.3.1.1	Method 1:	Formation of Enamines, Followed by Cyclization	976
15.8.1.2.3.1.2	Method 2:	Condensation of <i>o</i> -Methylpyridinecarbonitriles with Dialkyl Oxalates, Followed by Cyclization	977
15.8.1.2.3.2	By the Cyclization of Dinitriles		978
15.8.1.2.3.2.1	Method 1:	Ring Closure with Hydrogen Bromide	979
15.8.1.2.3.2.2	Method 2:	Ring Closure with an Alkoxide	980
15.8.2	Synthesis by Ring Transformation		981
15.8.2.1	By Ring Enlargement		981
15.8.2.1.1	Method 1:	Ring Opening, Followed by Ring Condensation	981
15.8.2.1.2	Method 2:	Carbene Insertion	982
15.8.2.1.3	Method 3:	By the Ring Expansion of Cyclic Imides	982
15.8.2.2	By Formal Exchange of Ring Members with Retention of the Ring Size		983
15.8.2.2.1	Method 1:	Conversion of Pyrido[2,3- <i>d</i>]pyrimidine and Its 3-Oxide into 2,3-Disubstituted 1,8-Naphthyridines	983
15.9	Product Class 9: Acridines R. H. Prager and C. M. Williams		
15.9	Product Class 9: Acridines		987
15.9.1	Product Subclass 1: Acridines		987
15.9.1.1	Synthesis by Ring-Closure Reactions		988
15.9.1.1.1	By Formation of One N—C and One C—C Bond		988
15.9.1.1.1.1	Method 1:	Acridines from 2-Aminophenylcarbonyl Compounds and Phenols	988
15.9.1.1.1.2	Method 2:	Electrocyclic Reactions	994
15.9.1.1.1.2.1	Variation 1:	From Quinomethane Precursors	994
15.9.1.1.1.2.2	Variation 2:	From Benzynes and Imines	995
15.9.1.1.1.2.3	Variation 3:	From Nitroarenes	996
15.9.1.1.2	By Formation of Two C—C Bonds		997
15.9.1.1.2.1	Method 1:	Synthesis from Diphenylamines and Carboxylic Acids: The Bernthsen Reaction	997
15.9.1.1.2.2	Method 2:	Radical Reactions of Quinones	998
15.9.1.1.2.3	Method 3:	Aza-Diels–Alder Reactions	999

15.9.1.1.2.4	Method 4: Synthesis from Acylated Diphenylamines	1000
15.9.1.1.3	By Formation of One N—C Bond	1000
15.9.1.1.3.1	Method 1: Electrocyclic Reactions	1000
15.9.1.1.3.2	Method 2: Arylation of Phenylacetonitriles	1001
15.9.1.1.3.3	Method 3: Photolysis or Pyrolysis of Aryl Azides	1001
15.9.1.1.3.4	Method 4: Cyclization of Bis(aminoaryl)methanes	1002
15.9.1.1.4	By Formation of One C—C Bond	1002
15.9.1.1.4.1	Method 1: Electrocyclic Reactions	1002
15.9.1.1.4.2	Method 2: Synthesis from Diarylamines	1003
15.9.1.1.4.2.1	Variation 1: From 2-(Arylamino)benzaldehydes	1005
15.9.1.1.4.2.2	Variation 2: By the McFadyen–Stevens Reaction	1007
15.9.1.1.4.2.3	Variation 3: From (2-Arylamino)acetophenones	1008
15.9.1.1.4.2.4	Variation 4: From 2-Anilino benzoic Acids	1010
15.9.1.1.4.3	Method 3: Synthesis from Enamines and Imines	1011
15.9.1.2	Synthesis by Ring Transformation	1011
15.9.1.2.1	By Ring Enlargement	1011
15.9.1.2.2	By Ring Contraction	1012
15.9.1.3	Aromatization	1012
15.9.1.3.1	Method 1: Dehydrogenation of Dihydroacridines and Tetrahydroacridines	1012
15.9.1.4	Synthesis by Substituent Modification	1013
15.9.1.4.1	Substitution of Existing Substituents	1013
15.9.1.4.1.1	Of Hydrogen	1013
15.9.1.4.1.2	Of C9 Substituents	1015
15.9.1.4.2	Modification of Substituents	1019
15.9.1.4.2.1	Method 1: Reduction of Acridinones	1019
15.9.2	Product Subclass 2: Acridine 10-Oxides	1021
15.9.3	Product Subclass 3: Acridinium Salts	1022
15.10	Product Class 10: Acridin-9(10<i>H</i>)-ones and Related Systems R. H. Prager and C. M. Williams	
15.10	Product Class 10: Acridin-9(10<i>H</i>)-ones and Related Systems	1029
15.10.1	Product Subclass 1: Acridin-9(10<i>H</i>)-ones	1031
15.10.1.1	Synthesis by Ring-Closure Reactions	1031
15.10.1.1.1	By Formation of Two N—C Bonds	1031
15.10.1.1.1.1	Method 1: Reaction of 2,2'-Dinitrobenzophenones with Aromatic Amines	1031
15.10.1.1.2	By Formation of One N—C and One C—C Bond	1031
15.10.1.1.2.1	Method 1: Reaction of 2-Aminobenzoic Acids and Phenols	1031
15.10.1.1.2.2	Method 2: Reaction of 2-Aminobenzoic Acids and Benzenes	1032

15.10.1.1.2.3	Method 3: Reaction of Isoatoic Anhydrides and Cyclohexenones	1033
15.10.1.1.3	By Formation of Two C—C Bonds	1035
15.10.1.1.3.1	Method 1: Lithiation of Quinolinones	1035
15.10.1.1.3.2	Method 2: Synthesis from Quinolin-4-ones and 4-Alkoxyquinolinium Salts	1036
15.10.1.1.3.3	Method 3: Electrocyclic Reactions	1036
15.10.1.1.4	By Formation of One N—C Bond	1036
15.10.1.1.4.1	Method 1: Nucleophilic Substitution	1036
15.10.1.1.4.1.1	Variation 1: Intramolecular Substitution of Fluoroarenes	1036
15.10.1.1.4.1.2	Variation 2: Intramolecular Addition to Benzynes	1037
15.10.1.1.4.1.3	Variation 3: Chapman Rearrangement	1037
15.10.1.1.4.2	Method 2: Photochemical and Pyrolytic Methods	1038
15.10.1.1.5	By Formation of One C—C Bond	1040
15.10.1.1.5.1	Method 1: Synthesis from 2-Anilinobenzoic Acids: The Ullmann–Jourdan Procedure	1040
15.10.1.1.5.1.1	Variation 1: Reaction of 2-Halobenzoic Acids	1040
15.10.1.1.5.1.2	Variation 2: Reaction of 2-Aminobenzoic Acids and Haloarenes	1043
15.10.1.1.5.1.3	Variation 3: Directed Metalation	1044
15.10.1.1.5.1.4	Variation 4: Reaction of 2-(Phenyliodonio)benzoates	1046
15.10.1.1.5.2	Method 2: Reaction of 2-Aminobenzamides and Ketones	1047
15.10.1.1.5.3	Method 3: Intramolecular Cycloaddition	1048
15.10.1.1.5.3.1	Variation 1: From 2-Styrylquinolin-4(1 <i>H</i>)-ones	1048
15.10.1.1.5.3.2	Variation 2: From 3-Styrylquinolin-4(1 <i>H</i>)-ones	1050
15.10.1.1.5.4	Method 4: Synthesis from Quinolin-4(1 <i>H</i>)-ones by Acylation at C3	1051
15.10.1.1.5.5	Method 5: Synthesis from Ketene Intermediates	1052
15.10.1.2	Aromatization	1054
15.10.1.2.1	Method 1: Dehydrogenation of Dihydroacridinones and Tetrahydroacridinones	1054
15.10.1.2.2	Method 2: Oxidation of Acridines and Acridinium Salts	1055
15.10.1.3	Synthesis by Substituent Modification	1056
15.10.2	Product Subclass 2: Acridine-9(10<i>H</i>)-thiones	1056
15.10.3	Product Subclass 3: Acridine-9(10<i>H</i>)-selones	1059
15.11	Product Class 11: Phenanthridines P. A. Keller	
15.11	Product Class 11: Phenanthridines	1065
15.11.1	Product Subclass 1: Phenanthridines	1065
15.11.1.1	Synthesis by Ring-Closure Reactions	1065
15.11.1.1.1	By Annulation to an Arene	1065
15.11.1.1.1.1	By Formation of One N—C and One C—C Bond	1066
15.11.1.1.1.1.1	Method 1: Resin-Supported Palladium-Based Cross-Coupling Reactions with Spontaneous Cyclization	1066

15.11.1.1.1.2	By Formation of Two C—C Bonds	1067
15.11.1.1.1.2.1	Method 1: Tandem [4+2] Cycloadditions and Cyclization	1067
15.11.1.1.1.3	By Formation of One N—C Bond	1067
15.11.1.1.1.3.1	Method 1: Generation of an Imine	1067
15.11.1.1.1.4	By Formation of One C—C Bond	1068
15.11.1.1.1.4.1	By Formation of the Biaryl Bond	1068
15.11.1.1.1.4.1.1	Method 1: Palladium-Based Couplings	1068
15.11.1.1.1.4.1.2	Method 2: The Use of Pschorr Ring-Closure Reactions	1069
15.11.1.1.1.4.1.3	Method 3: Photochemical Conditions	1070
15.11.1.1.1.4.1.3.1	Variation 1: Oxidative Photochemical Reactions	1071
15.11.1.1.1.4.1.4	Method 4: By Elimination–Addition	1072
15.11.1.1.1.4.1.5	Method 5: From 1,2,3-Benzotriazines with the Elimination of Nitrogen Gas	1073
15.11.1.1.1.4.2	By Ring Closure To Form the Heterocycle	1073
15.11.1.1.1.4.2.1	Method 1: Electrophilic Cyclization	1073
15.11.1.1.1.4.2.2	Method 2: Cyclization From Imines	1075
15.11.1.1.2	By Annulation to the Heterocyclic Ring	1075
15.11.1.1.2.1	Method 1: Using the Pschorr Cyclization	1075
15.11.1.1.2.2	Method 2: Using Oxidative Photochemical Reactions	1076
15.11.1.2	Synthesis by Ring Transformation	1077
15.11.1.2.1	By Ring Enlargement	1077
15.11.1.2.1.1	Method 1: From 9 <i>H</i> -Fluoren-9-ols	1077
15.11.1.2.2	By Ring Contraction	1077
15.11.1.2.2.1	Method 1: From Thiazepines	1077
15.11.1.2.2.2	Method 2: From 1,4-Diazocines	1078
15.11.1.2.2.3	Method 3: From Azepines	1079
15.11.1.3	Aromatization	1079
15.11.1.3.1	Method 1: Oxidation of 5,6-Dihydrophenanthridines	1079
15.11.1.4	Synthesis by Substituent Modification	1079
15.11.1.4.1	Of Heteroatoms	1079
15.11.1.4.1.1	Method 1: Reduction of Phenanthridine 5-Oxides	1079
15.11.1.4.2	Modification of Substituents	1080
15.11.1.4.2.1	Reduction of Phenanthridinones	1080
15.11.1.4.2.1.1	Method 1: By Reduction of the Corresponding Thioxo Compound	1080
15.11.1.4.2.1.1.1	Variation 1: By O-Substitution	1081
15.11.1.4.2.1.1.2	Variation 2: By Halogenation of Carbonyls	1082
15.11.2	Product Subclass 2: Phenanthridine 5-Oxides	1082
15.11.2.1	Synthesis by Ring-Closure Reactions	1082
15.11.2.1.1	By Annulation to Arene	1082
15.11.2.1.1.1	By Formation of One N—C Bond	1082

15.11.2.1.1.1	Method 1: Cyclization of Nitrobiphenyls	1082
15.11.2.2	Synthesis by Substituent Modification	1083
15.11.2.2.1	Addition to Hetarenes	1083
15.11.2.2.1.1	Method 1: Oxidation of Phenanthridines	1083
15.11.3	Product Subclass 3: Phenanthridinium Salts	1084
15.11.3.1	Synthesis by Ring-Closure Reactions	1084
15.11.3.1.1	By Annulation to an Arene	1084
15.11.3.1.1.1	By Formation of One N—C Bond	1084
15.11.3.1.1.1.1	Method 1: Phosphoryl Chloride Induced Cyclization	1084
15.11.3.2	Synthesis by Substituent Modification	1084
15.11.3.2.1	Addition of Organic Groups	1084
15.11.3.2.1.1	Method 1: Alkylation of Phenanthridines	1084
15.11.3.2.2	Modification of Substituents	1085
15.11.3.2.2.1	Method 1: De-ethoxylation of 5-Methylbenzo[c]phenanthridine	1085
15.12	Product Class 12: Phenanthridinones and Related Systems P. A. Keller	
15.12	Product Class 12: Phenanthridinones and Related Systems	1089
15.12.1	Product Subclass 1: Phenanthridinones	1089
15.12.1.1	Synthesis by Ring-Closure Reactions	1089
15.12.1.1.1	By Annulation to an Arene	1089
15.12.1.1.1.1	By Formation of Two C—C Bonds	1089
15.12.1.1.1.1.1	Method 1: Thermal Tandem Cyclization of an Allene-1,1-dicarboxanilide	1089
15.12.1.1.1.2	By Formation of One N—C Bond	1090
15.12.1.1.1.2.1	Method 1: Radical Cyclization of 2-(Formylamino)biaryls	1090
15.12.1.1.1.2.1.1	Variation 1: Photochemically Induced Cyclization	1090
15.12.1.1.1.3	By Formation of One C—C Bond	1091
15.12.1.1.1.3.1	By Formation of the Biaryl Bond	1091
15.12.1.1.1.3.1.1	Method 1: Palladium-Based Couplings	1091
15.12.1.2	Synthesis by Ring Transformation	1092
15.12.1.2.1	By Ring Enlargement	1092
15.12.1.2.1.1	Method 1: By Acid-Catalyzed Ring Expansion of Fluorenone Oximes	1092
15.12.1.3	Synthesis by Substituent Modification	1092
15.12.1.3.1	Method 1: Photochemical Rearrangement of Phenanthridine 5-Oxides	1092
15.12.1.3.1.1	Variation 1: Oxidation of Phenanthridinium Salts	1093
15.12.1.3.1.2	Variation 2: Oxidation of Phenanthridines	1094
15.12.2	Product Subclass 2: Phenanthridinethiones	1094
15.12.2.1	Synthesis by Ring-Closure Reactions	1095

15.12.2.1.1	By Annulation to an Arene	1095
15.12.2.1.1.1	Formation of One C—C Bond	1095
15.12.2.1.1.1.1	Method 1: Cyclization of Isothiocyanates	1095
15.13	Product Class 13: 1λ^3-Phosphinines F. Mathey and P. Le Floch	
15.13	Product Class 13: 1λ^3-Phosphinines	1097
15.13.1	Product Subclass 1: 1λ^3-Phosphinines	1100
15.13.1.1	Synthesis by Ring Transformation	1102
15.13.1.1.1	Ring Enlargement	1102
15.13.1.1.1.1	Method 1: From 1,2-Dihydrophosphetes	1102
15.13.1.1.1.2	Method 2: From Five-Membered Phosphorus Heterocycles	1103
15.13.1.1.2	Formal Exchange of Ring Members with Retention of the Ring Size	1107
15.13.1.1.2.1	Method 1: Tin–Phosphorus Exchange from 1,1-Dibutyl-1,4-dihydrostannines	1107
15.13.1.1.2.2	Method 2: Oxygen–Phosphorus Exchange from Pyrylium Salts	1108
15.13.1.1.2.3	Method 3: From Aza- and Diazaphosphinines	1110
15.13.1.2	Synthesis by Ring-Closure Reactions	1119
15.13.1.2.1	By Formation of One P—C and One C—C Bond	1119
15.13.1.2.1.1	Fragments P—C and C—C—C	1119
15.13.1.2.1.1.1	Method 1: From Phosphaalkynes	1119
15.13.1.2.1.1.2	Method 2: From Phosphaalkenes	1122
15.13.1.3	Aromatization	1124
15.13.1.3.1	Method 1: From 1 λ^5 -Phosphinines	1124
15.13.1.3.2	Method 2: From Dihydro-1 λ^3 -phosphinines	1125
15.13.1.3.3	Method 3: From Dihydrophosphinine Oxides and Sulfides	1126
15.13.1.4	Synthesis by Substituent Modification	1128
15.13.1.4.1	Method 1: Nucleophilic Aromatic Substitution	1128
15.13.1.4.2	Method 2: By Transformation of 2-Metalated Phosphinines	1128
15.13.1.4.2.1	Variation 1: Lithium and Magnesium Derivatives	1128
15.13.1.4.2.2	Variation 2: Zinc Derivatives	1129
15.13.1.4.2.3	Variation 3: Zirconium(IV) Derivatives	1130
15.13.1.4.2.4	Variation 4: Metal-Catalyzed Cross-Coupling Reactions with 2-Halophosphinines	1134
15.13.1.4.3	Method 3: Reactions at the P=C Bond	1136
15.13.1.4.3.1	Variation 1: Bromination of the Ring	1136
15.13.1.4.3.2	Variation 2: Phosphination and Ethylation of the Ring	1136
15.13.1.4.4	Methods 4: Miscellaneous Transformations	1138
15.13.2	Product Subclass 2: 1λ^3-Phosphinine Complexes	1140
15.13.2.1	σ -Complexes	1141
15.13.2.1.1	Method 1: By Ligand Displacement with 1 λ^3 -Phosphinines	1141

15.13.2.2	π -Complexes	1148
15.13.2.2.1	Method 1: By Ligand Displacement with $1\lambda^3$ -Phosphinines	1148
15.14	Product Class 14: $1\lambda^5$-Phosphinines R. Streubel	
15.14	Product Class 14: $1\lambda^5$-Phosphinines	1157
15.14.1	Product Subclass 1: $1\lambda^5$-Phosphinines	1162
15.14.1.1	Synthesis by Ring Transformation	1162
15.14.1.1.1	Method 1: Reaction of Primary Phosphinines with Pirylium Salts in the Presence of Nucleophiles	1162
15.14.1.1.2	Method 2: Oxidation of $1\lambda^3$ -Phosphinines to $1\lambda^5$ -Phosphinines	1163
15.14.1.1.2.1	Variation 1: Consecutive Reaction of $1\lambda^3$ -Phosphinines with Nucleophiles and Electrophiles	1163
15.14.1.1.2.2	Variation 2: Addition of Radicals to $1\lambda^3$ -Phosphinines	1165
15.14.1.1.2.3	Variation 3: Consecutive Reaction of $1\lambda^3$ -Phosphinines with Electrophiles and Nucleophiles	1165
15.14.1.1.2.4	Variation 4: Oxidative Addition of Protic Nucleophiles	1166
15.14.1.1.2.5	Variation 5: Reaction of $1\lambda^3$ -Phosphinine Dianions with Electrophiles	1167
15.14.1.1.3	Method 3: Thermal Ring Contraction of a 1-Phosphacyclonona- 2,4,8-triene	1167
15.14.1.2	Aromatization	1167
15.14.1.2.1	Method 1: Dehydrobromination of a 1,1-Diphenylphosphanium Bromide	1167
15.14.1.2.2	Method 2: Reaction of 1,2,3,4-Tetrahydro- and 1,2- and 1,4-Dihydro- $1\lambda^5$ -phosphinine 1-Oxides with Alkylating Agents in the Presence of Base	1168
15.14.1.2.3	Method 3: Reaction of 1,2,5,6-Tetrahydro- $1\lambda^5$ -phosphinine 1-Oxides with Phosphorus Pentachloride	1169
15.14.1.2.4	Method 4: Rearrangement of 1,2- and 1,4-Dihydro- $1\lambda^3$ -phosphinines ..	1169
15.14.1.2.4.1	Variation 1: Acid-Catalyzed Rearrangement	1169
15.14.1.2.4.2	Variation 2: Thermal Rearrangement	1170
15.14.1.3	Synthesis by Substituent Modification	1170
15.14.1.3.1	Substitution of Existing Substituents	1170
15.14.1.3.1.1	Method 1: C2 Acylation of 4-Aryl-Substituted $1\lambda^5$ -Phosphinines	1170
15.14.1.3.1.2	Method 2: Transformation of 4-Methyl- into 4-(Arylazo)- $1\lambda^5$ -phos- phinines	1171
15.14.1.3.1.3	Method 3: Substitution of <i>P</i> -Substituents of $1\lambda^5$ -Phosphinines	1171
15.14.1.3.1.3.1	Variation 1: Substitution of Halogen Substituents	1171
15.14.1.3.1.3.2	Variation 2: Substitution of Other Functional Groups	1174
15.14.1.3.2	Decomplexation of (2,3,4,5,6- η)- $1\lambda^5$ -Phosphinine Complexes	1174
15.14.2	Product Subclass 2: (2,3,4,5,6-η)-$1\lambda^5$-Phosphinine Complexes	1175
15.14.2.1	Synthesis by Complexation of $1\lambda^5$ -Phosphinines	1175

15.14.2.1.1	Method 1: Reaction of $1\lambda^5$ -Phosphinines with Metal(0)-Carbonyl Complexes	1175
15.14.2.2	Synthesis by Reaction of η^6 - $1\lambda^3$ -Phosphinine Complexes	1176
15.14.2.2.1	Method 1: Consecutive Reaction of $1\lambda^3$ -Phosphinine Complexes with Nucleophiles and Electrophiles	1176
15.15	Product Class 15: Benzo-Fused And Other Annulated Phosphinines U. Bergsträßer	
15.15	Product Class 15: Benzo-Fused And Other Annulated Phosphinines	1181
15.15.1	Product Subclass 1: Phosphinolines	1182
15.15.1.1	Synthesis by Ring-Closure Reactions	1182
15.15.1.1.1	By Formation of One P—C and One C—C Bond	1182
15.15.1.1.1.1	Method 1: From a Phosphaalkyne	1182
15.15.1.2	Synthesis by Ring Transformation	1182
15.15.1.2.1	Formal Exchange of Ring Members with Retention of the Ring Size	1182
15.15.1.2.1.1	Method 1: Oxygen–Phosphorus Exchange from a Naphthopyrylium Salt	1182
15.15.1.3	Aromatization	1183
15.15.1.3.1	Method 1: From a 1,2-Dihydrophosphinoline	1183
15.15.1.4	Synthesis by Substituent Modification	1183
15.15.1.4.1	Method 1: Substitution of Hydrogen	1183
15.15.2	Product Subclass 2: Isophosphinolines	1184
15.15.2.1	Synthesis by Ring-Closure Reactions	1184
15.15.2.1.1	By Formation of One P—C and One C—C Bond	1184
15.15.2.1.1.1	Method 1: From Phosphaalkenes and Phosphaalkynes	1184
15.15.2.2	Synthesis by Ring Transformation	1186
15.15.2.2.1	By Ring Contraction	1186
15.15.2.2.1.1	Method 1: From Polycyclic Phosphaalkenes	1186
15.15.2.3	Aromatization	1186
15.15.2.3.1	Method 1: Desulfuration of a 1,2-Dihydroisophosphinoline 2-Sulfide ...	1186
15.15.2.3.2	Method 2: Dehydrohalogenation of 2-Chloro-1,2-dihydroisophos- phinolines	1187
15.15.3	Product Subclass 3: Acridophosphanes	1188
15.15.3.1	Aromatization	1188
15.15.3.1.1	Method 1: From 5-Chloro-5,10-dihydroacridophosphanes	1188
15.15.4	Product Subclass 4: Benzo[c]phosphinolines	1189
15.15.4.1	Aromatization	1189
15.15.4.1.1	Method 1: From 5,6-Dihydrobenzo[c]phosphinolines	1189

Keyword Index	1191
Author Index	1249
Abbreviations	1315