

## Table of Contents

### Volume 2: Compounds of Groups 7–3 (Mn..., Cr..., V..., Ti..., Sc..., La..., Ac...)

|                   |   |      |
|-------------------|---|------|
| 2.10              | <b>Product Class 10: Organometallic Complexes of Titanium</b>                           |      |
| 2.10.19           | <b>Organometallic Complexes of Titanium (Update 1)</b>                                  | 2012 |
|                   | P. Bertus, F. Boeda, and M. S. M. Pearson-Long  |      |
| 2.10.19           | <b>Organometallic Complexes of Titanium (Update 1)</b>                                  | 1    |
| 2.10.19.1         | Titanium-Mediated Synthesis of Cyclopropyl Derivatives                                  | 1    |
| 2.10.19.1.1       | Synthesis of Cyclopropanes without Heteroatom Substitution                              | 2    |
| 2.10.19.1.1.1     | Method 1: Synthesis from Thioacetals and Thioethers                                     | 2    |
| 2.10.19.1.1.2     | Method 2: Synthesis from 1,1-Dihalides  | 7    |
| 2.10.19.1.2       | Synthesis of Cyclopropanols   | 8    |
| 2.10.19.1.2.1     | Synthesis from Carboxylic Acid Esters   | 8    |
| 2.10.19.1.2.1.1   | Method 1: Use of Grignard Reagents without Ligand Exchange                              | 8    |
| 2.10.19.1.2.1.2   | Method 2: Use of Alkenes and Grignard Reagents  | 13   |
| 2.10.19.1.2.1.2.1 | Variation 1: Bicyclo[n.1.0]alkan-1-ols from Unsaturated Carboxylic Acid Esters          | 13   |
| 2.10.19.1.2.1.2.2 | Variation 2: 2-(Hydroxyalkyl)cyclopropanols from Unsaturated Carboxylic Acid Esters     | 16   |
| 2.10.19.1.2.1.2.3 | Variation 3: Cyclopropanols from Carboxylic Acid Esters and Alkenes                     | 17   |
| 2.10.19.1.2.2     | Synthesis from Lactones and Other Acid Derivatives                                      | 20   |
| 2.10.19.1.2.2.1   | Method 1: Synthesis from Lactones   | 20   |
| 2.10.19.1.2.2.2   | Method 2: Synthesis from Other Acid Derivatives   | 21   |
| 2.10.19.1.3       | Synthesis of Cyclopropanone Hemiacetals   | 22   |
| 2.10.19.1.3.1     | Method 1: Synthesis from Cyclic Carbonates  | 22   |
| 2.10.19.1.4       | Synthesis of Cyclopropylamines  | 23   |
| 2.10.19.1.4.1     | Synthesis from Tertiary Amides  | 23   |
| 2.10.19.1.4.1.1   | Method 1: Use of Grignard Reagents without Ligand Exchange                              | 24   |
| 2.10.19.1.4.1.2   | Method 2: Use of Organozinc Reagents without Ligand Exchange                            | 28   |
| 2.10.19.1.4.1.3   | Method 3: Use of Alkenes and Grignard Reagents  | 28   |
| 2.10.19.1.4.1.3.1 | Variation 1: Bicyclo[n.1.0]alkan-1-amine Derivatives from Unsaturated Carboxylic Amides | 29   |
| 2.10.19.1.4.1.3.2 | Variation 2: 2-Azabicyclo[n.1.0]alkane Derivatives from Unsaturated Carboxylic Amides   | 31   |
| 2.10.19.1.4.1.3.3 | Variation 3: Cyclopropylamines from Tertiary Amides and Alkenes                         | 32   |

|                   |   |    |
|-------------------|---|----|
| 2.10.19.1.4.2     | Synthesis from Nitriles   | 35 |
| 2.10.19.1.4.2.1   | Method 1: Use of Grignard Reagents without Ligand Exchange                    | 35 |
| 2.10.19.1.4.2.1.1 | Variation 1: Alkylcyclopropylamines from Aliphatic Nitriles                   | 35 |
| 2.10.19.1.4.2.1.2 | Variation 2: 1-Aryl- and 1-Alkenylcyclopropylamines from Unsaturated Nitriles | 41 |
| 2.10.19.1.4.2.2   | Method 2: Use of Alkenes and Grignard Reagents                                | 43 |
| 2.10.19.1.4.2.2.1 | Variation 1: Bicyclic Cyclopropylamines from Unsaturated Nitriles             | 43 |
| 2.10.19.1.4.2.2.2 | Variation 2: Cyclopropylamines from Nitriles and Alkenes                      | 44 |
| 2.10.19.1.4.3     | Synthesis from Imides   | 45 |
| 2.10.19.1.4.3.1   | Method 1: Synthesis from Formimides or Cyclic Imides                          | 45 |

## Volume 4: Compounds of Group 15 (As, Sb, Bi) and Silicon Com- pounds

### 4.4 Product Class 4: Silicon Compounds

#### 4.4.1 Product Subclass 1: Disilenes 2012 A. Meltzer and D. Scheschkewitz

|           |  |    |
|-----------|--|----|
| 4.4.1     | Product Subclass 1: Disilenes  | 51 |
|           | Synthesis of Product Subclass 1                                      | 53 |
| 4.4.1.1   | Method 1: Synthesis of Acyclic Disilenes                             | 53 |
| 4.4.1.1.1 | Variation 1: Photolysis of Linear Trisilanes                         | 53 |
| 4.4.1.1.2 | Variation 2: Photolysis of Cyclotrisilanes                           | 54 |
| 4.4.1.1.3 | Variation 3: Reductive Dehalogenation of 1,1-Dihalosilanes           | 54 |
| 4.4.1.1.4 | Variation 4: Reductive Dehalogenation of 1,2-Dihalodisilanes         | 57 |
| 4.4.1.1.5 | Variation 5: Coupling of a 1,1-Dilithiosilane with Dihalosilanes     | 57 |
| 4.4.1.1.6 | Variation 6: Coupling of Alkali Metal Disilenides with Electrophiles | 58 |
| 4.4.1.1.7 | Variation 7: Addition to Disilynes                                   | 60 |
| 4.4.1.1.8 | Variation 8: Other Methods   | 60 |
| 4.4.1.2   | Method 2: Synthesis of Tetrasilabutadienes                           | 62 |
| 4.4.1.3   | Method 3: Synthesis of Cyclic Disilenes                              | 63 |

## Volume 8: Compounds of Group 1 (Li...Cs)

### 8.1 Product Class 1: Lithium Compounds

#### 8.1.31 Functionalized Organolithiums by Ring Opening of Heterocycles New M. Yus and F. Foubelo

|        |   |    |
|--------|---|----|
| 8.1.31 | Functionalized Organolithiums by Ring Opening of Heterocycles | 69 |
|--------|---|----|

|                |  |     |
|----------------|--|-----|
| 8.1.31.1       | Three-Membered Heterocycles .....  | 70  |
| 8.1.31.1.1     | Method 1: Oxiranes .....   | 70  |
| 8.1.31.1.2     | Method 2: Aziridines .....   | 75  |
| 8.1.31.2       | Four-Membered Heterocycles .....   | 77  |
| 8.1.31.2.1     | Method 1: Oxetanes .....   | 77  |
| 8.1.31.2.2     | Method 2: Azetidines .....   | 81  |
| 8.1.31.2.3     | Method 3: Thietanes .....  | 82  |
| 8.1.31.3       | Five-Membered Heterocycles .....   | 82  |
| 8.1.31.3.1     | Method 1: Oxygen-Containing Compounds .....  | 84  |
| 8.1.31.3.1.1   | Variation 1: Tetrahydrofurans .....  | 84  |
| 8.1.31.3.1.2   | Variation 2: Dioxolanes and Oxazolidines .....   | 85  |
| 8.1.31.3.1.3   | Variation 3: Benzo[ <i>b</i> ]furans .....   | 89  |
| 8.1.31.3.1.4   | Variation 4: Phthalans .....   | 91  |
| 8.1.31.3.2     | Method 2: Nitrogen-Containing Compounds: Pyrrolidines .....  | 94  |
| 8.1.31.3.3     | Method 3: Sulfur-Containing Compounds .....  | 96  |
| 8.1.31.4       | Six-Membered Heterocycles .....  | 97  |
| 8.1.31.4.1     | Method 1: Oxygen-Containing Compounds .....  | 97  |
| 8.1.31.4.1.1   | Variation 1: Saturated Oxygen-Containing Heterocycles .....  | 97  |
| 8.1.31.4.1.2   | Variation 2: 1-Benzopyrans .....   | 98  |
| 8.1.31.4.1.3   | Variation 3: 2-Benzopyrans .....   | 100 |
| 8.1.31.4.2     | Method 2: Nitrogen-Containing Compounds: Tetrahydroisoquinolines ..  | 101 |
| 8.1.31.4.3     | Method 3: Sulfur-Containing Compounds .....  | 103 |
| 8.1.31.5       | Seven-Membered Heterocycles .....  | 104 |
| 8.1.31.5.1     | Method 1: Dibenzo Oxygen-, Nitrogen-, and Sulfur-Containing Seven-Membered Heterocycles .....  | 104 |
| 8.1.31.5.2     | Method 2: Dinaphtho Oxygen- and Sulfur-Containing Seven-Membered Heterocycles .....  | 106 |
| 8.1.31.6       | Other Heterocycles .....   | 107 |
| 8.1.31.6.1     | Method 1: Benzodioxins, Benzoxathiins, Dihydrobenzodioxepins, and Dihydronaphthodioxocins .....  | 107 |
| 8.1.31.6.2     | Method 2: Phenoxathiin, Phenothiazine, and Thianthrene .....   | 110 |
| 8.1.32         | <b>Syntheses Mediated by <math>\alpha</math>-Lithiated Epoxides and Aziridines</b> <span style="border: 1px solid black; border-radius: 50%; padding: 2px;">New</span> |     |
|                | L. Degennaro, F. M. Perna, and S. Florio   |     |
| 8.1.32         | <b>Syntheses Mediated by <math>\alpha</math>-Lithiated Epoxides and Aziridines</b> .....   | 115 |
| 8.1.32.1       | Oxiranyllithiums .....   | 115 |
| 8.1.32.1.1     | 2-Alkyl-Substituted Oxiranyllithiums .....   | 116 |
| 8.1.32.1.1.1   | Method 1: Reactions with Electrophiles .....   | 116 |
| 8.1.32.1.1.1.1 | Variation 1: Stereospecific Trapping .....   | 116 |
| 8.1.32.1.1.1.2 | Variation 2: Asymmetric Lithiation and Trapping of <i>meso</i> -Oxiranes .....   | 117 |
| 8.1.32.1.1.1.3 | Variation 3: Coupling with Boronic Esters: Synthesis of Polyoxygenated Compounds .....   | 118 |

|                |              |  |     |
|----------------|--------------|--|-----|
| 8.1.32.1.1.2   | Method 2:    | Enantioselective $\alpha$ -Deprotonation–Rearrangement of <i>meso</i> -Oxiranes                                | 119 |
| 8.1.32.1.1.2.1 | Variation 1: | Synthesis of Bicyclic Alcohols by Transannular C–H Insertion   | 119 |
| 8.1.32.1.1.2.2 | Variation 2: | Synthesis of (–)-Xialenon  | 120 |
| 8.1.32.1.1.2.3 | Variation 3: | Effect of Lewis Acids on Transannular C–H Insertion Reactions  | 120 |
| 8.1.32.1.1.2.4 | Variation 4: | Enantioselective Rearrangement of <i>exo</i> -Norbornene Oxide to Nortricyclanol                               | 121 |
| 8.1.32.1.1.2.5 | Variation 5: | Transannular C–H Insertion in Lithiated 7-( <i>tert</i> -Butoxycarbonyl)-7-azanorbornene Oxide                 | 122 |
| 8.1.32.1.1.3   | Method 3:    | Construction of Nitrogen-Containing Heterocyclic Compounds by Desymmetrization of <i>meso</i> -Epoxides        | 123 |
| 8.1.32.1.1.4   | Method 4:    | Synthesis of Alkenes by Reductive Alkylation   | 124 |
| 8.1.32.1.1.4.1 | Variation 1: | Synthesis of Enantioenriched Unsaturated Diols   | 125 |
| 8.1.32.1.1.4.2 | Variation 2: | Synthesis of Enamines from Terminal Epoxides   | 126 |
| 8.1.32.1.1.5   | Method 5:    | Intramolecular Cyclopropanation of Unsaturated Terminal Epoxides (C=C Insertion)                               | 127 |
| 8.1.32.1.1.5.1 | Variation 1: | Stereospecific Synthesis of Bicyclic Alcohols  | 127 |
| 8.1.32.1.1.5.2 | Variation 2: | Synthesis of (–)-Sabina Ketone   | 128 |
| 8.1.32.1.1.6   | Method 6:    | Isomerization of $\alpha$ -Lithiated Epoxides to Ketones: 1,2-Hydrogen Migration                               | 129 |
| 8.1.32.1.2     |              | 2-Aryl-Substituted Oxiranyllithiums  | 130 |
| 8.1.32.1.2.1   | Method 1:    | Reactions of 2-Phenyloxiran-2-yllithiums   | 130 |
| 8.1.32.1.2.1.1 | Variation 1: | Stereospecific Trapping with Electrophiles   | 130 |
| 8.1.32.1.2.1.2 | Variation 2: | Stereoselective Synthesis of Antifungal Agents   | 131 |
| 8.1.32.1.2.2   | Method 2:    | Reactions of 3-Substituted 2-Phenyloxiran-2-yllithiums   | 132 |
| 8.1.32.1.2.2.1 | Variation 1: | Of 3-Methyl-2-phenyloxiran-2-yllithiums  | 132 |
| 8.1.32.1.2.2.2 | Variation 2: | Of 2-Lithiated 2,3-Diphenyloxiranes  | 132 |
| 8.1.32.1.2.3   | Method 3:    | Reactions of 2-Aryloxiran-2-yllithiums   | 134 |
| 8.1.32.1.2.4   | Method 4:    | Stereoselective Synthesis of Cyclopropanes   | 136 |
| 8.1.32.1.2.5   | Method 5:    | Stereoselective Synthesis of $\beta,\gamma$ -Epoxyhydroxylamines and 4-(Hydroxyalkyl)-1,2-oxazetidines         | 137 |
| 8.1.32.1.2.6   | Method 6:    | Stereocontrolled Synthesis of 1,2-Diols by Homologation of Boronic Esters with Lithiated 2-Phenyloxirane       | 138 |
| 8.1.32.1.2.7   | Method 7:    | Oxiranyl Anion Methodology Using Microflow Systems   | 140 |
| 8.1.32.1.3     |              | Lithiated Dihydrooxazol-2-yloxiranes   | 141 |
| 8.1.32.1.3.1   | Method 1:    | Reactions of 2-Lithiated 2-(4,5-Dihydrooxazol-2-yl)oxiranes  | 141 |
| 8.1.32.1.3.1.1 | Variation 1: | Configurational Stability of $\alpha$ -Lithiated (4,5-Dihydrooxazol-2-yl)oxiranes: Trapping with Electrophiles | 141 |
| 8.1.32.1.3.1.2 | Variation 2: | Synthesis of 2-Acyldihydrooxazoles   | 142 |
| 8.1.32.1.3.1.3 | Variation 3: | Synthesis of Cyclopropane-Fused $\gamma$ -Lactones   | 143 |
| 8.1.32.1.3.1.4 | Variation 4: | Synthesis of $\alpha$ -Epoxy- $\beta$ -amino Acids   | 144 |
| 8.1.32.1.3.2   | Method 2:    | Reactions of 2-Lithiated 3-(4,5-Dihydrooxazol-2-yl)oxiranes  | 147 |
| 8.1.32.1.3.2.1 | Variation 1: | Configurational Stability of 2-Lithiated 3-(4,5-Dihydrooxazol-2-yl)oxiranes: Trapping with Electrophiles       | 147 |
| 8.1.32.1.3.2.2 | Variation 2: | Synthesis of $\alpha,\beta$ -Epoxy- $\gamma$ -butyrolactones   | 148 |
| 8.1.32.1.3.2.3 | Variation 3: | Synthesis of $\alpha,\beta$ -Epoxy- $\gamma$ -amino Acids and $\alpha,\beta$ -Epoxy- $\gamma$ -butyrolactams   | 149 |

|                 |   |     |
|-----------------|---|-----|
| 8.1.32.1.3.3    | Method 3: 2-Lithiation of Terminal 3-(4,5-Dihydrooxazol-2-yl)oxiranes   | 152 |
| 8.1.32.1.4      | 2-Trifluoromethyl-Stabilized Oxiranyllithium  | 153 |
| 8.1.32.1.4.1    | Method 1: 2-(Trifluoromethyl)oxiranyllithium: Stereospecific Trapping with Electrophiles  | 153 |
| 8.1.32.1.4.2    | Method 2: 2-(Trifluoromethyl)oxiranyllithium as Precursor of the Oxiranylzinc   | 153 |
| 8.1.32.1.5      | Lactone-Derived Oxiranyllithiums  | 154 |
| 8.1.32.1.5.1    | Method 1: Reactions of Oxiranyllithiums Derived from $\alpha,\beta$ -Epoxy- $\gamma$ -butyrolactones  | 154 |
| 8.1.32.1.6      | Silyloxiranyllithiums   | 156 |
| 8.1.32.1.6.1    | Method 1: Lithiation of 3-Vinyloxiran-2-ylsilanes   | 156 |
| 8.1.32.1.6.2    | Method 2: Lithiation of 3-Alkyloxiran-2-ylsilanes   | 157 |
| 8.1.32.1.7      | 2-Sulfonyloxiranyllithiums  | 158 |
| 8.1.32.1.7.1    | Method 1: 2-Sulfonyloxiranyllithiums: Configurational Stability and Trapping with Electrophiles   | 158 |
| 8.1.32.1.7.2    | Method 2: Construction of Polycyclic Ethers   | 159 |
| 8.1.32.1.8      | 2-Lithiated 2-(Benzotriazol-1-yl)oxiranes   | 161 |
| 8.1.32.1.8.1    | Method 1: Synthesis of 2-(Benzotriazol-1-yl)oxiranyllithiums with Subsequent Trapping   | 161 |
| 8.1.32.1.9      | 2-Lithiated 2-(Benzothiazol-2-yl)oxiranes   | 161 |
| 8.1.32.1.9.1    | Method 1: Synthesis of 2-(Benzothiazol-2-yl)oxiranyllithiums with Subsequent Trapping   | 161 |
| 8.1.32.1.10     | Oxiranyllithiums by Transmetalation   | 163 |
| 8.1.32.1.10.1   | Method 1: Lithium–Tin Transmetalation   | 163 |
| 8.1.32.1.10.2   | Method 2: Lithium–Aluminum and Lithium–Zirconium Transmetalation  | 163 |
| 8.1.32.1.10.2.1 | Variation 1: Synthesis of Alkylated (Triphenylsilyl)alkenes: Reaction of a 2-Lithiated 2-(Triphenylsilyl)oxirane with Organoaluminum Reagents | 163 |
| 8.1.32.1.10.2.2 | Variation 2: Insertion of Metalated Epoxides into Zirconacycles   | 164 |
| 8.1.32.1.10.2.3 | Variation 3: Insertion of Metalated Epoxynitriles into Chlorobis( $\eta^5$ -cyclopentadienyl)organozirconium Reagents                         | 165 |
| 8.1.32.1.10.2.4 | Variation 4: Insertion of 2-Lithiated 2-(Phenylsulfonyl)oxiranes into Alkenylchlorobis( $\eta^5$ -cyclopentadienyl)zirconium Reagents         | 166 |
| 8.1.32.1.11     | Remotely Stabilized Lithiated Epoxides  | 168 |
| 8.1.32.1.11.1   | Method 1: Remotely Stabilized Lithiated Epoxides: Reactions with Aldehydes  | 168 |
| 8.1.32.1.11.2   | Method 2: Synthesis of Xylobovide   | 169 |
| 8.1.32.2        | $\alpha$ -Lithiated Aziridines  | 170 |
| 8.1.32.2.1      | Lithiation–Trapping Sequence of Aziridines with an Electron-Withdrawing Group at the Carbon Atom  | 171 |
| 8.1.32.2.1.1    | Method 1: Synthesis of 1-Alkylaziridine-2-carboxylates  | 171 |
| 8.1.32.2.1.2    | Method 2: Synthesis of 1-Alkylaziridine-2-carbothioates   | 172 |

|                |              |   |     |
|----------------|--------------|---|-----|
| 8.1.32.2.1.3   | Method 3:    | Reactions of 2-(4,5-Dihydrooxazol-2-yl)-Substituted 1-Phenylaziridines .....  | 173 |
| 8.1.32.2.1.4   | Method 4:    | Reactions of 2,3-Dihetaryl-Substituted 1-Phenylaziridines ...   | 174 |
| 8.1.32.2.1.5   | Method 5:    | Reactions of 2-(4,5-Dihydrooxazol-2-yl)-Substituted 1-Tritylaziridines .....  | 176 |
| 8.1.32.2.1.5.1 | Variation 1: | Synthesis of <i>N</i> -Tritylepimino- $\gamma$ -butyrolactones .....  | 177 |
| 8.1.32.2.1.6   | Method 6:    | Reactions of 2-(4,5-Dihydrooxazol-2-yl)-Substituted 1-Benzylaziridines .....  | 178 |
| 8.1.32.2.1.7   | Method 7:    | Reactions of 2-(4,5-Dihydrooxazol-2-yl)-Substituted 1-(1-Phenylethyl)aziridines .....   | 179 |
| 8.1.32.2.1.8   | Method 8:    | Synthesis of <i>C</i> -Substituted 1-Phenyl-2-sulfonylaziridines .....  | 181 |
| 8.1.32.2.2     |              | Lithiation of Aziridines with an Electron-Withdrawing Group at Nitrogen .....   | 182 |
| 8.1.32.2.2.1   | Method 1:    | <i>C</i> -Alkylation of 1-( <i>tert</i> -Butylsulfonyl)-2-phenylaziridine .....   | 182 |
| 8.1.32.2.2.1.1 | Variation 1: | Synthesis of 2-Substituted 1-( <i>tert</i> -Butylsulfonyl)-2-phenylaziridines Using Microreactor Technology .....                           | 183 |
| 8.1.32.2.2.1.2 | Variation 2: | Synthesis of ( <i>tert</i> -Butylsulfonyl)amino Alcohols .....  | 184 |
| 8.1.32.2.2.2   | Method 2:    | Synthesis of 2-Alkyl-Substituted 1-( <i>tert</i> -Butylsulfonyl)-3-(trimethylsilyl)aziridines .....   | 185 |
| 8.1.32.2.2.3   | Method 3:    | Synthesis of 2-Substituted 1-( <i>tert</i> -Butylsulfonyl)aziridines ...  | 186 |
| 8.1.32.2.2.3.1 | Variation 1: | Synthesis of 2-Substituted 1-( <i>tert</i> -Butylsulfonyl)aziridines Using Microreactor Technology .....                                    | 187 |
| 8.1.32.2.2.4   | Method 4:    | Synthesis of <i>trans</i> -2,3-Disubstituted 1-( <i>tert</i> -Butylsulfonyl)aziridines .....  | 188 |
| 8.1.32.2.2.5   | Method 5:    | Synthesis of <i>C</i> -Substituted 1-(2,4,6-Triisopropylphenylsulfonyl)aziridines .....   | 189 |
| 8.1.32.2.2.6   | Method 6:    | Reductive Alkylation (or Alkylative Ring Opening) of 1-Sulfonylaziridinylolithiums .....  | 191 |
| 8.1.32.2.2.6.1 | Variation 1: | Synthesis of Allylic Sulfonamides .....   | 191 |
| 8.1.32.2.2.6.2 | Variation 2: | Synthesis of Alkynylamines .....  | 193 |
| 8.1.32.2.2.6.3 | Variation 3: | Allylic Amino Alcohols and Amino Ethers by Organolithium-Induced Alkylative Ring Opening of 1-Sulfonyl-Protected Aziridinyl Ethers .....    | 195 |
| 8.1.32.2.2.6.4 | Variation 4: | Allylic Amino Alcohols and Amino Ethers by Organolithium-Induced Alkylative Ring Opening of 1,4-Dimethoxybut-2-ene-Derived Aziridines ..... | 196 |
| 8.1.32.2.2.7   | Method 7:    | Eliminative Dimerization of Lithiated 1-( <i>tert</i> -Butylsulfonyl)aziridines .....   | 197 |
| 8.1.32.2.2.7.1 | Variation 1: | Synthesis of 2-Ene-1,4-diamines .....   | 197 |
| 8.1.32.2.2.8   | Method 8:    | Intramolecular Cyclopropanation of Lithiated 1-( <i>tert</i> -Butylsulfonyl)aziridines .....  | 198 |
| 8.1.32.2.2.8.1 | Variation 1: | Synthesis of 2-Aminobicyclo[3.1.0]hexanes .....   | 198 |
| 8.1.32.2.2.9   | Method 9:    | Lithiation of 1-( <i>tert</i> -Butoxycarbonyl)aziridines .....  | 200 |
| 8.1.32.2.2.9.1 | Variation 1: | Synthesis of 2-Silylaziridines .....  | 200 |
| 8.1.32.2.2.9.2 | Variation 2: | Synthesis of <i>trans</i> -Configured Aziridine-2-carboxylates .....  | 200 |
| 8.1.32.2.2.9.3 | Variation 3: | Synthesis of <i>trans</i> -Configured Aziridin-2-ylphosphonates .....   | 202 |
| 8.1.32.2.2.9.4 | Variation 4: | Synthesis of <i>N</i> - <i>tert</i> -Butoxycarbonyl 1,2-Amino Alcohols .....  | 202 |
| 8.1.32.2.3     |              | Lithiation of Aziridines with an Electron-Donating Group on Nitrogen .....  | 204 |

|                |  |   |            |
|----------------|--|---|------------|
| 8.1.32.2.3.1   | Method 1:  | Synthesis of <i>cis</i> - and <i>trans</i> -Configured C-Substituted 1-Alkyl-2,3-diphenylaziridines | 204        |
| 8.1.32.2.3.2   | Method 2:  | Synthesis of C-Substituted 1-Alkyl-2-methyleneaziridines  | 205        |
| 8.1.32.2.3.2.1 | Variation 1:   | Synthesis of Chiral Nonracemic C-Substituted 1-Alkyl-2-methyleneaziridines                          | 206        |
| 8.1.32.2.3.3   | Method 3:  | Synthesis of C-Substituted Aziridine–Borane Complexes   | 207        |
| 8.1.32.2.3.3.1 | Variation 1:   | Of 1-[2-( <i>tert</i> -Butyldimethylsiloxy)ethyl]aziridine–Borane Complexes                         | 207        |
| 8.1.32.2.3.3.2 | Variation 2:   | Of 1-Alkyl-2-phenylaziridine–Borane Complexes   | 209        |
| 8.1.33         | <b>Transition-Metal-Catalyzed Carbon–Carbon Bond Formation with Organolithiums</b> |   | <b>New</b> |
|                | G. Manolikakes   |   |            |
| 8.1.33         | <b>Transition-Metal-Catalyzed Carbon–Carbon Bond Formation with Organolithiums</b> |   | 215        |
| 8.1.33.1       | Copper-Catalyzed Reactions   |   | 215        |
| 8.1.33.1.1     | Method 1:  | Copper-Catalyzed Conjugate Addition   | 216        |
| 8.1.33.1.2     | Method 2:  | Copper-Catalyzed Alkylation   | 217        |
| 8.1.33.1.2.1   | Variation 1:   | Copper-Catalyzed Asymmetric Allylation  | 217        |
| 8.1.33.1.3     | Method 3:  | Copper-Catalyzed Coupling of Organolithium Reagents with $\alpha$ -Lithiated Cyclic Enol Ethers     | 219        |
| 8.1.33.2       | Palladium-Catalyzed Reactions  |   | 219        |
| 8.1.33.2.1     | Method 1:  | Palladium-Catalyzed Cross-Coupling Reactions with Aryl and Vinyl Halides                            | 220        |
| 8.1.33.2.2     | Method 2:  | Palladium-Catalyzed Coupling Reaction of Aryllithium Reagents with 1-Bromo-2-methylbut-3-en-2-ol    | 221        |
| 8.1.33.3       | Iron-Catalyzed Reactions   |   | 222        |
| 8.1.33.3.1     | Method 1:  | Iron-Catalyzed Cross-Coupling Reactions   | 222        |
| 8.1.33.3.2     | Method 2:  | Iron-Catalyzed Carbolithiation of Alkynes   | 223        |

## Volume 16: Six-Membered Heteroarenes with Two Identical Heteroatoms

|              |  |  |             |
|--------------|--|--|-------------|
| 16.2         | <b>Product Class 2: 1,4-Dioxins and Benzo- and Dibenzo-Fused Derivatives</b> |  |             |
| 16.2.4       | <b>1,4-Dioxins and Benzo- and Dibenzo-Fused Derivatives</b>                  |  | <b>2012</b> |
|              | S. M. Sakya and J. Yang  |  |             |
| 16.2.4       | <b>1,4-Dioxins and Benzo- and Dibenzo-Fused Derivatives</b>                  |  | 227         |
| 16.2.4.1     | Synthesis by Ring-Closure Reactions  |  | 227         |
| 16.2.4.1.1   | By Formation of Two O–C Bonds  |  | 227         |
| 16.2.4.1.1.1 | Fragments O–C–C–O and C–C  |  | 227         |

|                |   |     |
|----------------|---|-----|
| 16.2.4.1.1.1.1 | Method 1: Dibenzo[ <i>b,e</i> ][1,4]dioxins by Base-Induced Coupling of Benzene-1,2-diols with Activated Fluorobenzenes                             | 227 |
| 16.2.4.1.1.2   | Fragments O—C—C and O—C—C   | 229 |
| 16.2.4.1.1.2.1 | Method 1: Substituted 1,4-Dioxins by Reaction of Methyl 3-Chloro-2-oxo-3-phenylpropanoate with Potassium Phthalimide or Sodium Imidazolid           | 229 |
| 16.2.4.1.2     | By Formation of One C—C Bond  | 230 |
| 16.2.4.1.2.1   | Fragment C—O—C—C—O—C  | 230 |
| 16.2.4.1.2.1.1 | Method 1: 1,4-Benzodioxins by Ring-Closing Metathesis of Divinyl Ethers   | 230 |
| 16.2.4.2       | Aromatization   | 231 |
| 16.2.4.2.1     | Method 1: 1,4-Benzodioxins by Isomerization of Exocyclic Alkenes  | 231 |
| 16.2.4.2.2     | Method 2: Dibenzo[ <i>b,e</i> ][1,4]dioxins from the Diels–Alder Reactions of 1,4-Benzodioxin and Benzo[ <i>b</i> ]furo[3,4- <i>e</i> ][1,4]dioxins | 232 |
| 16.2.4.3       | Synthesis by Substituent Modification   | 235 |
| 16.2.4.3.1     | Substitution of Existing Substituents   | 235 |
| 16.2.4.3.1.1   | Of Hydrogen   | 235 |
| 16.2.4.3.1.1.1 | Method 1: Vilsmeier Reaction of 2-Phenyl-1,4-benzodioxin  | 235 |
| 16.2.4.3.1.1.2 | Method 2: Diels–Alder Reaction of 2,2'-Bi-1,4-benzodioxin   | 235 |
| 16.2.4.3.1.2   | Of Metals   | 236 |
| 16.2.4.3.1.2.1 | Method 1: Stille Coupling of 2-(Trimethylstannyl)-1,4-benzodioxin with a Bromoalkene  | 236 |
| 16.2.4.3.1.3   | Of Halogens   | 238 |
| 16.2.4.3.1.3.1 | Method 1: Alkylation of 2-Bromo-1,4-benzodioxin by Lithium–Halogen Exchange   | 238 |
| 16.2.4.3.2     | Modification of Substituents  | 238 |
| 16.2.4.3.2.1   | Method 1: Alkylation of 1,4-Benzodioxin-6,7-dicarbaldehyde  | 238 |

### 16.3 Product Class 3: 1,2-Dithiins

|                |  |             |
|----------------|--|-------------|
| 16.3.5         | <b>1,2-Dithiins</b>  | <b>2012</b> |
|                | F. K. Yoshimoto and Q. Li  |             |
| 16.3.5         | <b>1,2-Dithiins</b>  | 241         |
| 16.3.5.1       | Synthesis by Ring-Closure Reactions  | 244         |
| 16.3.5.1.1     | By Formation of One S—S and Two S—C Bonds                                  | 244         |
| 16.3.5.1.1.1   | Fragment C—C—C—C and Two S Fragments                                       | 244         |
| 16.3.5.1.1.1.1 | Method 1: Addition of Sulfur to 6-Nitroperlylo[1,12- <i>bcd</i> ]thiophene | 244         |
| 16.3.5.1.1.1.2 | Method 2: Addition of Sulfur to 2-(Trimethylsiloxy)buta-1,3-diene          | 244         |
| 16.3.5.1.2     | By Formation of One S—S and One S—C Bond                                   | 245         |
| 16.3.5.1.2.1   | Fragments S—C—C—C and S  | 245         |



|                |  |     |
|----------------|--|-----|
| 16.3.5.1.2.1.1 | Method 1: Reaction of 2-(2-Phenylvinyl)-3-vinylthiirane Promoted by Acetonitrile(pentacarbonyl)tungsten(0) ..... | 245 |
| 16.3.5.1.3     | By Formation of One S—S and One C—C Bond .....   | 245 |
| 16.3.5.1.3.1   | Fragments S—C—C—C and S—C .....  | 245 |
| 16.3.5.1.3.1.1 | Method 1: Thermal Dimerization of $\alpha,\beta$ -Unsaturated $\beta$ -Arylsulfanyl Thioketones .....            | 245 |
| 16.3.5.1.3.1.2 | Method 2: Cobalt(II)-Mediated Dimerization of $\alpha,\beta$ -Unsaturated Thioacylsilanes .....                  | 246 |
| 16.3.5.1.3.1.3 | Method 3: Dimerization via Diels–Alder Reaction of Thioaldehydes .....   | 246 |
| 16.3.5.1.3.2   | Fragments S—C—C and S—C—C .....  | 247 |
| 16.3.5.1.3.2.1 | Method 1: Thionation–Dimerization of 1,3-Dihydro-2 <i>H</i> -indol-2-one .....                                   | 247 |
| 16.3.5.1.3.2.2 | Method 2: Manganese(IV) Oxide Promoted Oxidative Dimerization .....  | 248 |
| 16.3.5.1.4     | By Formation of One S—S Bond .....   | 248 |
| 16.3.5.1.4.1   | Fragment S—C—C—C—C—S .....   | 248 |
| 16.3.5.1.4.1.1 | Method 1: Polycyclization of Diynes .....  | 248 |
| 16.3.5.1.4.1.2 | Method 2: <i>N</i> -Bromosuccinimide-Induced Ring Formation .....  | 249 |
| 16.3.5.1.4.1.3 | Method 3: Cyclization of Dibromides Promoted by Phase-Transfer Catalysts .....                                   | 250 |
| 16.3.5.1.4.1.4 | Method 4: Cyclization of Dichlorides by Tandem Michael–Nucleophilic Substitution Processes .....                 | 251 |
| 16.3.5.1.5     | By Formation of One C—C Bond .....   | 252 |
| 16.3.5.1.5.1   | Fragment C—C—S—S—C—C .....   | 252 |
| 16.3.5.1.5.1.1 | Method 1: Cyclization via Ring-Closing Metathesis of Alkenes .....   | 252 |
| 16.3.5.2       | Synthesis by Ring Transformation .....   | 252 |
| 16.3.5.2.1     | Method 1: Ring Contraction Promoted by Photolysis .....  | 252 |
| 16.3.5.3       | Synthesis by Other Methods .....   | 253 |
| 16.3.5.3.1     | Method 1: Rearrangement Promoted by Photolysis .....   | 253 |
| 16.3.5.4       | Applications of 1,2-Dithiins in Organic Synthesis .....  | 253 |
| 16.3.5.4.1     | Reaction with Transition Metals .....  | 254 |
| 16.3.5.4.1.1   | Method 1: Reaction with Organometallic Complexes .....   | 254 |
| 16.3.5.4.1.2   | Method 2: Reaction with Copper Metal .....   | 255 |
| 16.3.5.4.2     | Reaction with Lewis Acids .....  | 255 |
| 16.3.5.4.2.1   | Method 1: Reaction Promoted by Aluminum Trichloride .....  | 255 |
| 16.3.5.4.2.2   | Method 2: Reaction Promoted by Boron Trifluoride .....   | 256 |
| 16.3.5.4.3     | Reaction with Diazo Compounds .....  | 257 |
| 16.3.5.4.3.1   | Method 1: Reaction Promoted by Rhodium(II) Acetate .....   | 257 |
| 16.3.5.4.3.2   | Method 2: Reaction Promoted by Copper(I) Chloride .....  | 258 |
| 16.3.5.4.4     | Reaction with Alkynes .....  | 259 |
| 16.3.5.4.4.1   | Method 1: Reaction Promoted by Bis(acetylacetonato)nickel(II) .....  | 259 |
| 16.3.5.4.5     | Reaction with Enzymes .....  | 260 |
| 16.3.5.4.5.1   | Method 1: Reaction with a Toluene Dioxygenase .....  | 260 |

## Volume 17: Six-Membered Hetarenes with Two Unlike or More than Two Heteroatoms and Fully Unsaturated Larger-Ring Heterocycles

|                |   |     |
|----------------|---|-----|
| 17.4           | <b>Product Class 4: Seven-Membered Hetarenes with One Heteroatom</b>                                    |     |
| 17.4.1.5       | <b>Oxepins</b> <span style="float: right;">2012</span>  |     |
|                | J. Hong   |     |
| 17.4.1.5       | <b>Oxepins</b> .....  | 263 |
| 17.4.1.5.1     | Synthesis by Ring-Closure Reactions .....   | 263 |
| 17.4.1.5.1.1   | By Formation of One O—C and One C—C Bond .....  | 263 |
| 17.4.1.5.1.1.1 | Method 1: From a 3-(Dimethylamino)-1-phenylprop-2-en-1-one and<br>Arylidene malononitriles .....        | 263 |
| 17.4.1.5.2     | Synthesis by Ring Transformation .....  | 263 |
| 17.4.1.5.2.1   | Method 1: By Valence Isomerization of 3-Oxaquadricyclanes .....   | 263 |
| 17.4.1.5.2.2   | Method 2: By Valence Isomerization of 7-Oxanorbornadienes .....   | 264 |
| 17.4.1.5.2.3   | Method 3: By Ring Enlargement of Furans with Diethyl Acetylenedicarboxylate .....                       | 265 |
| 17.4.1.5.2.4   | Method 4: By Valence Isomerization of Benzene Oxide .....   | 265 |
| 17.4.1.5.2.5   | Method 5: By Ring Enlargement of a 2-[(Prop-2-ynoxy)methyl]furan ..                                     | 266 |
| 17.4.1.5.2.6   | Method 6: By Ring Enlargement of a Cyclohexa-2,5-diene-1,4-diol via<br>S <sub>N</sub> 2' Reaction ..... | 267 |
| 17.4.1.5.3     | Aromatization .....   | 267 |
| 17.4.1.5.3.1   | Method 1: By Dehydrogenation .....  | 267 |
| 17.4.2.5       | <b>Benzoxepins</b> <span style="float: right;">2012</span>  |     |
|                | J. Hong   |     |
| 17.4.2.5       | <b>Benzoxepins</b> .....  | 271 |
| 17.4.2.5.1     | Synthesis by Ring-Closure Reactions .....   | 271 |
| 17.4.2.5.1.1   | By Formation of One O—C and Two C—C Bonds .....   | 271 |
| 17.4.2.5.1.1.1 | Method 1: From a Betaine and Diethyl Acetylenedicarboxylate .....                                       | 271 |
| 17.4.2.5.1.2   | By Formation of One O—C and One C—C Bond .....  | 271 |
| 17.4.2.5.1.2.1 | Method 1: From Dinrotoluenes and Salicylaldehydes .....   | 271 |
| 17.4.2.5.1.3   | By Formation of Two C—C Bonds .....   | 273 |
| 17.4.2.5.1.3.1 | Method 1: By Annulation of a Boronic Acid with Dimethyl Acetylenedicarboxylate .....                    | 273 |
| 17.4.2.5.1.4   | By Formation of One O—C Bond .....  | 274 |
| 17.4.2.5.1.4.1 | Method 1: From 2-Alkyl-3-[2-(iodoethynyl)phenyl]oxiranes .....  | 274 |

|                    |   |      |
|--------------------|---|------|
| 17.4.2.5.1.4.2     | Method 2: By Cyclization of 2-[2-(2-Bromophenyl)vinyl]phenols and Related Compounds   | 275  |
| 17.4.2.5.1.5       | By Formation of One C—C Bond  | 277  |
| 17.4.2.5.1.5.1     | Method 1: By Base-Catalyzed Cyclocondensation of Methyl (2 <i>E</i> )-4-(2-Formylphenoxy)but-2-enoate   | 277  |
| 17.4.2.5.2         | Synthesis by Ring Transformation  | 278  |
| 17.4.2.5.2.1       | By Ring Enlargement   | 278  |
| 17.4.2.5.2.1.1     | Method 1: Of a Benzofuran with 1-Phenyl-2-tosylacetylene  | 278  |
| 17.4.2.5.2.1.2     | Method 2: Of Xanthenes by Dehydration   | 278  |
| 17.4.2.5.2.1.3     | Method 3: Of 2-Diazo-3',6'-bis(diethylamino)spiro[indene-1,9'-xanthen]-3(2 <i>H</i> )-one (Rhodamine BBN)   | 280  |
| 17.4.2.5.2.1.4     | Method 4: Of a Dihydrofuran by Rearrangement  | 281  |
| 17.4.2.5.2.1.5     | Method 5: Of Tetrahydrobenzo[ <i>b</i> ]cyclopropa[ <i>e</i> ]pyran-1-carboxylates  | 282  |
| 17.4.2.5.3         | Synthesis by Substituent Modification   | 283  |
| 17.4.2.5.3.1       | Substitution of Existing Substituents   | 283  |
| 17.4.2.5.3.1.1     | Method 1: Condensation of Dibenz[ <i>b,f</i> ]oxepin-10(11 <i>H</i> )-one with 3-Methylbut-2-enal   | 283  |
| 17.4.2.5.3.1.2     | Method 2: Vilsmeier-Type Chloroformylation of Dibenz[ <i>b,f</i> ]oxepin-10(11 <i>H</i> )-ones  | 283  |
| 17.4.2.5.3.1.3     | Method 3: Reaction of Dibenz[ <i>b,f</i> ]oxepin-10(11 <i>H</i> )-one with Base and Carbon Disulfide/Iodomethane or Dimethyl Trithiocarbonate; Annulation of the Products | 284  |
| 17.4.2.5.3.1.4     | Method 4: 1 <i>H</i> -Dibenz[2,3:6,7]oxepino[4,5- <i>b</i> ]pyrroles by Annulation of 11-(Hydrazonoethylidene)dibenz[ <i>b,f</i> ]oxepin-10-ones                          | 290  |
| 17.4.2.5.3.1.5     | Method 5: 1 <i>H</i> -Dibenz[2,3:6,7]oxepino[4,5- <i>d</i> ]imidazoles by Oxidation/Annulation of Dibenz[ <i>b,f</i> ]oxepin-10(11 <i>H</i> )-ones                        | 291  |
| 17.4.5.5           | <b>Azepines, Cyclopentazepines, and Phosphorus Analogues</b>  | 2012 |
|                    | J. E. Camp  |      |
| 17.4.5.5           | <b>Azepines, Cyclopentazepines, and Phosphorus Analogues</b>  | 295  |
| 17.4.5.5.1         | Synthesis by Ring-Closure Reactions   | 295  |
| 17.4.5.5.1.1       | By Formation of One C—C Bond  | 295  |
| 17.4.5.5.1.1.1     | Method 1: Azepine Formation via Copper-Mediated Cyclization of 2-Azahepta-2,4-dien-6-ynyl Anions  | 295  |
| 17.4.5.5.1.1.2     | Method 2: 3 <i>H</i> -Azepines via Deprotonation of 2-Aza-1,3,5-trienes   | 296  |
| 17.4.5.5.2         | Synthesis by Ring Transformation  | 297  |
| 17.4.5.5.2.1       | By Ring Enlargement   | 297  |
| 17.4.5.5.2.1.1     | Of Five-Membered Heterocycles   | 297  |
| 17.4.5.5.2.1.1.1   | Method 1: Thermal Isomerization of 3-Azaquadracyclanes  | 297  |
| 17.4.5.5.2.1.1.1.1 | Variation 1: Thermal Isomerization of a Cyclobutane 3-Azaquadracyclane  | 297  |
| 17.4.5.5.2.1.2     | Of Six-Membered Arenes  | 298  |
| 17.4.5.5.2.1.2.1   | Method 1: Intramolecular Insertion of Arylnitrenes  | 298  |

|                    |  |      |
|--------------------|--|------|
| 17.4.5.5.2.1.2.1.1 | Variation 1: Photolytic Decomposition of Aryl Azides                                     | 298  |
| 17.4.5.5.2.1.2.1.2 | Variation 2: Rearrangement of Nitroarenes  | 300  |
| 17.4.5.5.3         | Synthesis by Substituent Modification  | 301  |
| 17.4.5.5.3.1       | Substitution of Existing Substituents  | 301  |
| 17.4.5.5.3.1.1     | Of Hydrogen  | 301  |
| 17.4.5.5.3.1.1.1   | Method 1: Tautomerization  | 301  |
| 17.4.5.5.3.1.1.1.1 | Variation 1: Rearrangement of 2 <i>H</i> - to 3 <i>H</i> -Azepines                       | 301  |
| 17.4.5.5.3.1.1.2   | Method 2: C-Halogenation   | 302  |
| 17.4.5.5.3.1.1.2.1 | Variation 1: C-Halogenation with <i>N</i> -Bromosuccinimide                              | 302  |
| 17.4.5.5.3.1.1.3   | Method 3: C-Alkylsulfanylation   | 302  |
| 17.4.5.5.3.1.1.4   | Method 4: C-Amination  | 303  |
| 17.4.5.5.3.1.1.5   | Method 5: C-Alkoxylation   | 304  |
| 17.4.5.5.3.1.2     | Of Heteroatoms   | 305  |
| 17.4.5.5.3.1.2.1   | Method 1: Of Alkoxy Groups   | 305  |
| 17.4.5.5.3.1.2.1.1 | Variation 1: Of Activated Organooxy Groups   | 307  |
| 17.4.6.10          | <b>Benzazepines and Their Group 15 Analogues</b>   | 2012 |
|                    | J. E. Camp   |      |
| 17.4.6.10          | <b>Benzazepines and Their Group 15 Analogues</b>   | 311  |
| 17.4.6.10.1        | 1 <i>H</i> -1-Benzazepines   | 312  |
| 17.4.6.10.1.1      | Synthesis by Ring-Closure Reactions  | 312  |
| 17.4.6.10.1.1.1    | By Formation of Two C—C Bonds and One C—N Bond   | 312  |
| 17.4.6.10.1.1.1.1  | Method 1: By Condensation between 2-Fluoroaniline and Aryl Methyl Ketones                | 312  |
| 17.4.6.10.1.1.2    | By Formation of One N—C and One C—C Bond   | 312  |
| 17.4.6.10.1.1.2.1  | Method 1: From 5-[( <i>E</i> )-(2-Dimethylamino)vinyl]-2,1,3-benzoselenadiazol-4-amine   | 312  |
| 17.4.6.10.1.1.3    | By Formation of Two C—C Bonds  | 313  |
| 17.4.6.10.1.1.3.1  | Method 1: By Thermal Cycloaddition of Dimethyl Acetylenedicarboxylate with Methylindoles | 313  |
| 17.4.6.10.1.1.3.2  | Method 2: From the Reaction of Phosphonium Ylides  | 314  |
| 17.4.6.10.1.1.4    | By Formation of One C—N Bond   | 316  |
| 17.4.6.10.1.1.4.1  | Method 1: By Intramolecular Addition of Anilines   | 316  |
| 17.4.6.10.1.2      | Synthesis by Ring Transformation   | 318  |
| 17.4.6.10.1.2.1    | By Ring Enlargement  | 318  |
| 17.4.6.10.1.2.1.1  | Method 1: By Ring Expansion of Activated Quinolines                                      | 318  |
| 17.4.6.10.2        | 2-Benzazepines   | 319  |
| 17.4.6.10.2.1      | Synthesis by Ring-Closure Reactions  | 319  |
| 17.4.6.10.2.1.1    | By Formation of One N—C and One C—C Bond   | 319  |

|                     |              |   |     |
|---------------------|--------------|---|-----|
| 17.4.6.10.2.1.1.1   | Method 1:    | By a Tandem Ritter/Houben–Hoesch Process  | 319 |
| 17.4.6.10.2.1.1.2   | Method 2:    | By Reaction of 4-Chloro-2-oxo-2 <i>H</i> -1-benzopyran-3-carbaldehyde with Benzylamines   | 320 |
| 17.4.6.10.2.1.2     |              | By Formation of One C—C Bond  | 322 |
| 17.4.6.10.2.1.2.1   | Method 1:    | By Cyclization of 2-Azahepta-2,4-dien-6-ynyls   | 322 |
| 17.4.6.10.3         |              | 5 <i>H</i> -Dibenz[ <i>b,d</i> ]azepines  | 323 |
| 17.4.6.10.3.1       |              | Synthesis by Substituent Modification   | 323 |
| 17.4.6.10.3.1.1     | Method 1:    | By Rhodium-Catalyzed Decarbonylative Cycloaddition  | 323 |
| 17.4.6.10.4         |              | 11 <i>H</i> -Dibenz[ <i>b,e</i> ]azepines   | 324 |
| 17.4.6.10.4.1       |              | Synthesis by Ring-Closure Reactions   | 324 |
| 17.4.6.10.4.1.1     |              | By Formation of One N—C and One C—C Bond  | 324 |
| 17.4.6.10.4.1.1.1   | Method 1:    | By Condensation of 2,6-Dimethylaniline with Phenanthrene-9,10-dione   | 324 |
| 17.4.6.10.4.1.2     |              | By Formation of One C—C Bond  | 324 |
| 17.4.6.10.4.1.2.1   | Method 1:    | By Bischler–Napieralski Cyclodehydration of <i>N</i> -(2-Benzylphenyl)-2-chloroacetamide  | 324 |
| 17.4.6.10.4.1.2.2   | Method 2:    | By Friedel–Crafts Cyclization of 2-Allyl- <i>N</i> -benzylanilines  | 325 |
| 17.4.6.10.4.1.2.3   | Method 3:    | By Acid-Mediated Cyclization of Benzylic Alcohols   | 326 |
| 17.4.6.10.5         |              | 5 <i>H</i> -Dibenz[ <i>c,e</i> ]azepines  | 328 |
| 17.4.6.10.5.1       |              | Synthesis by Ring-Closure Reactions   | 328 |
| 17.4.6.10.5.1.1     |              | By Formation of Two N—C Bonds   | 328 |
| 17.4.6.10.5.1.1.1   | Method 1:    | Ring Closure of 2,2'-Difunctionalized Biaryls with Chiral Amines under Acidic Conditions  | 328 |
| 17.4.6.10.6         |              | 5 <i>H</i> -Dibenz[ <i>b,f</i> ]azepines  | 333 |
| 17.4.6.10.6.1       |              | Synthesis by Ring-Closure Reactions   | 333 |
| 17.4.6.10.6.1.1     |              | By Formation of One C—C and One C—N Bond  | 333 |
| 17.4.6.10.6.1.1.1   | Method 1:    | By a Palladium-Catalyzed Tandem Process   | 333 |
| 17.4.6.10.6.1.2     |              | By Formation of One C—C Bond  | 334 |
| 17.4.6.10.6.1.2.1   | Method 1:    | By Palladium-Catalyzed Intramolecular Amination   | 334 |
| 17.4.6.10.6.1.2.2   | Method 2:    | Friedel–Crafts Acylation  | 335 |
| 17.4.6.10.6.2       |              | Synthesis by Ring Transformation  | 336 |
| 17.4.6.10.6.2.1     |              | By Ring Enlargement   | 336 |
| 17.4.6.10.6.2.1.1   | Method 1:    | From 1-Arylindoles  | 336 |
| 17.4.6.10.6.2.1.1.1 | Variation 1: | Ring Expansion of 6-Methoxy-1-phenylindole  | 336 |
| 17.4.6.10.6.3       |              | Aromatization   | 336 |
| 17.4.6.10.6.3.1     | Method 1:    | Bromination–Dehydrobromination  | 336 |
| 17.4.6.10.6.3.1.1   | Variation 1: | Dehalogenation of 5-Acetyl-10,11-dibromo-10,11-dihydro-5 <i>H</i> -dibenz[ <i>b,f</i> ]azepine with 1,2-Diphenylethane-1,2-diyldisodium | 336 |

|                     |  |     |
|---------------------|--|-----|
| 17.4.6.10.6.4       | Synthesis by Substituent Modification  | 337 |
| 17.4.6.10.6.4.1     | Substitution of Hydrogen   | 337 |
| 17.4.6.10.6.4.1.1   | Method 1: N-Alkylation of 5 <i>H</i> -Dibenz[ <i>b,f</i> ]azepines                                       | 337 |
| 17.4.6.10.6.4.1.1.1 | Variation 1: By Phase-Transfer Catalysis   | 338 |
| 17.4.6.10.6.4.1.2   | Method 2: N-Acylation of 5 <i>H</i> -Dibenz[ <i>b,f</i> ]azepines  | 339 |
| 17.4.6.10.6.4.1.2.1 | Variation 1: By Reaction with Acid Chlorides   | 339 |
| 17.4.6.10.6.4.1.2.2 | Variation 2: By Reaction with Dimethyl Carbonate   | 340 |
| 17.4.6.10.6.4.1.2.3 | Variation 3: By Palladium-Mediated Carbonylative Benzoylation of 5 <i>H</i> -Dibenz[ <i>b,f</i> ]azepine | 340 |
| 17.4.6.10.6.4.1.2.4 | Variation 4: By Reaction with Trifluoroacetic Anhydride  | 341 |
| 17.4.6.10.6.4.1.2.5 | Variation 5: N-Formylation of 5 <i>H</i> -Dibenz[ <i>b,f</i> ]azepine                                    | 341 |
| 17.4.6.10.6.4.1.3   | Method 3: Chlorocarbonylation of 5 <i>H</i> -Dibenz[ <i>b,f</i> ]azepines with Phosgene Equivalents      | 342 |
| 17.4.6.10.6.4.1.3.1 | Variation 1: N-Acylation of 5 <i>H</i> -Dibenz[ <i>b,f</i> ]azepines Followed by Amination               | 342 |
| 17.4.6.10.6.4.1.4   | Method 4: Formation of N—P Bonds   | 342 |
| 17.4.6.10.6.4.1.5   | Method 5: By a Methoxy Group   | 344 |
| 17.4.6.10.6.4.1.6   | Method 6: Palladium-Catalyzed N-Arylation  | 345 |
| 17.4.6.10.6.4.2     | Substitution of Heteroatoms  | 346 |
| 17.4.6.10.6.4.2.1   | Method 1: Substitution of Bromine  | 346 |
| 17.4.6.10.6.4.3     | Addition Reactions   | 347 |
| 17.4.6.10.6.4.3.1   | Method 1: Formation of Epoxides by Oxidation   | 347 |
| 17.4.6.10.6.4.3.2   | Method 2: Formation of Diols by Oxidation  | 347 |
| 17.4.6.10.6.4.3.3   | Method 3: [2 + 2] Photodimerization of 5-Acetyl-5 <i>H</i> -dibenz[ <i>b,f</i> ]azepine                  | 348 |
| 17.4.6.10.6.4.3.3.1 | Variation 1: [2 + 2] Photodimerization of 5 <i>H</i> -Dibenz[ <i>b,f</i> ]azepine Derivatives            | 348 |
| 17.4.6.10.7         | 9 <i>H</i> -Tribenz[ <i>b,d,f</i> ]azepines  | 349 |
| 17.4.6.10.7.1       | Synthesis by Ring-Closure Reactions  | 349 |
| 17.4.6.10.7.1.1     | By Formation of One N—C Bond   | 349 |
| 17.4.6.10.7.1.1.1   | Method 1: From <i>N</i> -(2-Bromophenyl)biphenyl-2-amine   | 349 |
| 17.4.6.10.8         | Other Group 15 Benzoheteropins   | 350 |
| 17.4.6.10.8.1       | 3-Benzoheteropins  | 350 |
| 17.4.6.10.8.1.1     | Synthesis by Ring-Closure Reactions  | 350 |
| 17.4.6.10.8.1.1.1   | By Formation of Two Heteroatom—Carbon Bonds  | 350 |
| 17.4.6.10.8.1.1.1.1 | Method 1: Potassium Hydroxide Catalyzed Addition of Metal Complexed Phosphines to 1,2-Diethynylbenzene   | 350 |
| 17.4.6.10.8.1.1.1.2 | Method 2: From 1,2-Bis[( <i>Z</i> )-2-bromovinyl]benzenes and Metal Halides                              | 351 |

## Volume 19: Three Carbon—Heteroatom Bonds: Nitriles, Isocyanides, and Derivatives

|               |  |       |     |
|---------------|--|-------|-----|
| 19.5          | <b>Product Class 5: Nitriles</b>   |       |     |
| 19.5.17       | <b>Synthesis of Nitriles Using Cross-Coupling Reactions</b>                    |       | New |
|               | D. M. Rudzinski and N. E. Leadbeater   |       |     |
| 19.5.17       | <b>Synthesis of Nitriles Using Cross-Coupling Reactions</b>                    | ..... | 355 |
| 19.5.17.1     | Preparation of Aryl Cyanides   | ..... | 355 |
| 19.5.17.1.1   | Method 1: Use of Alkali Metal Cyanides   | ..... | 355 |
| 19.5.17.1.1.1 | Variation 1: Palladium-Catalyzed Approaches                                    | ..... | 356 |
| 19.5.17.1.1.2 | Variation 2: Copper-Catalyzed Approaches                                       | ..... | 358 |
| 19.5.17.1.1.3 | Variation 3: Dual Palladium- and Copper-Catalyzed Approaches                   | ..... | 359 |
| 19.5.17.1.1.4 | Variation 4: Nickel-Catalyzed Approaches                                       | ..... | 360 |
| 19.5.17.1.2   | Method 2: Use of Zinc(II) Cyanide  | ..... | 361 |
| 19.5.17.1.2.1 | Variation 1: Palladium-Catalyzed Approaches: Homogeneous Catalysis             | ..    | 361 |
| 19.5.17.1.2.2 | Variation 2: Palladium-Catalyzed Approaches: Heterogeneous Catalysis           | ..    | 369 |
| 19.5.17.1.2.3 | Variation 3: Copper-Mediated Approaches  | ..... | 371 |
| 19.5.17.1.3   | Method 3: Use of Nickel(II) Cyanide as the Cyanide Source                      | ..... | 372 |
| 19.5.17.1.4   | Method 4: Use of Copper(I) Cyanide as the Cyanide Source                       | ..... | 373 |
| 19.5.17.1.5   | Method 5: Use of Potassium Hexacyanoferrate(II) as the Cyanide Source          | ·     | 374 |
| 19.5.17.1.5.1 | Variation 1: Palladium-Catalyzed Approaches: Homogeneous Catalysis             | ..    | 374 |
| 19.5.17.1.5.2 | Variation 2: Palladium-Catalyzed Approaches: Heterogeneous Catalysis           | ..    | 381 |
| 19.5.17.1.5.3 | Variation 3: Copper-Catalyzed Approaches                                       | ..... | 384 |
| 19.5.17.1.5.4 | Variation 4: Dual Palladium- and Copper-Catalyzed Approaches                   | ..... | 387 |
| 19.5.17.1.6   | Method 6: Use of Organic Cyanide Sources                                       | ..... | 388 |
| 19.5.17.1.6.1 | Variation 1: Use of Cyanohydrins   | ..... | 388 |
| 19.5.17.1.6.2 | Variation 2: Use of Trimethylsilyl Cyanide                                     | ..... | 392 |
| 19.5.17.1.6.3 | Variation 3: Use of <i>N</i> -Cyano- <i>N</i> -phenyl-4-toluenesulfonamide     | ..... | 394 |
| 19.5.17.1.7   | Method 7: In Situ Generation of Cyanide from Non-Cyanide-Containing Precursors | ..... | 394 |
| 19.5.17.2     | Preparation of Hetaryl Cyanides  | ..... | 396 |
| 19.5.17.2.1   | Method 1: Palladium-Catalyzed Coupling Reactions                               | ..... | 400 |
| 19.5.17.2.2   | Method 2: Copper-Catalyzed Coupling Reactions                                  | ..... | 403 |
| 19.5.17.2.3   | Method 3: Dual Palladium- and Copper-Catalyzed Approaches                      | ..... | 406 |
| 19.5.17.3     | Preparation of Vinyl Cyanides  | ..... | 409 |
| 19.5.17.3.1   | Method 1: Cyanation of Vinyl Halides and Vinylboronic Acids                    | ..... | 409 |

## Volume 27: Heteroatom Analogues of Aldehydes and Ketones

|               |  |            |
|---------------|--|------------|
| 27.25         | <b>Product Class 25: <i>N</i>-Sulfanyl-, <i>N</i>-Selanyl-, and <i>N</i>-Tellanylimines, and Their Oxidation Derivatives</b> | <b>New</b> |
|               | F. Chemla, F. Ferreira, and A. Pérez-Luna  |            |
| 27.25         | <b>Product Class 25: <i>N</i>-Sulfanyl-, <i>N</i>-Selanyl-, and <i>N</i>-Tellanylimines, and Their Oxidation Derivatives</b> | 417        |
| 27.25.1       | <b>Product Subclass 1: <i>N</i>-Sulfanylimines</b>   | 417        |
| 27.25.1.1     | Synthesis of Product Subclass 1  | 417        |
| 27.25.1.1.1   | Method 1: Synthesis from Aldehydes or Ketones  | 417        |
| 27.25.1.1.1.1 | Variation 1: From Ammonia and a Thiol  | 417        |
| 27.25.1.1.1.2 | Variation 2: From Ammonia and a Disulfide  | 418        |
| 27.25.1.1.1.3 | Variation 3: From <i>N,N</i> -Bis(trimethylsilyl)sulfenamides  | 419        |
| 27.25.1.1.1.4 | Variation 4: From Sulfenamides   | 421        |
| 27.25.1.1.2   | Method 2: Synthesis from Imines and Imine Derivatives  | 424        |
| 27.25.1.1.2.1 | Variation 1: From <i>N</i> -Unsubstituted Imines   | 424        |
| 27.25.1.1.2.2 | Variation 2: From Oximes   | 425        |
| 27.25.1.1.2.3 | Variation 3: From Oxime Thiocarbamates   | 426        |
| 27.25.1.1.2.4 | Variation 4: From <i>O</i> -Tosyloximes  | 427        |
| 27.25.1.1.2.5 | Variation 5: From <i>N</i> -Chloroimines   | 428        |
| 27.25.1.1.3   | Method 3: Synthesis from $\alpha$ -Aminoalkanoates   | 430        |
| 27.25.1.1.4   | Method 4: Synthesis from Sulfinamides  | 431        |
| 27.25.1.1.5   | Method 5: Synthesis from Nitro Compounds   | 431        |
| 27.25.2       | <b>Product Subclass 2: <i>N</i>-Sulfinylimines</b>   | 432        |
| 27.25.2.1     | Synthesis of Product Subclass 2  | 432        |
| 27.25.2.1.1   | Method 1: Synthesis by Oxidation of <i>N</i> -Sulfanylimines   | 432        |
| 27.25.2.1.2   | Method 2: Synthesis from <i>N</i> -Metalated Imines  | 436        |
| 27.25.2.1.2.1 | Variation 1: From Sulfinate Esters   | 436        |
| 27.25.2.1.2.2 | Variation 2: From a Cyclic Sulfinamide   | 438        |
| 27.25.2.1.3   | Method 3: Synthesis from Ortho Esters  | 439        |
| 27.25.2.1.4   | Method 4: Synthesis from an Aldehyde Hydrate   | 440        |
| 27.25.2.1.5   | Method 5: Synthesis from Carbonyl Derivatives  | 440        |
| 27.25.2.1.5.1 | Variation 1: From 1,2,3-Oxathiazolidine 2-Oxides   | 440        |
| 27.25.2.1.5.2 | Variation 2: From Sulfinate Esters   | 442        |
| 27.25.2.1.5.3 | Variation 3: From an <i>N</i> -Sulfinylbornane-10,2-sultam   | 444        |
| 27.25.2.1.5.4 | Variation 4: From Sulfinamides   | 445        |
| 27.25.2.1.5.5 | Variation 5: From a Phosphazene  | 448        |
| 27.25.2.1.6   | Method 6: Synthesis from Sulfoximides  | 449        |
| 27.25.2.1.7   | Method 7: Synthesis from Other <i>N</i> -Sulfinylimines  | 450        |
| 27.25.2.2     | Applications of Product Subclass 2 in Organic Synthesis  | 451        |



|               |   |     |
|---------------|---|-----|
| 27.25.3       | <b>Product Subclass 3: <i>N</i>-Sulfonylimines</b> .....                        | 453 |
| 27.25.3.1     | Synthesis of Products of Subclass 3 .....                                       | 453 |
| 27.25.3.1.1   | Method 1: Synthesis from Acetals .....  | 453 |
| 27.25.3.1.1.1 | Variation 1: From <i>O,O</i> -Acetals and Sulfonamides .....                    | 453 |
| 27.25.3.1.1.2 | Variation 2: From <i>N</i> -[(Arylsulfonyl)methyl]arenesulfonamides .....       | 454 |
| 27.25.3.1.2   | Method 2: Synthesis from Alkenes and Allenes .....                              | 455 |
| 27.25.3.1.2.1 | Variation 1: By Oxidative Amination .....                                       | 455 |
| 27.25.3.1.2.2 | Variation 2: From <i>N,N</i> -Dihalosulfonamides .....                          | 456 |
| 27.25.3.1.2.3 | Variation 3: From Sulfonyl Azides .....   | 456 |
| 27.25.3.1.2.4 | Variation 4: From Oxazolidinones .....  | 458 |
| 27.25.3.1.3   | Method 3: Synthesis from Alkynes .....  | 459 |
| 27.25.3.1.3.1 | Variation 1: By a Hydroamination Reaction .....                                 | 459 |
| 27.25.3.1.3.2 | Variation 2: From Sulfonyl Azides .....   | 461 |
| 27.25.3.1.3.3 | Variation 3: Through Iminobismuthane Addition .....                             | 462 |
| 27.25.3.1.4   | Method 4: Synthesis from Aziridines .....                                       | 463 |
| 27.25.3.1.5   | Method 5: Synthesis from Carbonyl Compounds .....                               | 464 |
| 27.25.3.1.5.1 | Variation 1: From Sulfonamides .....  | 464 |
| 27.25.3.1.5.2 | Variation 2: From Isocyanates .....   | 469 |
| 27.25.3.1.5.3 | Variation 3: From <i>N</i> -Sulfinylsulfonamides .....                          | 470 |
| 27.25.3.1.5.4 | Variation 4: From Chloramine-T .....  | 471 |
| 27.25.3.1.6   | Method 6: Synthesis from Other Imines .....                                     | 473 |
| 27.25.3.1.6.1 | Variation 1: From Oximes .....  | 473 |
| 27.25.3.1.6.2 | Variation 2: From <i>N</i> -(Trimethylsilyl)imines .....                        | 475 |
| 27.25.3.1.6.3 | Variation 3: From <i>N</i> -Sulfanylimines .....                                | 475 |
| 27.25.3.1.6.4 | Variation 4: From <i>N</i> -Sulfinylimines .....                                | 476 |
| 27.25.3.1.6.5 | Variation 5: From Imidoyl Chlorides .....                                       | 477 |
| 27.25.3.1.7   | Method 7: Synthesis from Sulfinimides .....                                     | 478 |
| 27.25.3.1.8   | Method 8: Synthesis by Oxidation of <i>N</i> -Sulfonylanilines .....            | 480 |
| 27.25.3.1.9   | Method 9: Synthesis from <i>N</i> -Sulfonylamides .....                         | 487 |
| 27.25.3.2     | Applications of Product Subclass 3 in Organic Synthesis .....                   | 487 |
| 27.25.4       | <b>Product Subclass 4: <i>N</i>-Selanylimines</b> .....                         | 488 |
| 27.25.4.1     | Synthesis of Product Subclass 4 .....   | 488 |
| 27.25.4.1.1   | Method 1: Synthesis from <i>N</i> -Unsubstituted Imines .....                   | 488 |
| 27.25.4.1.2   | Method 2: Synthesis by Oxidation of Phenols .....                               | 488 |
| 27.25.4.1.3   | Method 3: Synthesis from <i>N</i> -Selenamides .....                            | 490 |
| 27.25.5       | <b>Product Subclass 5: <i>N</i>-Seleninylimines and Related Compounds</b> ..... | 490 |
| 27.25.5.1     | Synthesis of Product Subclass 5 .....   | 490 |
| 27.25.5.1.1   | Method 1: Synthesis from Imines and Imine Derivatives .....                     | 490 |
| 27.25.5.1.1.1 | Variation 1: From Imines .....  | 491 |
| 27.25.5.1.1.2 | Variation 2: From <i>N</i> -Chloroimines .....                                  | 491 |
| 27.25.6       | <b>Product Subclass 6: <i>N</i>-Tellanylimines</b> .....                        | 491 |
| 27.25.6.1     | Synthesis of Product Subclass 6 .....   | 491 |

|             |  |     |
|-------------|--|-----|
| 27.25.6.1.1 | Method 1: From <i>N</i> -Metalloimines .....         | 491 |
| 27.25.6.1.2 | Method 2: From Pentafluorotellurium Isocyanate ..... | 492 |
|             | <b>Author Index</b> .....                            | 501 |
|             | <b>Abbreviations</b> .....                           | 523 |