

Dipl. oec. troph. Franca Mangiameli  
Prof. Dr. Nicolai Worm

# Außen schlank – innen fett

## Literaturverzeichnis

### Vorwort

Sharma P, Kumar A. Metabolic dysfunction associated fatty liver disease increases risk of severe Covid-19  
Diabetes Metab Syndr. 2020;14(5):825-827.

### VERSTECKTE FETTDEPOTS MACHEN KRANK

#### Fettpolster – früher überlebenswichtig

Freese J, Klement RJ, Ruiz-Núñez B. The sedentary (r)evolution: Have we lost our metabolic flexibility?  
F1000Res. 2017;6:1787.

Magkos F. Metabolically healthy obesity: what's in a name? Am J Clin Nutr 2019; 00:1–7.

<https://www.ifb-adipositas.de/adipositas/entwicklungen> (letzter Zugriff: 12.08.2020)

The Global BMI Mortality Collaboration. Body-mass index and all-cause mortality: individual- participant-data meta-analysis of 239 prospective studies in four continents. Lancet 2016; 388: 776–86

<http://www.euro.who.int/de/health-topics/noncommunicable-diseases/obesity/news/news/2017/10/world-obesity-day-understanding-the-social-consequences-of-obesity> (letzter Zugriff: 12.08.2020)

Smith GI, Mittendorfer B, Klein S. Metabolically healthy obesity: facts and fantasies. J Clin Invest. 2019; 129(10): 3978–3989.

Echouffo-Tcheugui JB, Short MI, Xanthakos V et al. Natural history of obesity subphenotypes: dynamic changes over two decades and prognosis in the framingham heart study. J Clin Endocrinol Metab 2019; 104(3): 738–752.

Ortega FB, Lee DC, Katzmarzyk PT et al. The intriguing metabolically healthy but obese phenotype: cardiovascular prognosis and role of fitness. Eur Heart J 2013; 34(5): 389–97.

De Lorenzo A, Gratteri S, Gualtieri P et al. Why primary obesity is a disease? J Transl Med 2019; 17:169.

Iacobini C, Pugliese G, Blasetti Fantauzzi C et al. Metabolically healthy versus metabolically unhealthy obesity. Metabolism 2019; 92: 51-60.

Lin H, Zhang L; Zheng R et al. The prevalence, metabolic risk and effects of lifestyle intervention for metabolically healthy obesity, Medicine 2017; 96(47): e8838.

Matsuzawa J. Pathophysiology and molecular mechanisms of visceral fat syndrome: the Japanese experience. Diabetes Metab Rev. 1997; 13(1): 3-13.

Magkos F. Metabolically healthy obesity: what's in a name? Am J Clin Nutr 2019; 00:1–7.

## Dünne im Vorteil – schlank, gesund?

The Global BMI Mortality Collaboration. Body-mass index and all-cause mortality: individual- participant-data meta-analysis of 239 prospective studies in four continents. Lancet 2016; 388: 776–86

Eckel N, Li Y, Kuxhaus O et al. Transition from metabolic healthy to unhealthy phenotypes and association with cardiovascular disease risk across BMI categories in 90 257 women (the Nurses' Health Study): 30 year follow-up from a prospective cohort study. Lancet Diabetes Endocrinol 2018; 6(9): 714-724.

Kramer CK, Zinman B, Retnakaran R. Are metabolically healthy overweight and obesity benign conditions? A systematic review and meta-analysis. Ann Intern Med. 2013; 159: 758-769.

Sahakyan KR, Somers VK, JP Rodriguez-Escudero et al. Normal-weight central obesity: Implications for total and cardiovascular mortality. Ann Intern Med 2015; 163 (11): 827-835.

Bosomworth NJ. Normal-weight central obesity. Unique hazard of the toxic waist. Can Fam Physician 2019; 65(6): 399-408.

Coutinho T, Goel K, Corrêa de Sá D et al. Combining body mass index with measures of central obesity in the assessment of mortality in subjects with coronary disease: role of "normal weight central obesity". J Am Coll Cardiol 2013; 61(5): 553-60.

Thomas EL, Frost G, Taylor-Robinson SD et al. Excess body fat in obese and normal-weight subjects. Nutr Res Rev. 2012 Jun; 25(1): 150-61.

Kennedy AP, Shea JL, Sun G. Comparison of the classification of obesity by BMI vs. dual-energy X-ray absorptiometry in the Newfoundland population. Obesity (Silver Spring). 2009;17(11):2094-2099.

De Lorenzo A, Deurenberg P, Pietrantuono et al. How fat is obese? Acta Diabetol 2003; 40: 254–257.

Tomiyama AJ, Hunger JM, Nguyen-Cuu J et al. Misclassification of cardiometabolic health when using body mass index categories in NHANES 2005–2012. Int J Obes 2016; 40: 883–886.

Männistö, S, Harald, K, Kontto, J et al. Dietary and lifestyle characteristics associated with normal-weight obesity: The National FINRISK 2007 Study. 2014 British Journal of Nutrition, 111(5), 887-894.

Ohlsson B, Manjer J. Sociodemographic and Lifestyle Factors in relation to Overweight Defined by BMI and "Normal-Weight Obesity". J Obes. 2020; 2020:2070297.

## TOFI - thin outside and fat inside

Gujral UP, Vittinghoff E, Mongraw-Chaffin M et al. Cardiometabolic abnormalities among normal-weight persons from five racial/ethnic groups in the united states: A cross-sectional analysis of two cohort studies. Ann Intern Med 2017; 166(9): 628-636.

Mathew H, Farr OM, Mantzoros CS. Metabolic health and weight: Understanding metabolically unhealthy normal weight or metabolically healthy obese patients. Metabolism. 2016;65(1):73-80.

Stefan N, Schick F, Häring HU. Causes, characteristics, and consequences of metabolically unhealthy normal weight in humans. Cell Metab 2017; 26(2): 292-300.

Buscemi S, Chiarello P, Buscemi C, et al. Characterization of Metabolically Healthy Obese People and Metabolically Unhealthy Normal-Weight People in a General Population Cohort of the ABCD Study. J Diabetes Res. 2017;2017:9294038.

Stefan N. Causes, consequences, and treatment of metabolically unhealthy fat distribution. Lancet Diabetes Endocrinol 2020; 8: 616–27.

Choi SI, Chung D, Lim JS et al. Relationship between regional body fat distribution and diabetes mellitus: 2008 to 2010 Korean National Health and Nutrition Examination Surveys. *Diabetes Metab J* 2017; 41(1): 51-59.

Vasan SK, Osmond C, Canoy D et al. Comparison of regional fat measurements by dual-energy X-ray absorptiometry and conventional anthropometry and their association with markers of diabetes and cardiovascular disease risk. *Int J Obes* 2018; 42(4): 850-857.

Klitgaard, HB, Kilbak, JH, Nozawa, EA. et al. Physiological and Lifestyle Traits of Metabolic Dysfunction in the Absence of Obesity. *Curr Diab Rep* 2020, 20, 17.

Kuzik N, Carson V, Andersen LB et al. Physical activity and sedentary time associations with metabolic health across weight statuses in children and adolescents. *Obesity* 2017; 25: 1762-1769.

Nier A, Brandt A, Baumann A et al. Metabolic abnormalities in normalweight children are associated with increased visceral fat accumulation, elevated plasma endotoxin levels and a higher monosaccharide intake. *Nutrients* 2019; 11(3), 652.

Nemet D. Childhood obesity, physical activity, and exercise – the year that was 2017: normal-weight obese— are we missing a population in need? *Pediatr Exerc Sci* 2018; 30(1): 52-53.

Musálek M, Pařízková J, Godina E et al. Poor skeletal robustness on lower extremities and weak lean mass development on upper arm and calf: normal weight obesity in middle-school-aged children (9 to 12). *Front Pediatr* 2018; 6: 371.

Viitasalo A, Pitkänen N, Pahkala K. et al. Increase in adiposity from childhood to adulthood predicts a metabolically obese phenotype in normal-weight adults. *Int J Obes* 2020; 44, 848–851.

Van Hulst A, Ybarra M, Mathieu ME et al. Determinants of new onset cardiometabolic risk among normal weight children. *Int J Obes (Lond)*. 2020;44(4):781-789.

Bhargava SK, Sachdev HS, Fall CH, et al. Relation of serial changes in childhood body-mass index to impaired glucose tolerance in young adulthood. *N Engl J Med*. 2004;350(9):865-875.

Mokha JS, Srinivasan SR, Dasmahapatra P, et al. Utility of waist-to-height ratio in assessing the status of central obesity and related cardiometabolic risk profile among normal weight and overweight/obese children: the Bogalusa Heart Study. *BMC Pediatr*. 2010;10:73. Published 2010 Oct 11.

## Fettgewebe als Schutzfaktor

Blüher M, Laufs U. New concepts for body shape-related cardiovascular risk: role of fat distribution and adipose tissue function. European Heart J 2019; 40, 2856–2858.

Blüher M. Inflammation des Fettgewebes. Diabetologe 2019; 15, 296–304.

Bays H. Central obesity as a clinical marker of adiposopathy; increased visceral adiposity as a surrogate marker for global fat dysfunction. Curr Opin Endocrinol Diabetes Obes 2014, 21:345–351.

Kahn CR, Wang G, Lee KY. Altered adipose tissue and adipocyte function in the pathogenesis of metabolic syndrome. J Clin Invest. 2019; 129(10): 3990-4000.

Klöting N, Blüher M. Adipocyte dysfunction, inflammation and metabolic syndrome. Rev Endocr Metab Disord, 2014.

Neeland IJ, Ross R, Després JP. Visceral and ectopic fat, atherosclerosis, and cardiometabolic disease: a position statement. Lancet Diabetes Endocrinol 2019; 7(9): 715-725.

Moreno-Indias I, Oliva-Olivera W, Omiste A et al. Adipose tissue infiltration in normal-weight subjects and its impact on metabolic function. Transl Res 2016; 172:6-17.e3.

Carrobbio S, Pellegrinelli V, Vidal-Puig A. Adipose Tissue Function and Expandability as Determinants of Lipotoxicity and the Metabolic Syndrome. Engin A., Engin A. (eds) Obesity and Lipotoxicity. Advances in Experimental Medicine and Biology, 2020, vol 960. Springer.

Kang S, Kyung C, Park JS et al. Subclinical vascular inflammation in subjects with normal weight obesity and its association with body Fat: an 18 F-FDG-PET/CT study. Cardiovasc Diabetol 2014; 13, 70.

Kim S, Kyung C, Park JS. et al. Normal-weight obesity is associated with increased risk of subclinical atherosclerosis. Cardiovasc Diabetol 2015; 14, 58 (2015).

Bosy-Westphal A, Braun W, Albrecht V. Determinants of ectopic liver fat in metabolic disease EurJ Clin Nutr 2019;73:209-214.

Iyengar NM, Arthur R, Manson JE et al. Association of Body Fat and Risk of Breast Cancer in Postmenopausal Women With Normal Body Mass Index: A Secondary Analysis of a Randomized Clinical Trial and Observational Study. *JAMA Oncol.* 2019 Feb 1;5(2):155-163.

Ross R, Neeland I, Shizuya Y et al. Waist circumference as a vital sign in clinical practice: A consensus statement from the IAS and ICCR Working Group on visceral obesity. *Nat Rev Endocrinol.* 2020; 16(3):177-189.

Brøns C, Groth Grunnet L. Skeletal muscle lipotoxicity in insulin resistance and type 2 diabetes: a causal mechanism or an innocent bystander? *European Journal of Endocrinology*, 2017; 176, R67–R78.

## RAUS AUS DER TOFI-FALLE

### Was macht den TOFI zum TOFI?

Tan CY, Vidal-Puig A. Adipose tissue expandability: The metabolic problems of obesity may arise from the inability to become more obese. *Biochem Soc Trans.* 2008;36(Pt 5):935-940.

Osadnik, K, Osadnik T, Lonnie, M. et al. Metabolically healthy obese and metabolic syndrome of the lean: the importance of diet quality. Analysis of MAGNETIC cohort. *Nutr J* 19, 19 (2020).

Brandão, I, Martins, MJ, Monteiro, R. Metabolically Healthy Obesity—Heterogeneity in Definitions and Unconventional Factors. *Metabolites* 2020, 10, 48.

Tchernof A, Després JP. Pathophysiology of human visceral obesity: an update. *Physiol Rev.* 2013;93(1):359-404.

Paniagua JA. Nutrition, insulin resistance and dysfunctional adipose tissue determine the different components of metabolic syndrome. *World J Diabetes.* 2016;7(19).

Lotta L, Gulati P, Day F et al. Integrative genomic analysis implicates limited peripheral adipose storage capacity in the pathogenesis of human insulin resistance. *Nat Genet* 49, 17–26 (2017).

Scott RA, Fall T, Pasko. D Common Genetic Variants Highlight the Role of Insulin Resistance and Body Fat Distribution in Type 2 Diabetes, Independent of Obesity. *Diabetes* 2014, 63 (12) 4378-4387.

Stephensa BR, Granadosa K, Zdericb, TW. Effects of 1 day of inactivity on insulin action in healthy men and women: interaction with energy intake. *Metabolism Clinical and Experimental* 60 (2011) 941–949.

Dirks ML, Wall BT, van de Valk B, et al. One Week of Bed Rest Leads to Substantial Muscle Atrophy and Induces Whole-Body Insulin Resistance in the Absence of Skeletal Muscle Lipid Accumulation. *Diabetes*. 2016;65(10):2862-2875.

Stephens BR, Granados K, Zderic TW et al. Effects of 1 day of inactivity on insulin action in healthy men and women: interaction with energy intake. *Metabolism*. 2011;60(7):941-949.

Olsen RH, Krogh-Madsen R, Thomsen C. et al. Metabolic responses to reduced daily steps in healthy nonexercising men. *JAMA*. 2008;299(11):1261-1263.

Bowden Davies KA, Sprung VS, Norman JA, et al. Short-term decreased physical activity with increased sedentary behaviour causes metabolic derangements and altered body composition: effects in individuals with and without a first-degree relative with type 2 diabetes. *Diabetologia*. 2018;61(6):1282-1294.

Bowden Davies KA, Pickles S, Sprung VS, et al. Reduced physical activity in young and older adults: metabolic and musculoskeletal implications. *Ther Adv Endocrinol Metab*. 2019; 10:2042018819888824.

Wolfe R. The underappreciated role of muscle in health and disease. *Am J Clin Nutr* 2006; 84:475–82.

Seo YG, Song HJ, Song YR. Fat-to-muscle ratio as a predictor of insulin resistance and metabolic syndrome in Korean adults. *J Cachexia Sarcopenia Muscle*. 2020;11(3):710-725.

Kim K, Park SM. Association of muscle mass and fat mass with insulin resistance and the prevalence of metabolic syndrome in Korean adults: a cross-sectional study. *Sci Rep*. 2018; 8(1):2703.

Polyzos, S, Margioris, AN. Sarcopenic obesity. *Hormones* 2018, 17, 321–331.

Kim, K., Park, S.M. Association of muscle mass and fat mass with insulin resistance and the prevalence of metabolic syndrome in Korean adults: a cross-sectional study. *Sci Rep* 2018, 8, 2703.

Choi KM. Sarcopenia and sarcopenic obesity. *Korean J Intern Med*. 2016;31(6):1054-1060.

Stefan N, Schick F, Häring HU. Causes, characteristics, and consequences of metabolically unhealthy normal weight in humans. *Cell Metab* 2017; 26(2): 292-300.

Burrows, R, Correa, P, Reyes, M. et al. Low muscle mass is associated with cardiometabolic risk regardless of nutritional status in adolescents: A cross - sectional study in a Chilean birth cohort, *Pediatr Diabetes*, 2017; 18: 895– 902.

Mirmiran P, Moslehi N, Hosseinpanah F et al. Dietary determinants of unhealthy metabolic phenotype in normal weight and overweight/obese adults: results of a prospective study. *Int. J. Food Sci. Nutr* 2020 Apr 1;1-11.

Monteiro C, Moubarac JC, Bertazzi R et al. Household availability of ultra-processed foods and obesity in nineteen European countries, *Publ Health Nutr* 2018;21:18-26.

Ghanim H, Abuaysheh S, Sia CL et al. Increase in plasma endotoxin concentrations and the expression of Toll-like receptors and suppressor of cytokine signaling-3 in mononuclear cells after a high-fat, high-carbohydrate meal: implications for insulin resistance. *Diabetes Care*. 2009 Dec;32(12):2281-7.

Barr SB, Wright JC. Postprandial energy expenditure in whole-food and processed-food meals: implications for daily energy expenditure. *Food Nutr Res*. 2010; 54:10.3402/fnr.v54i0.5144.

Zinöcker MK, Lindseth IA. The Western Diet-Microbiome-Host Interaction and Its Role in Metabolic Disease. *Nutrients*. 2018;10(3):365.

Guerrero-Romero F, Rodriguez-Moran M. Serum magnesium in the metabolically-obese normal-weight and healthy-obese subjects. *Eur J Intern Med*. 2013; 24(7):639-643.

Wang X, Chang X, Zhu Y et al. Metabolically Obese Individuals of Normal Weight Have a High Risk of 25-Hydroxyvitamin D Deficiency. *Am J Med Sci*. 2016;352(4):360-367.

Wolfe MM, Boylan MO. Obesity and the gastrointestinal tract: you are what you eat. *J Clin Gastroenterol*. 2014;48(10):817-822.

Olechnowicz J, Tinkov A, Skalny A et al. Zinc status is associated with inflammation, oxidative stress, lipid, and glucose metabolism. *J Physiol Sci*. 2018; 68(1):19-31.

Yang HK, Lee SH, Han K, et al. Lower serum zinc levels are associated with unhealthy metabolic status in normal-weight adults: The 2010 Korea National Health and Nutrition Examination Survey. *Diabetes Metab.* 2015;41(4):282-290.

Wang Z, Wang D, Wang Y. Cigarette Smoking and Adipose Tissue: The Emerging Role in Progression of Atherosclerosis. *Mediators Inflamm.* 2017; 2017:3102737.

Kato A, Li Y, Ota A, et al. Smoking results in accumulation of ectopic fat in the liver. *Diabetes Metab Syndr Obes.* 2019; 12:1075-1080.

Graff-Iversen S, Hewitt S, Forsén L et al. Associations of tobacco smoking with body mass distribution; a population-based study of 65,875 men and women in midlife. *BMC Public Health.* 2019;19(1):1439.

Nakano Y. Adult-Onset Diseases in Low Birth Weight Infants: Association with Adipose Tissue Maldevelopment. *J Atheroscler Thromb.* 2020; 27(5):397-405.

Hovi P, Andersson S, Eriksson JG, et al. Glucose regulation in young adults with very low birth weight. *N Engl J Med.* 2007; 356(20):2053-2063.

Kuk JL, Saunders TJ, Davidson LE et al. Age-related changes in total and regional fat distribution. *Ageing Res Rev.* 2009; 8(4):339-348.

Ambikairajah A, Walsh E, Tabatabaei-Jafari H et al. Fat mass changes during menopause: a metaanalysis. *Am J Obstet Gynecol.* 2019;221(5):393-409.e50.

Dulloo AG, Jacquet J, Montani JP et al. How dieting makes the lean fatter: from a perspective of body composition autoregulation through adipostats and proteinstats awaiting discovery. *Obes Rev.* 2015; 16 Suppl 1:25-35.

Krsek M, Rosická M, Nedvídková J, et al. Increased lipolysis of subcutaneous abdominal adipose tissue and altered noradrenergic activity in patients with Cushing's syndrome: an in-vivo microdialysis study. *Physiol Res.* 2006;55(4):421-428.

Epel ES. Psychological and metabolic stress: a recipe for accelerated cellular aging? *Hormones (Athens).* 2009; 8(1):7-22.

## Bin ich ein TOFI?

Ding C, Chan Z, Magkos F. Lean, but not healthy: the 'metabolically obese, normal-weight' phenotype. *Curr Opin Clin Nutr Metab Care.* 2016; 19(6):408-417.

Stefan N, Schick F, Häring HU. Causes, characteristics, and consequences of metabolically unhealthy normal weight in humans. *Cell Metab* 2017; 26(2): 292-300.

Ding C, Chan Z, Chooi YC, et al. Regulation of glucose metabolism in nondiabetic, metabolically obese normal-weight Asians. *Am J Physiol Endocrinol Metab.* 2018; 314(5):E494-E502.

Lu YC, Lin YC, Yen AM et al. Dual-energy X-ray absorptiometry-assessed adipose tissues in metabolically unhealthy normal weight Asians. *Sci Rep.* 2019;9(1):17698.

Ströhle A, Worm N (2014): Gesundes Übergewicht!? Warum das Übergewichts-Paradox nur scheinbar paradox ist. *Med Monatsschr Pharm* 37 (2), S. 54-64.

Thomas EL, Frost G, Taylor-Robinson SD et al. Excess body fat in obese and normal-weight subjects. *Nutr Res Rev.* 2012 Jun; 25(1): 150-61.

Benites-Zapata VA, Toro-Huamanchumo CJ, Urrunaga-Pastor D et al. High waist-to-hip ratio levels are associated with insulin resistance markers in normal-weight women. *Diabetes Metab Syndr.* 2019;13(1):636-642.

Oliveros E, Somers VK, Sochor O et al. The concept of normal weight obesity. *Prog Cardiovasc Dis.* 2014;56(4):426-433.

Moltz L, Holl RW. Primäre und sekundäre Insulinresistenz: Konzentrationen von Glukose und Insulin bei normgewichtigen, anscheinend gesunden Probandinnen. *Diabetologie und Stoffwechsel* 2010; 5(6): 372-378.

Morales-Gurrola G, Simental-Mendía LE, Castellanos-Juárez FX et al. The triglycerides and glucose index is associated with cardiovascular risk factors in metabolically obese normal-weight subjects. *J Endocrinol Invest.* 2020; 43(7):995-1000.

Ahn N, Baumeister SE, Amann U, et al. Visceral adiposity index (VAI), lipid accumulation product (LAP), and product of triglycerides and glucose (TyG) to discriminate prediabetes and diabetes. *Sci Rep.* 2019;9(1):9693.

Toro-Huamanchumo CJ, Urrunaga-Pastor D, Guarnizo-Poma M, et al. Triglycerides and glucose index as an insulin resistance marker in a sample of healthy adults. *Diabetes Metab Syndr.* 2019; 13(1):272-277.

Guerrero-Romero F, Villalobos-Molina R, Jiménez-Flores JR, et al. Fasting Triglycerides and Glucose Index as a Diagnostic Test for Insulin Resistance in Young Adults. *Arch Med Res.* 2016;47(5):382-387.

<https://www.mdapp.co/tyg-index-calculator-359/>

Kim JW, Kim DH, Roh YK, et al. Serum Ferritin Levels Are Positively Associated With Metabolically Obese Normal Weight: A Nationwide Population-Based Study. *Medicine (Baltimore).* 2015; 94(52):e2335.

Ahl S, Guenther M, Zhao S, et al. Adiponectin Levels Differentiate Metabolically Healthy vs Unhealthy Among Obese and Nonobese White Individuals. *J Clin Endocrinol Metab.* 2015;100(11):4172-4180.

Frühbeck G, Catalán V, Rodríguez A, et al. Adiponectin-leptin Ratio is a Functional Biomarker of Adipose Tissue Inflammation. *Nutrients.* 2019;11(2):454.

<https://www.leberfasten.com/leberfasten-konzept/machen-sie-den-test/> (letzter Zugriff: 12.08.2020)

Clarke SL, Reaven GM, Leonard D, et al. Cardiorespiratory Fitness, Body Mass Index, and Markers of Insulin Resistance in Apparently Healthy Women and Men. *Am J Med.* 2020;133(7):825-830.e2.

## Schlank aussehen und schlank sein

Zhang M, Schumann M, Huang T et al. Normal weight obesity and physical fitness in Chinese university students: an overlooked association. *BMC Public Health.* 2018; 18(1):1334.

Levine JA. Nonexercise activity thermogenesis--liberating the life-force. *J Intern Med.* 2007;262(3):273-287.

Villalblanca PA, Alegria JR, Mookadam F et al. Nonexercise activity thermogenesis in obesity management. *Mayo Clin Proc.* 2015; 90(4):509-519.

Lee DC, Pate RR, Lavie CJ et al. Leisure-time running reduces all-cause and cardiovascular mortality risk. *Am Coll Cardiol.* 2014;64(5):472-481.

Westcott WL. Resistance training is medicine: effects of strength training on health. *Curr Sports Med Rep.* 2012;11(4):209-216.

Ferreira FC, de Medeiros AI, Nicioli C, et al. Circuit resistance training in sedentary women: body composition and serum cytokine levels. *Appl Physiol Nutr Metab.* 2010;35(2):163-171.

Liu JX, Zhu L, Li PJ et al. Effectiveness of high-intensity interval training on glycemic control and cardiorespiratory fitness in patients with type 2 diabetes: a systematic review and meta-analysis. *Aging Clin Exp Res.* 2019;31(5):575-593.

Lora-Pozo I, Lucena-Anton D, Salazar A et al. Anthropometric, Cardiopulmonary and Metabolic Benefits of the High-Intensity Interval Training Versus Moderate, Low-Intensity or Control for Type 2 Diabetes: Systematic Review and Meta-Analysis. *Int J Environ Res Public Health.* 2019;16(22):4524.

Keech A, Holgate K, Fildes J, et al. High-intensity interval training for patients with coronary artery disease: Finding the optimal balance. *Int J Cardiol.* 2020; 298:8-14.

Lettieri-Barbato D, Giovannetti E, Aquilano K. Effects of dietary restriction on adipose mass and biomarkers of healthy aging in human. *Aging (Albany NY).* 2016;8(12):3341-3355.

Chaston TB, Dixon JB. Factors associated with percent change in visceral versus subcutaneous abdominal fat during weight loss: findings from a systematic review. *Int J Obes (Lond).* 2008;32(4):619-628.

McLaughlin T, Abbasi F, Lamendola C et al. Dietary weight loss in insulin-resistant non-obese humans: Metabolic benefits and relationship to adipose cell size. *Nutr Metab Cardiovasc Dis.* 2019; 29(1):62-68.

Chooi YC, Ding C, Chan Z, et al. Moderate Weight Loss Improves Body Composition and Metabolic Function in Metabolically Unhealthy Lean Subjects. *Obesity (Silver Spring).*

Cava E, Yeat NC, Mittendorfer B. Preserving Healthy Muscle during Weight Loss. *Adv Nutr.* 2017;8(3):511-519.

Worm N. Flexi-Carb: Mediterran genießen. Lebensstil beachten – Kohlenhydrate anpassen. Schlank und gesund bleiben. Riva Verlag, 2015.

Hyde PN, Sapper TN, Crabtree CD, et al. Dietary carbohydrate restriction improves metabolic syndrome independent of weight loss. *JCI Insight*. 2019; 4(12):e128308..

Kelly T, Unwin D, Finucane F. Low-Carbohydrate Diets in the Management of Obesity and Type 2 Diabetes: A Review from Clinicians Using the Approach in Practice. *Int J Environ Res Public Health*. 2020;17(7):2557.

Hallberg SJ, Gershuni VM, Hazbun TL et al. Reversing Type 2 Diabetes: A Narrative Review of the Evidence. *Nutrients*. 2019;11(4):766. Published 2019 Apr 1.

Gepner Y, Shelef I, Schwarzfuchs D, et al. Effect of Distinct Lifestyle Interventions on Mobilization of Fat Storage Pools: CENTRAL Magnetic Resonance Imaging Randomized Controlled Trial. *Circulation*. 2018;137(11):1143-1157.

Wildman R, Bruno RS. *Handbook of Nutraceuticals and Functional Foods (Modern Nutrition)*. CRC Press; Auflage: 3 (19. November 2019).

Haghishat, N, Ashtary-Larky, D, Bagheri, R et al. The effect of 12 weeks of equicaloric high protein diet in regulating appetite and body composition of women with Normal Weight Obesity: A randomized controlled trial. *2020 British Journal of Nutrition*, 1-20.

Choi J, Se-Young O, Lee D, et al. Characteristics of diet patterns in metabolically obese, normal weight adults (Korean National Health and Nutrition Examination Survey III, 2005). *Nutr Metab Cardiovasc Dis*. 2012;22(7):567-574.

Nishino K, Sakurai M, Takeshita Y, Takamura T. Consuming Carbohydrates after Meat or Vegetables Lowers Postprandial Excursions of Glucose and Insulin in Nondiabetic Subjects. *J Nutr Sci Vitaminol (Tokyo)*. 2018;64(5):316-320.

Johnston CS, Kim CM, Buller AJ. Vinegar improves insulin sensitivity to a high-carbohydrate meal in subjects with insulin resistance or type 2 diabetes. *Diabetes Care*. 2004;27(1):281-282.

Leeman M, Ostman E, Björck I. Vinegar dressing and cold storage of potatoes lowers postprandial glycaemic and insulinaemic responses in healthy subjects. *Eur J Clin Nutr*. 2005;59(11):1266-1271.

Shukla AP, Mauer E, Igel LI et al. Effect of Food Order on Ghrelin Suppression. *Diabetes Care*. 2018; 41(5):e76-e77.

Burton P, Lightowler HJ. The impact of freezing and toasting on the glycaemic response of white bread. Eur J Clin Nutr. 2008;62(5):594-599.

Paniagua JA, Gallego de la Sacristana GA, Romero I et al. Monounsaturated Fat–Rich Diet Prevents Central Body Fat Distribution and Decreases Postprandial Adiponectin Expression Induced by a Carbohydrate–Rich Diet in Insulin–Resistant Subjects Diabetes Care Jul 2007, 30 (7) 1717-1723;

Gao H, Geng T, Huang T, Zhao Q. Fish oil supplementation and insulin sensitivity: a systematic review and meta-analysis. Lipids Health Dis. 2017;16(1):131.

Spencer M, Finlin BS, Unal R, et al. Omega-3 fatty acids reduce adipose tissue macrophages in human subjects with insulin resistance. Diabetes. 2013;62(5):1709-1717.

Albracht-Schulte K, Kalupahana NS, Ramalingam L, et al. Omega-3 fatty acids in obesity and metabolic syndrome: a mechanistic update. J Nutr Biochem. 2018; 58:1-16.

von Schacky, C. Verwirrung um die Wirkung von Omega-3-Fettsäuren. Internist 2019, 60, 1319–1327.

Oshima Y, Rin S, Kita H, et al. The Frequency of Fish-Eating Could Negatively Associate with Visceral Adiposity in Those Who Eat Moderately. J Nutr Sci Vitaminol (Tokyo). 2015;61(5):426-431.

Sochol KM, Johns TS, Buttar RS, et al. The Effects of Dairy Intake on Insulin Resistance: A Systematic Review and Meta-Analysis of Randomized Clinical Trials. Nutrients. 2019;11(9):2237.

Bordoni A, Danesi F, Dardevet D et al. Dairy products and inflammation: A review of the clinical evidence. Crit Rev Food Sci Nutr. 2017;57(12):2497-2525.

Ulven SM, Holven KB, Gil A, Rangel-Huerta OD. Milk and Dairy Product Consumption and Inflammatory Biomarkers: An Updated Systematic Review of Randomized Clinical Trials. Adv Nutr. 2019;10(suppl\_2):S239-S250.

Guo J, Givens DI, Astrup A, et al. The Impact of Dairy Products in the Development of Type 2 Diabetes: Where Does the Evidence Stand in 2019? Adv Nutr. 2019;10(6):1066-1075.

Di Renzo L, Rizzo M, Sarlo F, et al. Effects of dark chocolate in a population of normal weight obese women: a pilot study. Eur Rev Med Pharmacol Sci. 2013;17(16):2257-2266.

Zugravu C, Otelea MR. Dark Chocolate: To Eat or Not to Eat? A Review. J AOAC Int. 2019;102(5):1388-1396.

Alperet DJ, Rebello SA, Khoo EY, et al. A randomized placebo-controlled trial of the effect of coffee consumption on insulin sensitivity: Design and baseline characteristics of the Coffee for METabolic Health (COMETH) study. Contemp Clin Trials Commun. 2016;4:105-117.

Ma J, McKeown NM, Hwang SJ et al. Sugar-sweetened beverage consumption is associated with change of visceral adipose tissue over 6 years of follow-up. Circulation. 2016; 133(4):370-377.

Ma J, Fox CS, Jacques PF et al. Sugar-sweetened beverage, diet soda, and fatty liver disease in the Framingham Heart Study cohorts. J Hepatol. 2015;63(2):462-469.

Jensen PN, Howard BV, Best LG, et al. Associations of diet soda and non-caloric artificial sweetener use with markers of glucose and insulin homeostasis and incident diabetes: the Strong Heart Family Study. Eur J Clin Nutr. 2020;74(2):322-327.

Hyson DA. A review and critical analysis of the scientific literature related to 100% fruit juice and human health. Adv Nutr. 2015;6(1):37-51.

Hägele FA, Büsing F, Nas A, et al. High orange juice consumption with or in-between three meals a day differently affects energy balance in healthy subjects. Nutr Diabetes. 2018;8(1):19.

Petroni ML, Brodosi L, Marchignoli F et al. Moderate Alcohol Intake in Non-Alcoholic Fatty Liver Disease: To Drink or Not to Drink? Nutrients. 2019; 11(12):3048.

Polsky S, Akturk HK. Alcohol Consumption, Diabetes Risk, and Cardiovascular Disease Within Diabetes. Curr Diab Rep. 2017;17(12):136.

Kema VH, Mojerla NR, Khan I et al. Effect of alcohol on adipose tissue: a review on ethanol mediated adipose tissue injury. Adipocyte. 2015; 4(4):225-231.

Gill S, Panda S. A smartphone app reveals erratic diurnal eating patterns in humans that can be modulated for health benefits. Cell Metab. 2015;22(5):789-798.

Sutton EF, Beyl R, Early KS et al. Early time-restricted feeding improves insulin sensitivity, blood pressure, and oxidative stress even without weight loss in men with prediabetes. Cell Metab. 2018;27(6):1212-1221.e3.

Cienfuegos S, Gabel K, Kalam F et al. Effects of 4- and 6-h Time-Restricted Feeding on Weight and Cardiometabolic Health: A Randomized Controlled Trial in Adults with Obesity. *Cell Metab.* 2020; S1550-4131(20)30319-3.

Gabel K, Hoddy KK, Haggerty N, et al. Effects of 8-hour time restricted feeding on body weight and metabolic disease risk factors in obese adults: A pilot study. *Nutr Healthy Aging.* 2018;4(4):345-353.

Szymczak-Pajor I, Śliwińska A. Analysis of association between Vitamin D deficiency and insulin resistance. *Nutrients.* 2019;11(4):794.

Rodríguez-Moran M, Guerrero-Romero F. Oral magnesium supplementation improves the metabolic profile of metabolically obese, normal-weight individuals: a randomized double-blind placebo-controlled trial. *Arch Med Res.* 2014;45(5):388-393.

Phillips AJK, Vidafar P, Burns AC et al. High sensitivity and interindividual variability in the response of the human circadian system to evening light. *Proc Natl Acad Sci U S A.* 2019;116(24):12019-12024.

Witbracht MG et al: Dairy food consumption and meal-induced cortisol response interacted to influence weight loss in overweight women undergoing a 12-week, meal-controlled, weight loss intervention. *J Nutr.* 2013 Jan;143(1):46-52.